

Year 12 Extension 1 Test. Fri 4th September, 2009.

Name _____ Teacher _____ Class _____

All questions are 12 marks.

1. **Use a new booklet** Marked by SKB

(a) Two submarines A and B are located at $(0, -50)$ and $(45\sqrt{3}, -50)$ with respect to the origin. The x -axis is the reference sea-level line. Submarine A fires two torpedoes simultaneously with speed 30ms^{-1} at angles θ_1 and θ_2 to the positive direction of the x -axis at submarine B. Neglecting any water or air resistance and letting the acceleration due to gravity, $g = 10\text{ms}^{-2}$:

- (i) Prove that the parametric equations of motion of the torpedoes are given by:
 $x = 30t \cos \theta$ and $y = -5t^2 + 30t \sin \theta - 50$. 2
- (ii) Hence show that the Cartesian equation of motion for either torpedo is given by:
 $y = \frac{-x^2 \sec^2 \theta}{180} + x \tan \theta - 50$ 1
- (iii) Find the exact size of angles θ_1 and θ_2 . 3
- (iv) Determine the maximum heights reached by both torpedoes. 2
- (v) Hence determine the shortest time for a torpedo to strike Submarine B. 1

(b) A projectile is launched from the origin at an angle of 45° to the positive direction of the x -axis. It passes through the points $(12k^2, 8k^2)$ and $(24k^2, 8k^2)$ during its flight. Neglecting air resistance and letting the acceleration due to gravity be $g \text{ ms}^{-2}$, find the speed of projection V in ms^{-1} in terms of g and k . You may assume that the parametric equations of motion are $x = \frac{Vt}{\sqrt{2}}$ and $y = \frac{-gt^2}{2} + \frac{Vt}{\sqrt{2}}$. 3

2. **Use a new booklet** Marked by HRK

- a) In a side show, the probability of getting a prize is 0.4. If a boy tries 8 times what is the probability that he wins at least 3 prizes? 3
- b) Find the coefficient of x^{15} in $\left(x^3 - \frac{2}{x}\right)^9$ 3
- c) Show that if $(1+2x)^4 = \sum_{k=0}^4 {}^4C_k (2x)^k$ and $T_{k+1} = {}^4C_k (2x)^k$ then $\frac{T_{k+1}}{T_k} = \frac{5-k}{k} 2x$
 or show that if $(1+2x)^4 = \sum_{k=0}^4 T_k x^k$ and $T_k = {}^4C_k 2^k$ then $\frac{T_{k+1}}{T_k} = \frac{8-2k}{k+1}$ 3
- d) If $(1+x)^n = \sum_{k=0}^n {}^nC_k x^k$ show that $\sum_{k=1}^n k {}^nC_k = n2^{n-1}$ 3

3. **Use a new booklet** Marked by SKB

- (a) The letters of the statement "FRIDAY FROLICS" are arranged in a circle. Find the probability that the two F's are together. 3
- (b) Consider a pack of 50 playing cards consisting of 5 colours Red, Blue, Green, Yellow and Orange, with cards numbered from 1 to 10 for each colour.
 - (i) Find the number of six card combinations. 1
 - (ii) If six cards are dealt at random from the pack find the number of ways of obtaining four 9's and two eights. 1
 - (i) If six cards are dealt at random from the pack find the number of ways of obtaining four cards of the same number and two cards of a different number. 1
- (c) There are 10 books on a table of which 6 are Maths, 3 are English and 1 is Latin. From these books 3 Maths, 2 English and 1 Latin are chosen and placed on a book shelf. Determine the number of book shelf arrangements if the Maths books are to be together. 2
- (d) Ten molluscs of which two are sea-snails are arranged at random in a row.
 - (i) Determine the number of arrangements without restrictions. 1
 - (ii) Determine the probability that there are at least 5 other molluscs between the sea-snails. 3



Examination Booklet – 4 pages

Student Name or Number

Subject Ext 1 Test Unit Value (please circle) 1 2 3 4

Teacher
4-9-09


Section	Part	Question	Mark

Instructions:

- Clearly identify the questions you have attempted in each booklet
- If you have used more than one booklet for the same question, mark this clearly on each additional booklet you use
- Use the ruled pages only
- No part of this booklet may be torn out
- Under no circumstance may a booklet, used or unused, be removed from the examination by the student
- Marks may be deducted for poor or illegible work



1(a)



(i) $\ddot{x} = 0, \ddot{y} = -10$
 $\therefore \dot{x} = c_1, \dot{y} = -10t + c_2$
 when $t = 0, \dot{x} = 30 \cos \theta, \dot{y} = 30 \sin \theta$
 $\therefore 30 \cos \theta = c_1, 30 \sin \theta = c_2$
 $\therefore \dot{x} = 30 \cos \theta, \dot{y} = -10t + 30 \sin \theta$
 $\therefore x = 30t \cos \theta + c_3, y = -5t^2 + 30t \sin \theta + c_4$ ✓
 when $t = 0, x = 0, y = -50$
 $\therefore c_3 = 0, c_4 = -50$
 $\therefore x = 30t \cos \theta$ — (1)
 $y = -5t^2 + 30t \sin \theta - 50$ — (2) } Are the parametric eqns of motion of the torpedoes. ✓

(ii) From (1) $t = \frac{x}{30 \cos \theta}$ sub into (2)
 $\therefore y = -5 \left(\frac{x}{30 \cos \theta} \right)^2 + 30 \sin \theta \left(\frac{x}{30 \cos \theta} \right) - 50$
 $\therefore y = \frac{-x^2 \sec^2 \theta}{180} + x \tan \theta - 50$ ✓ is the Cartesian eqn of motion.



(iii) Torpedoes pass through $(45\sqrt{3}, -50)$

$$\therefore -50 = \frac{(45\sqrt{3})^2 \sec^2 \theta}{180} + 45\sqrt{3} \tan \theta - 50$$

$$\therefore \frac{(45\sqrt{3})^2 (1 + \tan^2 \theta)}{180} - 45\sqrt{3} \tan \theta = 0 \quad \checkmark$$

$$\therefore \frac{45\sqrt{3}}{180} (1 + \tan^2 \theta) - \tan \theta = 0$$

$$\therefore 45\sqrt{3} \tan^2 \theta - 180 \tan \theta + 45\sqrt{3} = 0$$

$$\therefore \tan \theta = \frac{180 \pm \sqrt{180^2 - 4(45\sqrt{3})(45\sqrt{3})}}{90\sqrt{3}} \quad \checkmark$$

$$= \frac{180 \pm \sqrt{8100}}{90\sqrt{3}}$$

$$= \frac{270}{90\sqrt{3}} \quad \text{or} \quad \frac{90}{90\sqrt{3}}$$

$$= \frac{3}{\sqrt{3}} \quad \text{or} \quad \frac{1}{\sqrt{3}} \quad \checkmark$$

$$\therefore \theta = 60^\circ \quad \text{or} \quad 30^\circ. \quad \Rightarrow \theta_1 = 60^\circ, \theta_2 = 30^\circ.$$

(iv) For max. height $y = 0 \quad \therefore -10t + 30 \sin \theta = 0$

$$\therefore t = 3 \sin \theta$$

$$\text{when } \theta = 60^\circ \quad t = \frac{3\sqrt{3}}{2} \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{sub into (2)}$$

$$\theta = 30^\circ \quad t = \frac{3}{2}$$

$$\text{Now when } t = \frac{3\sqrt{3}}{2} \quad y_{\max} = -5 \left(\frac{3\sqrt{3}}{2} \right)^2 + 30 \left(\frac{3\sqrt{3}}{2} \right) \frac{\sqrt{3}}{2} - 50$$


$$= -16 \frac{1}{2} \text{ m with respect to } O \quad \checkmark$$

$$\text{wh } t = \frac{3}{2} \quad y_{\max} = -5 \left(\frac{3}{2} \right)^2 + 30 \left(\frac{3}{2} \right) \frac{1}{2} - 50$$

$$= -38 \frac{1}{2} \text{ m with respect to } O \quad \checkmark$$

(v) As projection for either torpedo is symmetrical

\Rightarrow shortest time to strike submarine B is 3 seconds.



$$x = \frac{vt}{\frac{1}{\sqrt{2}}} \quad \text{--- (1)}$$

$$y = \frac{-gt^2}{2} + \frac{vt}{\sqrt{2}} \quad \text{--- (2)}$$

$$\text{from (1) } t = \frac{\sqrt{2}x}{v} \quad \text{sub into (2)}$$

$$\therefore y = \frac{-g}{2} \left(\frac{\sqrt{2}x}{v} \right)^2 + \frac{v}{\sqrt{2}} \left(\frac{\sqrt{2}x}{v} \right) \quad \checkmark$$

$$\therefore y = \frac{-gx^2}{v^2} + x$$

$$\text{Through } (12k^2, 8k^2) \quad \therefore 8k^2 = \frac{-g(12k^2)^2}{v^2} + 12k^2$$

$$\therefore g \left(\frac{144k^4}{v^2} \right) = 4k^2 \quad \checkmark$$

$$\therefore v^2 = \frac{g(144k^4)}{4k^2} = 36gk^2$$

$$\Rightarrow v = 6\sqrt{g}k$$

$$\therefore \text{Speed of projection, } v = 6\sqrt{g}k \text{ ms}^{-1}. \quad \checkmark$$

2 (a) $P(\text{prize}) = 0.4 = p$; $P(\overline{\text{prize}}) = 0.6 = q$

Consider $(q+p)^8$

$$\therefore P(\text{win at least 3 prizes}) = 1 - P(\text{win 0, 1, or 2 prizes})$$

$$= 1 - \left[q^8 + {}^8C_1 q^7 p + {}^8C_2 q^6 p^2 \right]$$

$$= 1 - \left[0.6^8 + 8(0.6)^7(0.4) + 28(0.6)^6(0.4)^2 \right]$$

$$= 0.68460544$$

$$= 0.68 \text{ (2dp)}$$



(b)

$$(x^3 - \frac{2}{x})^9$$

Now Term = ${}^9C_r (x^3)^{9-r} (\frac{-2}{x})^r$

$$= {}^9C_r x^{27-3r} (-2)^r x^{-r}$$

For x^{15} term: $27-3r-r=15 \quad \therefore 4r=12, r=3$

\therefore coeff of $x^{15} = {}^9C_3 (-2)^3 = -872$

(c)

$$(1+2x)^4 = \sum_{k=0}^4 {}^4C_k (2x)^k$$

Now Term = ${}^4C_k (2x)^k$

$\therefore T_k = {}^4C_{k-1} (2x)^{k-1}$

$$\frac{T_{k+1}}{T_k} = \frac{{}^4C_k (2x)^k}{{}^4C_{k-1} (2x)^{k-1}}$$

$$= \frac{4! (k-1)! (5-k)! 2x}{(4-k)! k! 4!}$$

$$= \left(\frac{5-k}{k}\right) 2x$$

(d)

$$(1+x)^n = \sum_{k=0}^n {}^nC_k x^k$$

d.b.s. with x :

$$n(1+x)^{n-1} = \sum_{k=0}^{n-1} {}^nC_k k x^{k-1}$$

let $x=1$

$$\therefore n(2)^{n-1} = \sum_{k=1}^n {}^nC_k \cdot k \cdot 1^{k-1}$$

$$\therefore n 2^{n-1} = \sum_{k=1}^n k {}^nC_k$$

(c*)

Alternative: $(1+2x)^4 = \sum_{k=0}^4 T_k x^k$. Now $T_k = {}^4C_k 2^k$

$\therefore T_{k+1} = {}^4C_{k+1} 2^{k+1}$

$$\frac{T_{k+1}}{T_k} = \frac{4! (4-k)! k! 2^{k+1}}{(3-k)! (k+1)! 4! 2^k}$$

$$= \frac{(4-k) 2}{k+1} = \frac{8-2k}{k+1}$$

2 are I's and 7 other single letters

\therefore No. of ways two F's together = 2! This leaves $(13-2)+1 = 12$ letter units

$$\therefore \text{Prob}(2 \text{ F's together in a circle}) = \frac{1 \times 2! \times 11!}{2! \cdot 2! \cdot 2!} = \frac{1 \times 12!}{2! \cdot 2! \cdot 2!} = \frac{1}{6}$$

(b) (i) ${}^{50}C_6 = 15,890,700$

(ii) No. of ways (4 a's and 2 s's) = ${}^5C_4 \times {}^5C_2 = 50$

(ii) No. of ways (any 4 of 1 number and any 2 of 2 different numbers) = $\frac{{}^5C_4 \times {}^5C_2 \times 2! \times {}^{10}C_2}{2! \times 2!} = 4500$

(c) No. of arrangements if 3 Maths books together, Consider 3 Maths books as a unit. This is done in 3! ways. This leaves $(6-3)+1 = 4$ book units.

\therefore No. of arrangements if 3 Maths books together = ${}^6C_3 \times 3! \times {}^3C_2 \times 2! \times 4! = 8640$

(d) (i) Number of arrangements without restrictions = $10! = 3,628,800$

(ii) Possible arrangements are:

$$\left. \begin{array}{l} S \text{ --- } S \text{ ---} \\ - S \text{ --- } S \text{ ---} \\ - - S \text{ --- } S \text{ ---} \\ - - - S \text{ --- } S \text{ ---} \end{array} \right\} \begin{array}{l} 5 \text{ spaces for} \\ \text{other molluscs} \\ = 2! \times 4 \times 8! \end{array}$$

\therefore similarly for 6 spaces: $2! \times 3 \times 8!$
for 7 spaces: $2! \times 2 \times 8!$
for 8 spaces: $2! \times 1 \times 8!$

\therefore Prob (at least 5 other molluscs between sea snails) = $\frac{2! \times 8! \cdot [4+3+2+1]}{10!} = \frac{2}{9}$