


Sydney Girls High School



Mathematics Department

HSC Extension 1 Half-Yearly Examination

2004

Topics Assessed:

- Polynomials*
- Circle Geometry*
- Inverse Trigonometric Functions*
- Integration II*

Time Allowed: 75 minutes

Instructions:

- There are 3 (THREE) questions of equal value.
- Start each question on a new page.

QUESTION 3:

a) For the function $f(x) = 2\sin^{-1}3x$:

- ii. State the domain.
- iii. State the range.
- iv. Sketch the graph of the function.
- v. Find the equation of the tangent to the curve at the point $y = \frac{\pi}{2}$.

b) For the function $y = \frac{1}{x+2}$:

- i. Find the inverse function.
- ii. Find the point(s) of intersection between the function and its inverse.

c) If $f(x) = \sin^{-1}x + \cos^{-1}x$, $-1 \leq x \leq 1$:

- ii. Show that $f'(x) = 0$ for all x .
- iii. Show that $f(x) = \frac{\pi}{2}$ for all x .

d) Find the derivative of $y = \cos^{-1}\sqrt{1-x}$.

e) Write down the general solution for $\sin\theta = \frac{1}{\sqrt{2}}$.

f) Evaluate $\int_0^{\frac{5}{3}} \frac{dx}{25+9x^2}$. Leave your answer in terms of π .

g) Evaluate, showing working, $\sin\left[\cos^{-1}\frac{4}{5} + \tan^{-1}\left(-\frac{4}{3}\right)\right]$.

END OF TEST ☺

Question 1:

$$P(x) = x^3 - 7x - 6$$

$$P(-1) = (-1)^3 - 7(-1) - 6$$

$$= -1 + 7 - 6$$

$$= 0$$


∴ $P(-1) = 0$ ∴ $x = -1$ is a root of $P(x)$. (2)

$$\begin{array}{r} x^2 - x - 6 \\ +1 \overline{) x^3 + 0x^2 - 7x - 6} \\ \underline{x^3 + x^2} \\ -x^2 - 7x \\ \underline{-x^2 - x} \\ -6x - 6 \\ \underline{-6x - 6} \\ 0 \end{array}$$

$$\therefore P(x) = (x+1)(x^2 - x - 6)$$

$$= (x+1)(x-3)(x+2)$$

∴ roots are $-1, 3$ and -2 . (3)



X is midpt of AB (given)

∴ $OX \perp AB$ (line which bisects a chord from the centre of the circle is perp. to the chord)

$$\therefore \angle OXB = 90^\circ$$

Similarly, $\angle OYB = 90^\circ$

$$\angle OXB + \angle OYB = 90^\circ + 90^\circ$$

$$= 180^\circ$$

∴ OXBY is a cyclic quadrilateral (opp. angles are supplementary). (3)

Let $\angle ADC = x$

$$\therefore \angle ABY = 180^\circ - x \text{ (opp. } \angle \text{s cyclic quad ABCD are supp.)}$$

In cyclic quad OXBY

$$\angle XOY = 180^\circ - \angle ABY \text{ (opp } \angle \text{s cyclic quad OXBY are supp.)}$$

$$= 180^\circ - (180^\circ - x)$$

$$= x$$

$$\therefore \angle ADC = \angle XOY. \quad (3)$$

b) $P(x) = x^2 - 4x$

$$P(-3) = (-3)^2 - 4(-3)$$

$$= -27 + 12$$

$$= -15$$

∴ remainder is -15 . (2)

c) Let roots be $\alpha - d, \alpha, \alpha + d$. (5)

$$\alpha - d + \alpha + \alpha + d = \frac{36}{8}$$

$$3\alpha = \frac{36}{8}$$

$$\alpha = 1\frac{1}{2}$$

$$\alpha(\alpha - d)(\alpha + d) = \frac{-21}{8}$$

$$\frac{3}{2} \left(\left\{ \frac{3}{2} \right\}^2 - d^2 \right) = \frac{-21}{8}$$

$$\frac{9}{4} - d^2 = -\frac{7}{4}$$

$$d^2 = 16$$

$$d = \pm 4$$

∴ roots are $-2\frac{1}{2}, 1\frac{1}{2}, 5\frac{1}{2}$

d) $P(x) = x^4 - 12x + 7$

$$P(0) = 0^4 - 12(0) + 7 = 7$$

$$P(1) = 1^4 - 12 + 7 = -4$$

Since $P(0) > 0$ and $P(1) < 0$, the root lies between 0 and 1. (5)

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$= 1 - \left(\frac{-4}{-8} \right)$$

$$= 1 - \frac{1}{2}$$

$$= 0.50 \text{ (2 dec. pl.)}$$

$f(x)$ always

$$f(1) = -4$$

$$f'(x) = 4x^3 - 12$$

$$f'(1) = -8$$

∴ root is 0.50.

ex 2

a) $\int \frac{t}{\sqrt{1+t}} dt$ $u = 1+t$
 $\frac{du}{dt} = 1$
 $dt = du$

$\int \frac{u-1}{\sqrt{u}} du$

$\int (u-1)u^{-\frac{1}{2}} du$

$= \int u^{\frac{1}{2}} - u^{-\frac{1}{2}} du$ (5)
 $= \frac{u^{\frac{3}{2}}}{\frac{3}{2}} - \frac{u^{\frac{1}{2}}}{\frac{1}{2}} + C$


$= \frac{2}{3} \sqrt{(1+t)^3} - 2\sqrt{1+t} + C$

b) $\int_0^4 x \sqrt{16-x^2} dx$ $u = 16-x^2$
 $\frac{du}{dx} = -2x$

$\int_0^4 2x \sqrt{16-x^2} dx$ $du = -2x dx$ (5)
 $x=4 \rightarrow u=0$
 $x=0 \rightarrow u=16$

$\int_0^4 x \sqrt{u} du$

$= \frac{1}{2} \int_0^4 u^{\frac{1}{2}} du = \frac{1}{2} \left[\frac{u^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^4 = \frac{1}{2} \left[\frac{2}{3} (16)^{\frac{3}{2}} \right]$
 $= \frac{64}{3} = 21\frac{1}{3}$



$\angle FAB = 2x$ ext. \angle of a cyclic quad ABCD
 $\angle FAE = x$ ext. \angle of a cyclic quad AEC D

$\angle EAC = 2x - x = x$ (5)

EA bisects $\angle FAB$

$\int_0^5 \sqrt{x+4} dx$ $u = x+4$
 $\frac{du}{dx} = 1$
 $dx = du$

$5 \int (u-4)u^{\frac{1}{2}} du$

$= 5 \int u^{\frac{3}{2}} - 4u^{\frac{1}{2}} du$


$= 5 \left[\frac{u^{\frac{5}{2}}}{\frac{5}{2}} - \frac{4u^{\frac{3}{2}}}{\frac{3}{2}} \right]$ (5)

$= 5 \left[\frac{2\sqrt{(x+4)^5}}{5} - \frac{8\sqrt{(x+4)^3}}{3} \right] + C$

$= 2\sqrt{(x+4)^5} - \frac{40\sqrt{(x+4)^3}}{3} + C$

7 d) Vol = $\pi \int_{\frac{1}{\sqrt{3}}}^{\sqrt{3}} \frac{1}{1+x^2} dx$

$= \pi \left[\tan^{-1} x \right]_{\frac{1}{\sqrt{3}}}^{\sqrt{3}}$



$= \pi \left[\frac{\pi}{3} - \frac{\pi}{6} \right]$ (5)

$= \frac{\pi}{6} \left[\frac{\pi}{6} \right]$

$= \frac{\pi^2}{6}$

QUESTION 3 (25 marks)


(a) $f(x) = 2\sin^{-1} 3x$

(ii) $-1 \leq 3x \leq 1$

D: $-\frac{1}{3} \leq x \leq \frac{1}{3}$ (1)

(iii) $-\frac{\pi}{2} \leq \frac{y}{2} \leq \frac{\pi}{2}$

R: $-\pi \leq y \leq \pi$ (1)



(v) $f'(x) = 2 \left[\frac{1}{\sqrt{1-u^2}} \times 3 \right]$
 $= \frac{6}{\sqrt{1-9x^2}}$
 $= \frac{6}{\sqrt{9-x^2}}$

If $y = \frac{\pi}{2}$ then $\frac{\pi}{2} = 2\sin^{-1} 3x$
 $\frac{\pi}{4} = \sin^{-1} 3x$
 $\frac{1}{3\sqrt{2}} = x$

If $x = \frac{1}{3\sqrt{2}}$ $f'(\frac{1}{3\sqrt{2}}) = \frac{2}{\sqrt{\frac{1}{9} - (\frac{1}{3\sqrt{2}})^2}}$
 $= \frac{2}{\sqrt{\frac{1}{9} - \frac{1}{18}}}$
 $= \frac{2}{\sqrt{\frac{1}{18}}}$
 $= 6\sqrt{2}$

$\therefore m = 6\sqrt{2}$

Using point gradient formula

$y - \frac{\pi}{2} = 6\sqrt{2}(x - \frac{1}{3\sqrt{2}})$

$y = 6\sqrt{2}x - 2 + \frac{\pi}{2}$

$0 = 12\sqrt{2}x - 2y + \pi - 4$ (3)

(b) $y = \frac{1}{x+2}$

(i) $y = \frac{1}{x+2}$
 $x = \frac{1}{y+2}$

$y = \frac{1}{x} - 2$

$f'(x) = \frac{1}{x^2} - 2$ (2)

(ii) $\frac{1}{x+2} = \frac{1}{x} - 2$

$\frac{1}{x+2} = \frac{1-2x}{x}$

$x = (x+2)(1-2x)$

$x = x - 2x^2 + 2 - 4x$

$2x^2 + 4x - 2 = 0$

$x^2 + 2x - 1 = 0$

$x = \frac{-4 \pm \sqrt{32}}{4}$

$= -1 \pm \sqrt{2}$ (3)

\therefore pts. of intersection are $(-1+\sqrt{2}, -1+\sqrt{2})$ & $(-1-\sqrt{2}, -1-\sqrt{2})$

(c) $f(x) = \sin^{-1} x + \cos^{-1} x$

(i) $\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$

$\frac{d}{dx} \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}}$ (2)

$\therefore f'(x) = \frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}} = 0$

3(c)(iii) Show that $f(x) = \frac{\pi}{2}$

for all x .

Method 1: let $a = \sin^{-1} x$


$x = \sin a$ ($-\frac{\pi}{2} \leq a \leq \frac{\pi}{2}$)

$= \cos(\frac{\pi}{2} - a)$

$\frac{\pi}{2} - a = \cos^{-1} x$

$\therefore \sin^{-1} x + \cos^{-1} x = a + \frac{\pi}{2} - a = \frac{\pi}{2}$ (2)

Method 2:



let $\beta = \sin^{-1} x$ & $\alpha = \cos^{-1} x$

$\alpha + \beta = 180^\circ - 90^\circ$ (angle sum of Δ)

$= 90^\circ$

$\therefore \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$

(d) $y = \cos^{-1} \sqrt{1-x}$

let $u = (1-x)^{\frac{1}{2}}$

$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

$= \frac{-1}{\sqrt{1-u^2}} \times \frac{-1}{2\sqrt{1-x}}$

$= \frac{1}{2\sqrt{1-(1-x)}} \times \frac{1}{\sqrt{1-x}}$

$= \frac{1}{2\sqrt{x}(\sqrt{1-x})}$ (2)

$= \frac{1}{2\sqrt{x-x^2}}$

(e) $\sin \theta = \frac{1}{\sqrt{2}}$

$\sin^{-1}(\frac{1}{\sqrt{2}}) = \frac{\pi}{4}$

for general solution

$\theta = n\pi + (-1)^n \sin^{-1} b$ (2)

$= n\pi + (-1)^n \frac{\pi}{4}$

(f) $\int_0^{\frac{\pi}{3}} \frac{dx}{25+9x^2}$

$= \frac{1}{9} \int_0^{\frac{\pi}{3}} \frac{dx}{\frac{25}{9} + x^2}$

$= \frac{1}{9} \cdot \frac{3}{5} \left[\tan^{-1} \frac{3x}{5} \right]_0^{\frac{\pi}{3}}$

$= \frac{1}{15} \left[\tan^{-1} 1 - \tan^{-1} 0 \right]$



$= \frac{\pi}{60}$ (3)

(g) $\sin^{-1} \left[\cos^{-1} \frac{4}{5} + \tan^{-1} \left(-\frac{4}{3} \right) \right]$

let $a = \cos^{-1} \frac{4}{5}$ & $b = \tan^{-1} \left(-\frac{4}{3} \right)$

$\therefore \cos a = \frac{4}{5}$

$\tan b = -\frac{4}{3}$



$\sin(a+b) = \sin a \cos b + \cos a \sin b$

$= \frac{3}{5} \times \frac{3}{5} + \frac{4}{5} \times \frac{-4}{5}$

$= -\frac{7}{25}$

$= \frac{-7}{25}$ (2)