

SYDNEY GIRLS HIGH SCHOOL



2006 HSC Assessment Task 3

June 13, 2006

MATHEMATICS Extension 1

Year 12

Time allowed: 80 minutes

Topics: Parametrics, Circle Geometry, Inverse Functions, Integration by Substitution

DIRECTIONS TO CANDIDATES:

- Attempt all questions
- Questions are of equal value
- There are 4 questions with part marks shown in brackets
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work
- Board approved calculators may be used
- Write on one side of the paper only

QUESTION 1

MARKS

- a) Find the inverse of the following functions and state the domain and range
- (i)  $y = \log_e(x-3)$  2
- (ii)  $y = x^2 - 4x + 5 \quad x \geq 2$  2
- b) Differentiate
- (i)  $y = \sin^{-1} 3x$  2
- (ii)  $y = \cos^{-1} \frac{x}{4}$  2
- c) Find the primitive function of
- (i)  $\int \frac{1}{4+x^2} dx$  1
- d)  $f(x) = x \sin^{-1} x$
- (i) what is the domain of  $f(x)$  1
- (ii) show that this is an even function 2
- (iii) verify that when  $x = 0$ ,  $f(x)$  is stationary 2
- (iv) sketch a graph of  $y = f(x)$  1
- e) (i) Show that  $\frac{d(x^2 \tan^{-1} x)}{dx} = 2x \tan^{-1} x + 1 - \frac{1}{1+x^2}$  2
- (ii) Hence or otherwise find  $\int_0^{\sqrt{5}} x \tan^{-1} x dx$  3


QUESTION 2

- a) Find the following indefinite integrals using the substitution given MARKS
- (i)  $\int x\sqrt{x^2+4} dx$   $u = x^2 + 4$  2
- (ii)  $\int \frac{dx}{x(\log x_e)^3}$   $u = \log x_e$  2
- (iii)  $\int \frac{e^x dx}{\sqrt{49-e^x}}$   $u = e^x$  2
- b) Evaluate the following definite integrals using the substitution given
- (i)  $\int_{-5}^0 \frac{t dt}{\sqrt{4-t}}$   $t = 4 - u^2$  4
- (ii)  $\int_0^{\frac{\pi}{2}} \frac{\sin \theta}{3-2 \cos \theta} d\theta$   $y = 3 - 2 \cos \theta$  4
- c) The region R is bounded by the curve  $y = \frac{x}{x+1}$  the x-axis and the vertical line  $x = 3$ .
- Use the substitution  $u = x + 1$  to find
- (i) the exact area R 3
- (ii) the exact volume generated when R is rotated about the x-axis 3

QUESTION 3

MARKS

- a) T  $(2at, at^2)$  is a variable point on the parabola  $x^2 = 4ay$  3
- (i) show that the gradient of the tangent at T is t.
- (ii) show that the equation of the tangent at T is  $y = tx - at^2$
- b) Write down the equation of the chord of contact of tangents drawn from a point  $(x_1, y_1)$  to the parabola  $x^2 = 4ay$  4
- (i) find the equation of the chord of contact from the point  $(3, -2)$  to the parabola  $x^2 = 8y$
- (ii) at what point does the line intersect the directrix
- c) If PM is a normal to the parabola  $x^2 = 4ay$  at a variable point  $P(2ap, ap^2)$  and FN is drawn through the focus F parallel to the tangent at P to cut the normal at N 4
- (i) prove that the locus of N(x, y) is  $x^2 = a(y-a)$



d) The tangents at  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  on the parabola  $x^2 = 4ay$  intersect at  $T(a(p+q), apq)$

9


(i) find  $M$  the midpoint of  $PQ$

Hence show that

(ii)  $TM$  is parallel to the axis of symmetry


(iii) if  $TM$  meets the parabola on  $R$ , then  $R$  bisects  $TM$

(iv) the tangent at  $R$  is parallel to the chord  $PQ$



QUESTION 4


MARKS



O is the centre of the circle 5  
Prove that


(i)  $x + y = 90$

(ii)  $z - y = 2x$



3

BA and BC are tangents to the circles  
 $DE = 5 \times BD$ . Prove  $BA = \sqrt{6} \times BD$




5

$PQ$  is a common tangent and  $PA \parallel QB$ . Prove that

(i)  $PC \parallel QD$

(ii)  $PQBC$  is a cyclic quadrilateral


d)



P and Q are the centres of the circles  
PAX and QAY are straight lines.

Prove that P, Q, X and Y are concyclic

e)



A and B are fixed points. P moves on the plane so that AB subtends an angle of  $30^\circ$  at P.

(i) describe the locus of P

(ii) describe what construction you would carry out to draw the locus of P

THE END

4

4/12 Ext 2, June 06

a) if  $y = \log(x-3)$

$f^{-1}: x = \log(y-3)$


$y-3 = e^x$

$y = e^x + 3$

Domain = the set of reals

Range =  $\{y: y > 3\}$

$y = e^x + 3$



d) i)  $f(x) = x \cdot \sin^{-1} x$

Domain =  $\{x: -1 \leq x \leq 1\}$  (1)

ii)  $f(a) = a \cdot \sin^{-1} a$

$f(-a) = -a \cdot \sin^{-1}(-a)$

$= -a \cdot (-\sin^{-1} a)$

$= a \cdot \sin^{-1} a$  (2)

$\therefore f(-a) = f(a)$

$\therefore f(x)$  is an even function

iii)  $f(x) = x \cdot \sin^{-1} x$

$f'(x) = \sqrt{1-x^2} + x \cdot \frac{-x}{\sqrt{1-x^2}}$

$= \sin^{-1} x + \frac{x^2}{\sqrt{1-x^2}}$

$= \sin^{-1} 0 + \frac{0}{\sqrt{1-0}}$

$f'(0) = 0 + 0$

$= 0 + 0$

$\therefore f'(0) = 0$

$\therefore$  When  $x=0$ ,  $(x, f(x))$  is stationary.

ii)  $y = x^2 - 4x + 5, x \geq 2$

$f^{-1}: x = y^2 - 4y + 5, y \geq 2$

$x-5 = y^2 - 4y$

$x-5 = (y-2)^2 - 4$

$x-1 = (y-2)^2, y \geq 2$

$y-2 = \sqrt{x-1}, y \geq 2$

$y = 2 + \sqrt{x-1}, y \geq 2$

Domain =  $\{x: x \geq 1\}$

Range =  $\{y: y \geq 2\}$  (2)

i)  $y = \sin^{-1} 3x$

$y' = \frac{1}{\sqrt{1-(3x)^2}} \times 3$


$= \frac{3}{\sqrt{1-9x^2}}$  (2)

ii)  $y = \cos^{-1} x$

$y' = \frac{-1}{\sqrt{1-x^2}}$  (2)

i)  $\int \frac{1}{4+2x} dx$

$= \frac{1}{2} \tan^{-1} \frac{x}{2} + C$  (1)



e) i)  $\frac{d}{dx} (x^2 \cdot \tan^{-1} x) = \sqrt{\frac{du}{dx} + u \frac{dv}{dx}}$

$= \tan^{-1} x \times 2x + x^2 \times \frac{1}{1+x^2}$

$\frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + \frac{x^2}{x^2+1}$

$= 2x \cdot \tan^{-1} x + \frac{x^2+1-1}{x^2+1}$

$\frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + \frac{x^2}{x^2+1} - \frac{1}{x^2+1}$

$\frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + 1 - \frac{1}{x^2+1}$

(2)

i)  $\frac{d}{dx}(x^2 \tan^{-1} x) = 2x \cdot \tan^{-1} x + 1 - \frac{1}{x^2+1}$

$x^2 \cdot \tan^{-1} x = \int 2x \cdot \tan^{-1} x dx + \int dx - \int \frac{dx}{x^2+1}$   
 $x^2 \cdot \tan^{-1} x = \int 2x \cdot \tan^{-1} x dx + (x - \tan^{-1} x + c)$   
 $2 \cdot \tan^{-1} x - x + \tan^{-1} x + c = \int 2x \cdot \tan^{-1} x dx$

$2x \cdot \tan^{-1} x dx = [x^2 \cdot \tan^{-1} x - x + \tan^{-1} x + c]$   
 $x \cdot \tan^{-1} x dx = [\frac{x^2}{2} \cdot \tan^{-1} x - \frac{x}{2} + \frac{1}{2} \tan^{-1} x + c]$   
 $= (\frac{3}{2} \cdot \tan^{-1} 3 - \frac{\sqrt{3}}{2} + \frac{1}{2} \tan^{-1} 3 + c) - (0 - 0 + 0 + c)$   
 $= \frac{3}{2} \times \frac{\pi}{3} - \frac{\sqrt{3}}{2} + \frac{1}{2} \times \frac{\pi}{3}$   
 $= \frac{4\pi}{6} - \frac{\sqrt{3}}{2}$   
 $= \frac{2\pi}{3} - \frac{\sqrt{3}}{2}$  (3)

ii)  $\int x \sqrt{x^2+4} dx$   $u = x^2+4$   
 $= \int \sqrt{u} \cdot \frac{1}{2} du$   
 $= \frac{1}{2} \int u^{1/2} du$   
 $= \frac{1}{2} \times \frac{2}{3} u^{3/2} + c$   
 $= \frac{1}{3} (x^2+4)^{3/2} + c$  (2)

ii)  $\int \frac{dx}{x(\log_2 x)^2}$   $u = \log_2 x$   
 $du = \frac{1}{x} dx$   
 $= \int \frac{du}{u^2}$   
 $= \int u^{-2} du$   
 $= -\frac{1}{u} + c$   
 $= -\frac{1}{\log_2 x} + c$  (2)


iii)  $\int \frac{e^x dx}{\sqrt{49-e^x}}$   $u = e^x$   
 $du = e^x dx$   
 $= \int \frac{du}{\sqrt{49-u}}$   
 $= \int (49-u)^{-1/2} du$   
 $= 2(49-u)^{1/2} + c$   
 $= 2\sqrt{49-e^x} + c$  (2)

b) i)  $\int \frac{t dt}{-5\sqrt{4-t}}$   $t = 4-u$   
 $dt = -2u du$   
 $= \int \frac{(4-u)(-2u) du}{\sqrt{4-u}}$   $u=2$   
 $= \int \frac{-2u(4-u)}{\sqrt{4-u}} du$   $u=-3$   
 $= \int \frac{-2u(4-u)}{\sqrt{4-u}} du$   $u=3$

$= \int_2^3 \frac{2u(4-u)}{\sqrt{4-u}} du$   
 $= \int_2^3 (8-2u) du$   
 $= [8u - \frac{2}{3}u^3]_2^3$   
 $= (8 \times 3 - \frac{2}{3} \times 3^3) - (8 \times 2 - \frac{2}{3} \times 2^3)$   
 $= (24 - 18) - (16 - \frac{16}{3})$   
 $= 6 - 16 + \frac{16}{3}$   
 $= -10 + \frac{16}{3}$   
 $= -4\frac{2}{3}$  (4)

ii)  $\int_0^{\pi/2} \frac{\sin \theta d\theta}{3-2\cos \theta}$   $y = 3-2\cos \theta$   
 $dy = 2\sin \theta d\theta$   
 $\theta = \frac{\pi}{3}, y = 3-2\cos \frac{\pi}{3} = 3-1 = 2$   
 $\theta = 0, y = 3-2\cos 0 = 3-2 = 1$

$= \int_1^2 \frac{dy}{2y}$   
 $= \frac{1}{2} \ln y + c$   
 $= (\frac{1}{2} \ln 2 + c) - (\frac{1}{2} \ln 1 + c)$   
 $= \frac{1}{2} \ln 2$  (4)




$A = \int_0^3 \frac{x}{x+1} dx$   $u = x+1$   
 $du = dx$   
 $= \int_1^4 \frac{u-1}{u} du$   
 $= \int_1^4 (1 - \frac{1}{u}) du$   
 $= [u - \ln u + c]_1^4$   
 $= (4 - \ln 4 + c) - (1 - \ln 1 + c)$   
 $= 4 - \ln 4 - 1 + \ln 1$   
 $= (3 - \ln 4)$  sq units (3)

$V = \pi \int_0^3 y^2 dx$   
 $= \pi \int_0^3 \frac{x^2}{(x+1)^2} dx$   
 $= \pi \int_1^4 \frac{(u-1)^2}{u^2} du$

$= \pi \int_1^4 \frac{u^2 - 2u + 1}{u^2} du$   
 $= \pi \int_1^4 (1 - \frac{2}{u} + \frac{1}{u^2}) du$   
 $= \pi [u - 2 \ln u - \frac{1}{u}]_1^4$   
 $= \pi [(4 - 2 \ln 4 - \frac{1}{4}) - (1 - \ln 1 - 1)]$

$= \pi (3\frac{3}{4} - 2 \ln 4 + 2 \ln 1)$   
 $= \pi (15/4 - 2 \ln 4)$   
 or units (3)



$$y = \frac{x^2}{4a}$$

$$\frac{dy}{dx} = \frac{2x}{4a} = \frac{x}{2a}$$


at  $x = 2at$ ,  $\frac{dy}{dx} = \frac{2at}{2a} = t$

$\therefore T$  is  $Y - Y_1 = m(X - X_1)$

$$Y - at^2 = t(X - 2at)$$

$$Y - at^2 = tX - 2at^2$$

$$Y = tX - at^2$$



$x_1 x_2 = 2a(y_1 + y_2)$

$x_1 = 3, y_1 = -2, a = 2$

$\therefore 3x_2 = 4(y_2 - 2)$

$$3x_2 = 4y_2 - 8$$


$$3x_2 - 4y_2 + 8 = 0$$

at  $y_2 = -2$ ,  $3x_2 + 8 + 8 = 0$

$$3x_2 = -16$$

$$x_2 = -16/3$$

$\therefore P$  is  $(-16/3, -2)$



$x^2 = 4ay$

$$2x = 4a \cdot \frac{dy}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{x}{2a}$$

at  $x = 2ap$ ,  $\frac{dy}{dx} = p$

$\therefore m_{n} = -1/p$

Eqn is  $Y - ap^2 = -1/p(X - 2ap)$

$$pY - ap^3 = -X + 2ap$$

$$\therefore X + pY = 2ap + ap^3$$

line through focus is  $Y - a = p(X - 0)$

$$Y = pX + a$$

Sub into normal.

$$\therefore X + p(pX + a) = 2ap + ap^3$$

$$X + p^2X = ap + ap^3$$

$$X(1 + p^2) = ap(1 + p^2)$$


$$\therefore X = ap \text{ \& } Y = ap^2 + a$$

$\therefore \frac{X}{a} = p \Rightarrow Y = a(\frac{X}{a})^2 + a$

$$Y = \frac{X^2}{a} + a$$

$$aY = X^2 + a^2$$

$$\therefore X^2 = a(Y - a)$$



i)  $M$  is  $\frac{ap + 2aq}{2}, \frac{ap^2 + aq^2}{2}$

$$= a(p+q), \frac{a}{2}(p^2 + q^2)$$

ii)  $MT$  have the same  $X$  value

$\therefore$  line is vertical (i.e. parallel to  $Y$  axis)

iii) Midpoint  $MT$  is  $a(p+q), \frac{1}{2} \{ \frac{a}{2}(p^2 + q^2) + a \}$

$$= a(p+q), \frac{1}{2} \{ \frac{a}{2}(p^2 + q^2) + 2ap \}$$

$$= a(p+q), \frac{a}{4}(p+q)^2$$

Sub into  $X^2 = 4ay$

$$LHS = a^2(p+q)^2$$

$$RHS = 4a \cdot \frac{a}{4}(p+q)^2 = a^2(p+q)^2$$

$\therefore R$  lies on  $X^2 = 4ay$ .

iv)  $\frac{dy}{dx} = \frac{x}{2a}$ . At  $R$ ,  $\frac{dy}{dx} = \frac{1}{2} \cdot \frac{2a(p+q)}{2} = \frac{p+q}{2}$

$$m_{PQ} = \frac{ap^2 - aq^2}{2ap - 2aq} = \frac{a(p-q)(p+q)}{2a(p-q)} = \frac{p+q}{2} = \text{slope}$$

$\therefore P, X, Q$  in a cyclic quad because  $\angle PXR$  &  $\angle PYQ$  are subtending equal angles.

$\angle PXR = \angle PYQ$


$\therefore \angle PXR = \angle PYQ$  (radii)

$\therefore \angle XPR = \angle QPY$  (radii)

$\therefore \angle XPR = \angle QPY$  (radii)

$\therefore \angle PXR = \angle PYQ$

$\therefore P, X, Q$  in a cyclic quad



$\therefore PCBD$  in a cyclic quad

$\therefore \angle PCB = 180 - X$  (op. angles)


$\therefore \angle ACP = X$

$\therefore \angle ACP = \angle BDE$  (radii)

$\therefore \angle ACP = \angle BDE$  (radii)

$\therefore \angle ACP = \angle BDE$

$\therefore \angle ACP = \angle BDE$



$\therefore$  With compass on  $P$  draw a circle of radius  $OP$ .

The construction lines meet at  $A$  &  $B$  & the centre of the circle is where  $\therefore$  construct  $60^\circ$  angles at  $A$  &  $B$ .

If the angle at  $P$  on the circumference is  $30^\circ$ , the angle at the centre is  $60^\circ$ .

