The frequency is the number of times each score, event or measurement occurs. A frequency distribution is shown in a table in which information or data is arranged in order. From the frequency distribution, the average and range can be found, and the data can be shown on graphs such as histograms, frequency curves and cumulative frequency graphs.

1	Definitions	Averages	<u>Histogram</u>	Frequency Curve
	Stem and Leaf Diagram	Cumulative Frequency Curve	Measures of Spread	Box and Whisker Graph

#### **Definitions**

- · Discrete data can be counted.
  - e.g. Number of people in a theatre, number of leaves on a tree.
- Continuous data is obtained by measuring.
  - e.g. Heights of people, lengths of leaves on a tree.
- Ungrouped data is used with small amounts of data where individual scores are listed.
  - e.g. Test results of five people: 15, 20, 18, 17, 16
- Grouped data is used with larger amounts of data where scores are placed in groups or classes.
  - e.g.Test results of 30 people

Score	Frequency
0 - 5	3
6 - 10	8
11 - 15	12
16 - 20	7

**A** 107

### Averages

An average is a number that typifies a set of numbers or statistics.

There are three commonly used types of average.

- Median. The median of a set of scores is the middle score when the scores have been arranged in order from smallest to largest.
- Mode. The mode is the most frequent score, that is, the one that occurs most often.
- · Mean.

The mean of a set of scores is the sum of all of the scores divided by the number of scores.

The mean is often known as the average.

The symbol for the mean is  $\bar{x}$ 

For ungrouped data,  $\overline{x} = \frac{\text{sum of scores}}{\text{number of scores}}$ For grouped data,  $\overline{x} = \frac{\text{sum of (middle score x frequency)}}{\text{number of scores}}$ 

Example	Answer
For the following set of test results of 10 people:	(a) The mean $= \underbrace{3+8+5+6+8+5+4+1+10+5}_{10}$
3, 8, 5, 6, 8, 5, 4, 1, 10, 5	<u>= 55</u> = 5.5 10
Find:	(b) The mode = 5 (occurs 3
(a) The mean	times)
(b) The mode	(c) Arrange in order:
(c) The median	1, 3, 4, 5, 5, 5, 6, 8, 8, 10
(c) The median	Median is 5

#### A 100

## Findng the mean on a calculator

The procedure for finding statistical values will vary slightly from calculator to calculator.

For a scientific calculator such as the Texas Instruments TI-30 IIB, to find the mean of 5, 7, and 12:

Task	Keys	Select
Select the statistical mode STAT	27H SID (=)	Choose 1-VAR
Enter the data	for each item of data	$x_1 = \dots$ and FRQ = 1 for each item of data.
Check number of items entered	atenues $n=3$	Select n
	FROMEN T	₹

a mu mean	**************************************	DOIGOL A
Clear old data	EXIT STAT	Always do this before entering a new set of data.

### Finding the mean on a spreadsheet

Enter the data:

	A	£
	Values	
	5	***************************************
	7	
	12	
S500	Average	8

The function entered in cell B5 to find the mean is =AVERAGE(A2..A4) this gives a mean of

#### Histogram

A histogram is a graph that is used to show the information from a frequency distribution.

It is similar to a column graph and the area of each column is proportional to the frequency of the score that it represents.

In a histogram the bars always touch, whereas in a bar graph the bars do not need to touch.

e.g Ages of workers in a company.

Note that this is grouped data.


Ages	Frequency
15 < 25	14
. 25 < 35	20
35 < 45	24
45 < 55	18
55 < 65	10

Note: 15 < 25 means 15 ≤ height < 25

i.e Greater or equal to 15 but less

than 25.

Different people use different ways of presenting this type of data.




## **Frequency Curve**

This is sometimes called a frequency polygon or line graph.

A frequency curve is obtained by joining up the mid-points of the tops of the columns of the histogram.

e.g. For the ages of the workers.



A 707

## Stem and Leaf Diagram

A stem and leaf diagram or chart provides a means of ordering and displaying data.

Numbers are split into categories of the first digit(s) and the last digits(s)

e.g. Given the following test scores:

Let the stem be the first digit (the tens column). Now enter each of the second digits (the units column) into the leaf part of the diagram.

Both parts of a stem and leaf diagram can contain more than one digit.

e.g. For numbers such as 3476, the stem could be 34 and the leaf could be 76.

A 102


#### **Cumulative Frequency Curve**

In a cumulative frequency table, each of the scores from the frequency distribution is added to the total of the ones before it.

e.g. For the ages of the workers:

Age (x)	Cumulative frequency
< 15	0
< 25	14 .
< 35	34
< 45	58
< 55	76
< 65	86

e.g. There are a total of 58 workers aged less than 45 years old.



& 10°

## Measures of Spread

As well as averages, it is necessary to have a measure of how spread out the scores are.

The range and quartiles are ways of measuring this spread.

#### Range


The range of a set of scores is the difference between the highest and the lowest scores.

e.g. For 
$$\{4, 5, 6, 7, 8, 9, 10\}$$
 the range is  $10 - 4 = 6$ 

#### Quartiles

The quartiles divide a distribution into four equal parts.

• For ungrouped data {4, 5, 6, 7, 8, 9, 10}:



- The interquartile range is the difference between the upper and the lower quartiles
  - i.e. Interquartile range = 9 5 = 4
- For grouped data the quartiles and median can be obtained by using the vertical axis of the cumulative frequency curve. The total number of scores is divided by four and then read off using the scale.



## **Box and Whisker Graphs**


Box and whisker graphs are useful for comparing the data in different frequency distributions.

The diagrams consist of a rectangle which indicates the quartiles and the median, and a line on each end to show the maximum and minimum values of the distribution.

e.g. Draw a box and whisker diagram to compare the following frequency distributions of two schools' School Certificate Mathematics results.

School A: Median, 51; Lower quartile, 42; Upper quartile, 65; Maximum, 95; Minimum, 23

School B: Median, 54; Lower quartile, 39; Upper quartile, 68; Maximum, 99; Minimum, 15



### **Box and Whisker Plots**

Box and whisker graphs are useful for comparing the data in different frequency distributions.


The diagrams consist of a rectangle which indicates the quartiles and the median, and a line on each end to show the maximum and minimum values of the distribution.

#### Example

Draw a box and whisker diagram to compare the following frequency distributions of two schools' School Certificate Mathematics results.

School A: Median, 51; Lower quartile, 42; Upper quartile, 65; Maximum, 95; Minimum, 23

School B: Median, 54; Lower quartile, 39; Upper quartile, 68; Maximum, 99; Minimum, 15



From the diagram above, the two sets of marks can now be compared.

School B would appear to have the higher marks with a higher median, upper quartile and maximum.

School A's results are less spread, with an interquartile range of 23, compared to School B's 29.

### **Comparing Data**

This <u>experiment</u> compares two sets of data created by you. The results are shown on histograms, a box plots and in tables.

(The table shows one or two features which are outside the scope of this course.)

AIV

u†

From the diagram above, the two sets of marks can now be compared.

School B would appear to have the higher marks with a higher median, upper quartile and maximum.


School A's results are less spread, with an interquartile range of 23, compared to School B's 29.



- 1. For the following scores, find:
- (a) The median
- (b) The mode
- (c) The mean
- (i) 8, 6, 4, 9, 3, 3, 2
- (ii) 101, 99, 100, 98, 101, 104, 96, 101
- (iii) 10, 12, 12, 12, 20
- (iv) 50, 30, 20, 20, 10, 5
- 2. A golfer plays seven rounds of golf in a week. His scores are: 70, 62, 62, 69, 71, 72, 80
- (a) Calculate the mean, median and mode of these seven scores.
- (b) Which of these averages would not be very representative of his week's golf.



- 3. A girl had a newspaper delivery round. Her weekly pay for 2 month's work was:
- \$12.50, \$11.80, \$10.70, \$13.10, \$4.50, \$12.00, \$9.20, \$9.60
- (a) What was her mean weekly pay?
- (b) What was her median pay?
- (c) What was the difference between her highest and lowest pays?
- 4. The diagram shows the height of plants, measured in centimetres.



- (a) How many plants were measured?
  - (b) What was the median plant height?
  - (c) What percentage of plants were over 45 cm high?
  - (d) What percentage of plants were less than 30 cm high?
  - (e) How many plants were between 50 and 60 cm high?
  - 5. The following set of marks were for a class of 20 pupils. The test was out of 10 marks.


- (a) Construct a frequency distribution table.
- (b) Draw a frequency histogram.
- (c) Draw a frequency curve.
- (d) Calculate the mean mark.
- (e) Find the median mark.
- (f) What is the mode?
- (g) What is the probability that a person picked at random would score
  - (i) Seven? (ii) Less than 5?
- 6. The frequency distribution for the length, in seconds, of 100 telephone calls was:

Time (seconds)	Frequency
0 - 20	0
21 - 40	5
41 - 60	7
61 - 80	14
81 - 100	28

101 - 120	21
121 - 140	13
141 - 160	9
161 - 180	3

- (a) Construct a cumulative frequency table.
- (b) Draw a cumulative frequency graph.
- (c) What number of calls lasted no more than 2 minutes?
- (d) What is the probability that a call picked at random lasted more than 140 seconds?
- (e) What percentage of calls lasted more than 1 minute?
- 7. State whether the following would be discrete or continuous data:
- (a) The weights of people.
- (b) The number of people at a concert.
- (c) The heights of trees.
- (d) A person's shoe size.
- 8. The following list is the temperatures, in °C, at 20 main centres of New Zealand for a day in January.
- 23, 27, 19, 28, 24, 26, 27, 24, 18, 23, 15, 24, 23, 22, 24, 29, 16, 24, 22, 25
- (a) Construct a frequency table with class intervals of 3°C.
- (b) Draw a histogram to illustrate this distribution.
- (c) What is the modal temperature range?
- (d) Calculate the mean temperature.
- (e) What would the median temperature be?
- (f) What is the probability that a centre picked at random would have a temperature greater than or equal to 21°C?

- 9. The cumulative frequency graph shows the volume of drink in cans of soft drink.
- (a) How many cans of soft drink were measured?
- (b) How many cans contained:
- (i) Less than 350 mL?
- (ii) More than 340 mL?
- (iii) Between 320 and 370 mL?
- (c) What is the probability that a randomly selected can of drink will contain less than 350 mL?
- (d) If 4000 cans were produced, how many would you expect to contain less than 350 mL?
- (e) What is the median amount of drink in a can?



10. The table shows the lengths of 17 New Zealand rivers.

Show this information on an ordered stem and leaf diagram, using the first two digits as the stem.

Length (in km)
169
217
161
209
209

vanditaiki	173
Manawatu	182
Waihou	175
Mohaka	172
Oreti	203
,Wairau	169
Rakaia	145
Whangehu	161
Patea	143
Mokau	158
Ngarurora	154
Buller	177

11. The final examination results of two classes are to be compared.

The results are shown below:

First class	53	43	67	87	36	-56	76	45	87	73
	· 45	53	74	65	38	71	40	33	59	
Second class	46	45	47	67	56	59	66	50	,79	80
	35	55	77	55	30	65	42	39	39	

- (a) Draw a stem and leaf diagram for each class.
- (b) Arrange each set of results in order and calculate the median and the upper and lower quartiles.
- (c) Draw box and whisker plots, side by side and compare the results of the two classes.



1.

- (i) (a) 4
- (b) 3
- (c) 5

- (ii) (a) 100.5
- (b) 101
- (c) 100

- (iii) (a) 12
- (b) 12
- (c) 13.2

- (iv) (a) 20
- (b) 20
- (c) 22.5
- 2. (a) Mean = 69.4 (to 1 d.p.) Median = 70 Mode = 62
- (b) The mode

3.

- (a) \$10.43
- (b) \$11.25
- (c) \$8.60
- 4. (a) 40 (b) 40 cm (c)  $37 \pm 2\%$  (d)  $21 \pm 2\%$  (e)  $7 \pm 1$

5.(a)

Score (x)	Frequency (f)	x.f
0	1	0
1	1	1
2	2	4
3	2	6
4	2	8
5 .	3	15
6	5	30
7	3	21
8	1	8
9	0	0
10	0	0
Totals	20	93

(b) Histogram of test results




(c) Frequency curve of test results.

Trequency \$ 4 4 2 2 1 0 0 1 2 3 4 5 6 7 8 9 10

- (d) 4.65 (e) 5 (f) 6
- (g) (i) 0.15 (ii) 0.4
- 6. (a)


Time (t) (seconds)	Cumulative frequency
≤ 20	0
≤ 40	5
≤ 60	12
≤ 80	26 .
≤ 100	54
≤ 120	75
≤ 140	88
≤ 160	97
≤ 180	100



- (c) 75 (d) 0.12 (e) 88%
- 7. (a) Continuous (b) Discrete
- (c) Continuous (d) Discrete
- 8. (a)

Temp <sup>0</sup> C	Frequency	Midpoint	f. x
15 - 17	2	16	32
18 - 20	2	19	38
21 - 23	5	22	110
24 - 26	7 .	25	175
27 - 29	4	28	112
	20		467

(b) Histogram of temperatures



(c) 24 - 26°C (d) 23.35°C (e) 24°C (f) 0.8

- (c) 0.7
- (d) 2800
- (e) 340 mL

10.	14 15 16 17 18 19 20 21	35 48 1199 257 2 3 399	Stem represents: hundreds and tens e.g. 17 represents 170
-----	--	--	---

11. (a)	First class	1 1	Second class
	863	3	0599
	5530	4	2567
	9633	5	05569
	75	6	567
	6431	7	79
	77	8	0
			Stem represents: tens
			e.g. 6 represents 60


(b) There are 19 results in each class, therefore the 10th score is the median, the 5th score is the lower quartile and the 15th score is the upper quartile.

### From diagram:

Γ	median = 56	•	median = 55	
First class:	upper quartile =	Second class:	upper quartile = 66	
Ciass.	lower quartile = 43		lower quartile = 42	

(c)

First class Second class



Medians are similar, the first class has a wider spread and range of marks although the second class has the lowest mark.



#### Unit Test #49

Select your answers to the following 10 questions from the pop-up menus in the right hand column. When you are satisfied with your answers, fill in your name in the space provided below the test, and click the "Submit Test" button. Clicking the "Begin Test Again" button will clear all the answers.

Q1:	Questions 1 to 4 refer to the following set of data: 2, 7, 7, 8, 9, 12, 15, 20  What is the mode?	A. 8.5 B. 7 C. 10 D. 18	Answer 1:	
Q2:	What is the mean?	A. 8.5 B. 7 C. 10 D. 18	Answer 2:	
Q3:	What is the median?	A 8.5 B. 7 C. 10 D. 18	Answer 3:	
Q4:	What is the range?	A. 8.5 B. 7 C. 10 D. 18	Answer 4:	
Q5:	The heights of 100 people are measured.  What type of data is this?	A. continuous B. discrete	Answer 5:	
Q6:	In one sample of 100 pods of peas, the mean was 6 peas per pod. In another sample of 50 pods, the mean was 8 peas per pod. The combined mean for the two samples taken together was	A. exactly 6 peas per pod B.between 6 and 7 peas per pod C. exactly 7 peas per pod D. between 7 and 8 peas per pod	Answer 6:	
Q7:	A group of students sat a test. Their mean score was 11. Another student who scored 29 joins the group. The mean is now 13. The number of students in the	A. 5 B. 8 C. 11 D. 18	Answer 7:	

Q8:	Original group was  Score 70 71 72 73 74  Frequency 1 3 5 4 7  David's best 20 golf scores are given in the table above.	A. 72 B. 72.65 C. 73 D. 74	Answer 8:	
Q9:	The median is  In the table given above, the mean is	A. 72 B. 72.65 C. 73 D. 74	Answer 9:	
Q10:	In the table above what is the mode score?	A. 71 B. 72 C. 73 D. 74	Answer 10:	\$ 35533

Enter your initial	
and surname here:	

Submit Test Begin Test Again

ANS

Notes | Test | Summary | FAQ | Exercise | Answers

## **Frequency Distributions**

Unit Test #49

Select your answers to the following 10 questions from the pop-up menus in the right hand column. When you are satisfied with your answers, fill in your name in the space provided below the test, and click the "Submit Test" button. Clicking the "Begin Test Again" button will clear all the answers.

Espere Wong

Q1:	Questions 1 to 4 refer to the following set of data: 2, 7, 7, 8, 9, 12, 15, 29  What is the mode?	A. 8.5 B. 7 C. 10 D. 18	Answer 1:	
Q2:	What is the mean?	A. 8.5 B. 7 C.10 D. 18	Answer 2:	
Q3;	What is the median?	A 8.5 √ B. 7 C. 10 D. 18	Answer 3:	
Q4:	What is the range?	A. 8.5 B. 7 C. 10 D 18 V	Answer 4:	
Q5;	The heights of 100 people are measured.  What type of data is this?	A)continuous  B. discrete	Answer 5:	
Q6:	In one sample of 100 pods of peas, the mean was 6 peas per pod. In another sample of 50 pods, the mean was 8 peas per pod. The combined mean for the two samples taken together was	A. exactly 6 peas per pod B.between 6 and 7 peas per pod C. exactly 7 peas per pod D. between 7 and 8 peas per pod	Answer 6:	
Q7:	A group of students sat a test. Their mean score was 11. Another student who scored 29 joins the group. The mean is now 13. The number of students in the	A. 5 B. 8 C. 11 D. 18	Answer 7:	

	original group was			
Q8:	Score 70 71 72 73 74 Frequency 1 3 5 4 7  David's best 20 golf scores are given in the table above. The median is	A. 72 B. 72.65 Ĉ 73 √ D. 74	Answer 8:	
Q9:	In the table given above, the mean is	A. 72 B) 72.65 C. 73 D. 74	Answer 9:	
Q10:	In the table above what is the mode score?	A. 71 B. 72 C. 73 D)74	Answer 10:	

Enter your initial	
and surname here.	

Submit Test

Begin Test Again