



**SYDNEY BOYS HIGH
SCHOOL**
MOORE PARK, SURRY HILLS

2009
YEAR 12 Mathematics Extension 1
HSC Task #2

Mathematics Extension 1

General Instructions

- Reading Time – 5 Minutes
- Working time – 90 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators may be used.
- Each Section is to be returned in a separate bundle.
- All answers must be given in exact simplified form unless otherwise stated.
- Marks may NOT be awarded for messy or badly arranged work.
- All necessary working should be shown in every question.

Total Marks – 61

- Attempt questions 1-3
- Start each new section of a separate answer booklet

Examiner: *D. McQuillan*

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, a > 0, -a < x < a$$


$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, x > 0$

Section A (21 marks)

- (1) O is the centre of the circle. Find the value of θ .



- (2) Find

(a) $\int 3x^5 dx$

(b) $\int \frac{(3x+2)^2}{3} dx$

(c) $\int \frac{dx}{\sqrt{7x-1}}$

- (3) Evaluate

(a) $\int_{-8}^8 \frac{x^3}{2} dx$

(b) $\int_3^9 (x^2 + 3) dx$

(c) $\int \sqrt{4-x} dx$

- (4) The new NSW number plates have three letters then two numbers followed by a letter.



How many number plates with this configuration are possible?

Marks

1


5

5

1

- (5) AB is a tangent. Find the value of x .

2



- (6) Given the parametric equations $x = 2t - 1$ and $y = 4t^2$ find $\frac{dy}{dx}$.


2

- (7) Find the volume when $y = x^2 - 3$ is rotated about the y-axis between $y = -3$ and $y = 2$.

2

- (8) ABCD is a trapezium in which $AB \parallel DC$. AD and BC are the diameters of their respective circles, and these circles cut DC at X and Y respectively. Prove that ABYX is a rectangle.

3



End of Section A

Start a new ANSWER BOOKLET

Section B (20 marks)

Marks

- (1) Use Simpson's Rule with 3 function values to find an approximation to the area under the arc of $y = \log_{10} x$ between $x = 2$ and $x = 4$. Round answer to 2 decimal places. 2
- (2) For the function $f(x) = x^3 - 3x + 1$. 5
- (a) Show that $f(x)$ has a zero for x between 0 and 1.
- (b) Take $x = 0$ as your first approximation and use two applications of Newton's method to find a better approximation to 3 decimal places.
- (c) Explain why you could not use $x = 1$ as your first approximation.
- (3) A five-card hand is dealt from a standard 52-card deck. How many ways can you be dealt, 3
- (a) three hearts and two clubs.
- (b) three of one suit and two of another.
- (4) Find the area enclosed between $y = x + 2$ and $y = x^2 - 2x + 2$. 3
- (5) If ${}^n P_k = 120 {}^n C_k$, find the value of k . 2
- (6) Given that the equation for the chord of contact is $xx_0 = 2a(y + y_0)$. Find the point of intersection of the tangents to $x^2 = 8y$ at the points where the chord of contact $y = x - 1$ intersects the parabola. 2

(7) For the parabola $x^2 = 4ay$.

3

- (a) Derive the equation of the chord joining the points $P = (2ap, ap^2)$ and $Q = (2aq, aq^2)$.
- (b) Show that as q approaches p the equation of the chord becomes the equation of the tangent.


End of Section B

Start a new ANSWER BOOKLET

Section C (20 marks)

Marks

- (1) Prove that ${}^n P_k = {}^{n-1} P_k + k \cdot {}^{n-1} P_{k-1}$. 2
- (2) AB is a chord of a circle and PA is a tangent at A. XY is a perpendicular bisector of AB and passes through the centre of the circle, O. PY is perpendicular to PA. 6
- (a) Copy the diagram into your answer booklet.
- (b) Show that triangle AYB is an isosceles triangle.
- (c) If $\angle PAY = \alpha$ show that $\angle YAB = \alpha$.
- (d) Prove that $PY = XY$.



- (3) In how many ways can the letters of the word TOMATO be arranged if the Ts are to be separated? 2
- (4) A Rugby League team of 13 players are to be selected at random from 20 boys. What is the probability that of 3 friends, 2 are selected and one misses out. 2

- (5) For the parabola $x = 4t, y = 2t^2$. 4
- (a) Derive the equations of the normals at $t = p$ and $t = -p$.
- (b) The normals intersect at R. Find the locus of R.

- (6) Use mathematical induction to prove that 4

$$\frac{1}{2} + \frac{2}{3} + \dots + \frac{n}{n+1} < \frac{n^2}{n+1}$$

for all integers $n \geq 2$.

End of Section C

End of Exam

1/ $\theta = 55^\circ$ 1

2/ (a) $\int 3x^5 dx = \frac{x^6}{2} + c$ 1

(b) $\int \frac{(3x+2)^2}{8} dx = \frac{(3x+2)^3}{27} + c$ 2

or $= x^3 + 2x^2 + \frac{4}{3}x + c$

(c) $\int \frac{dx}{(7x-1)^4} = \frac{2}{7} \sqrt{7x-1} + c$ 2

3/ (a) $\int_{-8}^8 \frac{x^3}{2} dx = \left[\frac{x^4}{8} \right]_{-8}^8 = 10$ 2

[Could use $\int_{-a}^a f(x) dx = 0$ if $f(x)$ odd]

(b) $\int_{-3}^3 (x^2+3) dx = \left[\frac{x^3}{3} + 3x \right]_{-3}^3 = 36$ 2

$= [(9+9) - (-9-9)]$

[Could use $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ if $f(x)$ even]

(c) $\int \sqrt{x-4} dx = \frac{2}{3} (x-4)^{3/2} + c$ 1

4/ $26.26.26.10.10.26 = 4569760$ 1

5/ $(AB)^2 = 2x \times x$ 1

$100 = 2x^2$

$x = \sqrt{50} = 5\sqrt{2}$ 1

6/ $\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dx}{dt}$

$= 8t \div 2$ 2

$= 4t$

$= 4 \left(\frac{x+1}{2} \right)$

$= 2x + 2$

7/ $V = \pi \int_{-3}^2 (y+3) dy$

$= \pi \left[\frac{y^2}{2} + 3y \right]_{-3}^2$ 1

$= \pi [(2+6) - (4.5-9)]$

$= 12.5\pi \text{ units}^3$ 1

8/ 3

$\hat{A}XD = 90^\circ$ (angle in semi-c)
 $\hat{A}XY = 90^\circ$ (adj. angles on str. line)

Similarly


$\hat{B}YC = 90^\circ$ (angle in semi-c)
 $\hat{B}YX = 90^\circ$ (adj. ang. on str. line)

$\Rightarrow AX \parallel BY$ since the co-interior angles AXY and BYX are supplementary.

$\therefore ABYX$ is a // gram with one angle a right angle.

ie a rectangle.

Section B



$\int_a^b f(x) dx \approx \frac{b-a}{6} [f(a) + 4f(\frac{a+b}{2}) + f(b)]$

x	2	3	4
f(x)	$\log_{10} 2$	$\log_{10} 3$	$\log_{10} 4$

$\int_2^4 \log_{10} x dx \approx \frac{4-2}{6} [\log_{10} 2 + 4 \log_{10} 3 + \log_{10} 4]$

$\approx 0.94 \text{ units}^2$

2) a) $f(x) = x^3 - 3x + 1$

$f(0) = 1$

$f(1) = -1$

since $f(x)$ is continuous and opposite in sign at endpoints, $f(x)$ has at least one zero when $0 \leq x \leq 1$.

b) $a_1 = a - \frac{f(a)}{f'(a)}$

$f(x) = x^3 - 3x + 1$

$f(0) = 1$

$f'(x) = 3x^2 - 3$

$f'(0) = -3$

$a_1 = 0 - \frac{1}{-3}$

$a_1 = \frac{1}{3}$

$f(\frac{1}{3}) = \frac{1}{27}$

$a_2 = \frac{1}{3} - \frac{(\frac{1}{27})}{(-\frac{8}{3})}$

$f'(\frac{1}{3}) = -\frac{8}{3}$


$a_2 = \frac{25}{72}$

$a_2 \approx 0.347$

c) $f'(1) = 0$ tangent when $x=1$ is horizontal
 doesn't cross the x -axis.
 \therefore Newton's method fails.


3) a) ${}^{13}C_3 \times {}^{13}C_2 = 22308$

b) 3 cards 2 cards



$12 \times 22308 = 267696$

4) $y = x^2 - 2x + 2$ — ① $y = x + 2$ — ②
 $y = x^2 - 2x + 1 + 1$
 $y = (x-1)^2 + 1$



sub ① into ②
 $x^2 - 2x + 2 = x + 2$
 $x^2 - 3x = 0$
 $x(x-3) = 0$
 $x = 0$ $x = 3$

Area = $\int_0^3 [x+2 - (x^2 - 2x + 2)] dx$
 $= \int_0^3 (3x - x^2) dx$
 $= \left[\frac{3x^2}{2} - \frac{x^3}{3} \right]_0^3$

$= \frac{3(3)^2}{2} - \frac{(3)^3}{3} - (0)$

$= \frac{9}{2} \text{ units}^2$

5) ${}^n P_k = 120 {}^n C_k$

$\frac{n!}{(n-k)!} = 120 \frac{n!}{(n-k)! k!}$

$k! = 120$

$\therefore k = 5$

6) $x^2 = 8y$
 $x^2 = 4ay$
 $a = 2$

$xx_1 = 2a(y+y_1)$

$xx_1 = 4(y+y_1)$


$y = x - 1$

$x = y + 1$

$4x = 4(y+1)$

$x_1 = 4, y_1 = 1$

∴ point of intersection of the tangents is (4, 1)



$m_{PQ} = \frac{ap^2 - aq^2}{2ap - 2aq}$
 $= \frac{a(p^2 - q^2)}{2a(p - q)}$
 $= \frac{(p - q)(p + q)}{2(p - q)}$
 $= \frac{p + q}{2}$

$y - y_1 = m(x - x_1)$
 $y - ap^2 = \frac{p+q}{2}(x - 2ap)$

$$y - ap^{\frac{1}{2}} = \frac{1}{2}(p+q)x - ap^{\frac{1}{2}} - apq$$

$$y - \frac{1}{2}(p+q)x + apq = 0$$

b) as $q \rightarrow p$

$$y - \frac{1}{2}(p+p)x + ap(p) = 0$$


$$y - px + ap^2 = 0$$

which is the equation of the tangent.

SECTION C. (EXT. 1.)

Q1

$$\begin{aligned} \text{RHS} &= {}^{n-1}P_k + k {}^{n-1}P_{k-1} \\ &= \frac{(n-1)!}{(n-1-k)!} + k \frac{(n-1)!}{(n-k)!} \\ &= \frac{(n-1)!}{(n-k)!} [n-k+k] \\ &= \frac{n(n-1)!}{(n-k)!} \\ &= \frac{n!}{(n-k)!} \\ &= {}^n P_r \\ &= \text{LHS.} \end{aligned}$$



(b) $AX = BX$ (data)
 $\widehat{AXY} = \widehat{BYX}$ (data)
 XY is common.
 $\therefore \triangle AXY \cong \triangle BYX$ (S.A.S)
 $\therefore AY = BY$ (corresponding sides of congruent triangles)
 $\therefore \triangle AYB$ is isosceles.

(c) $\angle PAY = d$.

$\therefore \angle ABY = d$ (angle between tangent and chord is equal to the angle in the alternate segment)

Now $\angle YAB = \angle ABY$ (isosceles triangle, base angles equal).

$\therefore \angle YAB = d$.

(d) APYX is a cyclic quadrilateral (opposite angles \widehat{APY} and \widehat{AXY} are supplementary)
 $\therefore \widehat{XY} = \widehat{PY}$ (equal angles are subtended by equal chords, at the circumference of a circle).

[OR. Show that $\triangle AXY \cong \triangle APY$ using (AAS) test]

Q3. $\frac{6!}{2! \times 2!} - \frac{5!}{2!} = 120$ OR. $\frac{4!}{2!} \times \binom{5}{2} = 120$.

Q4. $\frac{\binom{3}{2} \times \binom{17}{11}}{\binom{20}{13}} = \frac{91}{190}$.

Q5. Given $y = 2t^2$ & $x = 4t$.
 $\frac{dy}{dt} = 4t$, $\frac{dx}{dt} = 4$, $\therefore \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = 4t \times \frac{1}{4} = t$.

\therefore slope of normal where $t = p$ is $-\frac{1}{p}$.

\therefore eqn. of normal is $\frac{y - 2p^2}{x - 4p} = -\frac{1}{p}$.

$$py - 2p^3 = -x + 4p \quad \text{--- (1)}$$

$$x + py = 2p^3 + 4p$$

At $t = -p$, $\frac{y - 2p^2}{x + 4p} = \frac{1}{p}$ $\therefore py - 2p^3 = x + 4p$ --- (2)
 $-x + py = 2p^3 + 4p$ --- (2)

Add (1) + (2)

$$2py = 4p^3 + 8p$$


$$y = 2p^2 + 4$$

Subtract (2) from (1)

$$2x = 0$$

$$x = 0$$

$\therefore R$ is $(0, 2p^2 + 4)$



Q6. Step I. Consider $n = 2$.

$$LHS = \frac{1}{2} + \frac{2}{3} = \frac{7}{6} \quad RHS = \frac{4}{3}$$

\therefore true when $n = 2$ as $\frac{7}{6} < \frac{4}{3}$

Step II Assume the statement to be true when $n = k$.

$$\text{i.e. } \frac{1}{2} + \frac{2}{3} + \dots + \frac{k}{k+1} < \frac{k^2}{k+1}$$

Step III Required to prove the statement is true

for $n = k+1$ under the assumption in Step II.

$$\text{i.e. } \frac{1}{2} + \frac{2}{3} + \dots + \frac{k}{k+1} + \frac{k+1}{k+2} < \frac{(k+1)^2}{k+2}$$

±

now

$$\begin{aligned}
 \text{LHS} &= \frac{1}{2} + \frac{2}{3} + \dots + \frac{k}{k+1} + \frac{k+1}{k+2} \\
 &< \frac{k^2}{k+1} + \frac{k+1}{k+2} \quad (\text{from assumption}) \\
 &= \frac{k^2(k+2) + (k+1)^2}{(k+1)(k+2)} \\
 &= \frac{k^3 + 2k^2 + k^2 + 2k + 1}{(k+1)(k+2)} \\
 &< \frac{k^3 + 3k^2 + 3k + 1}{(k+1)(k+2)} \\
 &= \frac{(k+1)^3}{(k+1)(k+2)} \\
 &= \frac{(k+1)^2}{k+2} = \text{RHS}
 \end{aligned}$$

$$\therefore \dots \frac{1}{2} + \frac{2}{3} + \dots + \frac{k+1}{k+2} < \frac{(k+1)^2}{k+2}$$

Step IV. By the principle of mathematical induction the statement is true for all integral $n \geq 2$.