

BOARD OF STUDIES
NEW SOUTH WALES

2010
HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

Total marks – 120

- Attempt Questions 1–10
- All questions are of equal value

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a}\sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a}\cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a}\tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a}\sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a}\tan^{-1}\frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}\frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln\left(x + \sqrt{x^2 - a^2}\right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln\left(x + \sqrt{x^2 + a^2}\right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Total marks – 120

Attempt Questions 1–10

All questions are of equal value

Answer each question in the appropriate writing booklet. Extra writing booklets are available.

Question 1 (12 marks) Use the Question 1 Writing Booklet.

(a) Solve $x^2 = 4x$. 2

(b) Find integers a and b such that $\frac{1}{\sqrt{5}-2} = a + b\sqrt{5}$. 2

(c) Write down the equation of the circle with centre $(-1, 2)$ and radius 5. 1

(d) Solve $|2x + 3| = 9$. 2

(e) Differentiate $x^2 \tan x$ with respect to x . 2

(f) Find the limiting sum of the geometric series $1 - \frac{1}{3} + \frac{1}{9} - \frac{1}{27} + \dots$. 2

(g) Let $f(x) = \sqrt{x-8}$. What is the domain of $f(x)$? 1

Question 2 (12 marks) Use the Question 2 Writing Booklet.

(a) Differentiate $\frac{\cos x}{x}$ with respect to x . 2

(b) Solve the inequality $x^2 - x - 12 < 0$. 2

(c) Find the gradient of the tangent to the curve $y = \ln(3x)$ at the point where $x = 2$. 2


(d) (i) Find $\int \sqrt{5x+1} \, dx$. 2

(ii) Find $\int \frac{x}{4+x^2} \, dx$. 2

(e) Given that $\int_0^6 (x+k) \, dx = 30$, and k is a constant, find the value of k . 2

Question 3 (12 marks) Use the Question 3 Writing Booklet.

- (a) In the diagram A , B and C are the points $(-2, -4)$, $(12, 6)$ and $(6, 8)$ respectively. The point $N(2, 2)$ is the midpoint of AC . The point M is the midpoint of AB .



- | | |
|--|---|
| (i) Find the coordinates of M . | 1 |
| (ii) Find the gradient of BC . | 1 |
| (iii) Prove that $\triangle ABC$ is similar to $\triangle AMN$. | 2 |
| (iv) Find the equation of MN . | 2 |
| (v) Find the exact length of BC . | 1 |
| (vi) Given that the area of $\triangle ABC$ is 44 square units, find the perpendicular distance from A to BC . | 1 |

Question 3 continues on page 5

Question 3 (continued)

- (b) (i) Sketch the curve $y = \ln x$. 1

- (ii) Use the trapezoidal rule with three function values to find an approximation to 2


$$\int_1^3 \ln x \, dx.$$

- (iii) State whether the approximation found in part (ii) is greater than or less than the exact value of $\int_1^3 \ln x \, dx$. Justify your answer. 1

End of Question 3

Question 4 (12 marks) Use the Question 4 Writing Booklet.

- (a) Susannah is training for a fun run by running every week for 26 weeks. She runs 1 km in the first week and each week after that she runs 750 m more than the previous week, until she reaches 10 km in a week. She then continues to run 10 km each week.
- (i) How far does Susannah run in the 9th week? **1**
- (ii) In which week does she first run 10 km? **1**
- (iii) What is the total distance that Susannah runs in 26 weeks? **2**
- (b) The curves $y = e^{2x}$ and $y = e^{-x}$ intersect at the point $(0, 1)$ as shown in the diagram. **3**



Find the exact area enclosed by the curves and the line $x = 2$.

Question 4 continues on page 7


Question 4 (continued)

- (c) There are twelve chocolates in a box. Four of the chocolates have mint centres, four have caramel centres and four have strawberry centres. Ali randomly selects two chocolates and eats them.
- (i) What is the probability that the two chocolates have mint centres? **1**
- (ii) What is the probability that the two chocolates have the same centre? **1**
- (iii) What is the probability that the two chocolates have different centres? **1**
- (d) Let $f(x) = 1 + e^x$. **2**
- Show that $f(x) \times f(-x) = f(x) + f(-x)$.

End of Question 4

Question 5 (12 marks) Use the Question 5 Writing Booklet.

- (a) A rainwater tank is to be designed in the shape of a cylinder with radius r metres and height h metres.



The volume of the tank is to be 10 cubic metres. Let A be the surface area of the tank, including its top and base, in square metres.

- (i) Given that $A = 2\pi r^2 + 2\pi r h$, show that $A = 2\pi r^2 + \frac{20}{r}$. 2
- (ii) Show that A has a minimum value and find the value of r for which the minimum occurs. 3

- (b) (i) Prove that 1

$$\sec^2 x + \sec x \tan x = \frac{1 + \sin x}{\cos^2 x}$$

- (ii) Hence prove that 1

$$\sec^2 x + \sec x \tan x = \frac{1}{1 - \sin x}$$

- (iii) Hence use the table of standard integrals to find the exact value of 2


$$\int_0^{\frac{\pi}{4}} \frac{1}{1 - \sin x} dx.$$

Question 5 continues on page 9

Question 5 (continued)

- (c) The diagram shows the curve $y = \frac{1}{x}$, for $x > 0$. 3

The area under the curve between $x = a$ and $x = 1$ is A_1 . The area under the curve between $x = 1$ and $x = b$ is A_2 .



The areas A_1 and A_2 are each equal to 1 square unit.

Find the values of a and b .

End of Question 5


Question 6 (12 marks) Use the Question 6 Writing Booklet.

(a) Let $f(x) = (x + 2)(x^2 + 4)$.

- (i) Show that the graph $y = f(x)$ has no stationary points. 2
- (ii) Find the values of x for which the graph $y = f(x)$ is concave down, and the values for which it is concave up. 2
- (iii) Sketch the graph $y = f(x)$, indicating the values of the x and y intercepts. 2

(b) The diagram shows a circle with centre O and radius 5 cm.

The length of the arc PQ is 9 cm. Lines drawn perpendicular to OP and OQ at P and Q respectively meet at T .



NOT TO SCALE

- (i) Find $\angle POQ$ in radians. 1
- (ii) Prove that $\triangle OPT$ is congruent to $\triangle OQT$. 2
- (iii) Find the length of PT . 1
- (iv) Find the area of the shaded region. 2

Question 7 (12 marks) Use the Question 7 Writing Booklet.

(a) The acceleration of a particle is given by


$$\ddot{x} = 4 \cos 2t$$

where x is displacement in metres and t is time in seconds.

Initially the particle is at the origin with a velocity of 1 m s^{-1} .

- (i) Show that the velocity of the particle is given by 2
- $$\dot{x} = 2 \sin 2t + 1.$$
- (ii) Find the time when the particle first comes to rest. 2
- (iii) Find the displacement, x , of the particle in terms of t . 2

(b) The parabola shown in the diagram is the graph $y = x^2$. The points $A(-1, 1)$ and $B(2, 4)$ are on the parabola.



- (i) Find the equation of the tangent to the parabola at A . 2
- (ii) Let M be the midpoint of AB .
There is a point C on the parabola such that the tangent at C is parallel to AB .
Show that the line MC is vertical. 2
- (iii) The tangent at A meets the line MC at T . 2
Show that the line BT is a tangent to the parabola.

Question 8 (12 marks) Use the Question 8 Writing Booklet.

- (a) Assume that the population, P , of cane toads in Australia has been growing at a rate proportional to P . That is, $\frac{dP}{dt} = kP$ where k is a positive constant. 4

There were 102 cane toads brought to Australia from Hawaii in 1935.

Seventy-five years later, in 2010, it is estimated that there are 200 million cane toads in Australia.

If the population continues to grow at this rate, how many cane toads will there be in Australia in 2035?


- (b) Two identical biased coins are tossed together, and the outcome is recorded. After a large number of trials it is observed that the probability that both coins land showing heads is 0.36. 2

What is the probability that both coins land showing tails?

Question 8 continues on page 13

Question 8 (continued)

- (c) The graph shown is $y = A \sin bx$.



- (i) Write down the value of A . 1
- (ii) Find the value of b . 1
- (iii) Copy or trace the graph into your writing booklet. 2

On the same set of axes, draw the graph $y = 3 \sin x + 1$, for $0 \leq x \leq \pi$.

- (d) Let $f(x) = x^3 - 3x^2 + kx + 8$, where k is a constant. 2

Find the values of k for which $f(x)$ is an increasing function.

End of Question 8

Question 9 (12 marks) Use the Question 9 Writing Booklet.

- (a) (i) When Chris started a new job, \$500 was deposited into his superannuation fund at the beginning of each month. The money was invested at 0.5% per month, compounded monthly. 2

Let \$ P be the value of the investment after 240 months, when Chris retires.

Show that $P = 232\,175.55$.

- (ii) After retirement, Chris withdraws \$2000 from the account at the end of each month, without making any further deposits. The account continues to earn interest at 0.5% per month.

Let \$ A_n be the amount left in the account n months after Chris's retirement.

(1) Show that $A_n = (P - 400\,000) \times 1.005^n + 400\,000$. 3


- (2) For how many months after retirement will there be money left in the account? 2

Question 9 continues on page 15

Question 9 (continued)

- (b) Let $y = f(x)$ be a function defined for $0 \leq x \leq 6$, with $f(0) = 0$.

The diagram shows the graph of the derivative of f , $y = f'(x)$.




The shaded region A_1 has area 4 square units. The shaded region A_2 has area 4 square units.

- (i) For which values of x is $f(x)$ increasing? 1
- (ii) What is the maximum value of $f(x)$? 1
- (iii) Find the value of $f(6)$. 1
- (iv) Draw a graph of $y = f(x)$ for $0 \leq x \leq 6$. 2

End of Question 9

Question 10 (12 marks) Use the Question 10 Writing Booklet.

- (a) In the diagram ABC is an isosceles triangle with $AC = BC = x$. The point D on the interval AB is chosen so that $AD = CD$. Let $AD = a$, $DB = y$ and $\angle ADC = \theta$.



(i) Show that $\triangle ABC$ is similar to $\triangle ACD$.

2

(ii) Show that $x^2 = a^2 + ay$.

1

(iii) Show that $y = a(1 - 2 \cos \theta)$.

2


(iv) Deduce that $y \leq 3a$.

1

Question 10 continues on page 17

Question 10 (continued)

- (b) The circle $x^2 + y^2 = r^2$ has radius r and centre O . The circle meets the positive x -axis at B . The point A is on the interval OB . A vertical line through A meets the circle at P . Let $\theta = \angle OPA$.




- (i) The shaded region bounded by the arc PB and the intervals AB and AP is rotated about the x -axis. Show that the volume, V , formed is given by

3

$$V = \frac{\pi r^3}{3} (2 - 3 \sin \theta + \sin^3 \theta).$$

(ii)



A container is in the shape of a hemisphere of radius r metres. The container is initially horizontal and full of water. The container is then tilted at an angle of θ to the horizontal so that some water spills out.

- (1) Find θ so that the depth of water remaining is one half of the original depth.

1

- (2) What fraction of the original volume is left in the container?

2

End of paper

2010 Higher School Certificate Solutions Mathematics

Question 1

(a) $x^2 = 4x$
 $x(x-4) = 0$
 $x = 0, 4.$

(b) $\frac{1}{\sqrt{5}-2} = \frac{1}{\sqrt{5}-2} \times \frac{\sqrt{5}+2}{\sqrt{5}+2}$
 $= \frac{\sqrt{5}+2}{5-4}$
 $= \sqrt{5}+2$
 $= 2+\sqrt{5}$
 $= a+b\sqrt{5}$
 $\therefore a=2$ and $b=1.$

(c) Circle whose centre is $(-1, 2)$ and radius 5 has equation
 $(x+1)^2 + (y-2)^2 = 25.$

(d) $|2x+3|=9$
 $2x+3=9$ or $2x+3=-9$
 $2x=6$ $2x=-12$
 $x=3$ $x=-6$
 $\therefore x=3, -6$

(e) Using the product rule
 $\frac{d}{dx} x^2 \tan x = u'v + uv'$
 $= 2x \tan x + x^2 \sec^2 x$
 $= x(2 \tan x + x \sec^2 x)$

(f) $1 - \frac{1}{3} + \frac{1}{9} - \frac{1}{27} + \dots$ is a geometric sequence where $a=1$ and $r=-\frac{1}{3}$

$$S_{\infty} = \frac{a}{1-r}$$

$$= \frac{1}{1-(-\frac{1}{3})}$$

$$= \frac{1}{(\frac{4}{3})}$$


$$= \frac{3}{4}$$

(g) $f(x) = \sqrt{x-8}$ is defined for $x-8 \geq 0$
 ie $x \geq 8$
 Hence the domain is $x \geq 8.$

Question 2

(a) Using the quotient rule
 $\frac{d}{dx} \left(\frac{\cos x}{x} \right) = \frac{u'v - uv'}{v^2}$
 $= \frac{(-\sin x) \cdot x - (\cos x) \cdot 1}{x^2}$
 $= \frac{-x \sin x - \cos x}{x^2}$

(b) $x^2 - x - 12 < 0$
 $(x-4)(x+3) < 0$
 $-3 < x < 4$



(c) $\frac{dy}{dx} = \frac{1}{3x} \times 3$
 $= \frac{3x}{3}$
 $= \frac{1}{x}$
 when $x=2$
 $\frac{dy}{dx} = \frac{1}{2}$

The gradient of the tangent is $\frac{1}{2}$

(d) (i) $\int \sqrt{5x+1} dx = \int (5x+1)^{\frac{1}{2}} dx$
 $= \frac{(5x+1)^{\frac{3}{2}}}{\frac{3}{2} \times 5} + C$
 $= \frac{2}{15} \sqrt{(5x+1)^3} + C$

(ii) $\int \frac{x}{4+x^2} dx = \frac{1}{2} \int \frac{x}{4+x^2} dx$
 $= \frac{1}{2} \ln(4+x^2) + C$
 $= \ln \sqrt{4+x^2} + C$

(e) $\int_0^6 (x+k) dx = 30$
 $\left[\frac{1}{2}x^2 + kx \right]_0^6 = 30$
 $\frac{1}{2}(6)^2 + 6k - (0+0) = 30$
 $18 + 6k = 30$
 $6k = 12$
 $k = 2$

Question 3

(a) (i) $M = \left(\frac{-2+12}{2}, \frac{-4+6}{2} \right)$
 $= (5, 1)$
 \therefore The coordinates of M are $(5, 1).$

(ii) $m = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{8-6}{6-12}$
 $= \frac{2}{-6}$
 $= -\frac{1}{3}$
 $\therefore m_{BC} = -\frac{1}{3}$

(iii) Prove $\triangle ABC$ is similar to $\triangle AMN.$
 $N(2,2)$ $M(5,1)$

Gradient of $MN = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{2-1}{2-5}$
 $= -\frac{1}{3}$

$\therefore m_{BC} = m_{MN} \therefore BC$ is parallel to MN

In $\triangle ABC$ and $\triangle AMN$
 $\angle CAB$ is common
 $\angle ANM = \angle ACB$ (corres. \angle s in \parallel)
 $\angle AMN = \angle ABC$ (corres. \angle s in \parallel)
 $\therefore \triangle ABC \sim \triangle AMN$ (equi-angular)

(iv) gradient of $MN = -\frac{1}{3}$ [from (iii)]
 $y - y_1 = m(x - x_1)$
 $y - 2 = -\frac{1}{3}(x - 2)$
 $3y - 6 = -x + 2$
 $x + 3y - 8 = 0$

$$\begin{aligned}
 (v) \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(12 - 6)^2 + (6 - 8)^2} \\
 &= \sqrt{6^2 + (-2)^2} \\
 &= \sqrt{36 + 4} \\
 &= \sqrt{40} \\
 &= 2\sqrt{10}
 \end{aligned}$$

∴ the distance is $2\sqrt{10}$ units

$$(vi) \quad \text{Area of } \triangle ABC = \frac{1}{2} \times BC \times \perp \text{ distance}$$

$$\frac{1}{2} \times 2\sqrt{10} \times h = 44$$

$$\sqrt{10} \times h = 44$$


$$h = \frac{44}{\sqrt{10}} \times \frac{\sqrt{10}}{\sqrt{10}}$$

$$= \frac{44\sqrt{10}}{10}$$

$$= \frac{22\sqrt{10}}{5}$$


∴ the perpendicular distance is $\frac{22\sqrt{10}}{5}$ units.

(b) (i)



$$\begin{aligned}
 (ii) \quad \text{Area} &= \frac{1}{2} [(\ln 1 + \ln 2) + (\ln 2 + \ln 3)] \\
 &= \frac{1}{2} [0 + 2\ln 2 + \ln 3] \\
 &= 1.24 \text{ units}^2 \quad (2 \text{ dp})
 \end{aligned}$$

(iii)



It is less than the exact value because $y = \ln x$ is concave down and the trapezium are beneath the curve.

Question 4

(a) (i) This is an arithmetic series where $a = 1$, $d = 0.75$ and $n = 9$.

$$T_n = a + (n-1)d$$

$$T_9 = 1 + (9-1) \times 0.75$$

$$T_9 = 7$$

∴ She runs 7 km in the 9th week.

(ii) when $T_n = 10$,

$$10 = a + (n-1)d$$

$$10 = 1 + (n-1) \times 0.75$$

$$10 = 1 + 0.75 \times n - 0.75$$

$$0.75 \times n = 9.75$$

$$n = 13$$

∴ She runs 10 km in the 13th week.

(iii) For the first 13 weeks:

$$S_n = \frac{n}{2}(a+l)$$

$$= \frac{13}{2}(1+10)$$

$$= 71.5 \text{ km}$$

For the second 13 weeks, she does 10 km each week.

∴ total distance for 26 weeks is

$$71.5 + 10 \times 13 = 201.5 \text{ km.}$$

(b)

$$\begin{aligned}
 \text{Area} &= \int_0^2 (e^{2x} - e^{-x}) dx \\
 &= \left[\frac{e^{2x}}{2} + e^{-x} \right]_0^2 \\
 &= \left(\frac{e^4}{2} + e^{-2} \right) - \left(\frac{e^0}{2} + e^0 \right) \\
 &= \frac{e^4}{2} + \frac{1}{e^2} - \frac{1}{2} - 1 \\
 &= \frac{e^4}{2} + \frac{1}{e^2} - 1 \frac{1}{2} \text{ units}^2
 \end{aligned}$$

(c)

$$\begin{aligned}
 (i) \quad P(2 \text{ mints}) &= \frac{4}{12} \times \frac{3}{11} \\
 &= \frac{1}{11}
 \end{aligned}$$

$$(ii) \quad P(\text{same}) = P(mm) + P(cc) + P(ss)$$

$$= \frac{1}{11} + \frac{1}{11} + \frac{1}{11}$$

$$= \frac{3}{11}$$

$$(iii) \quad P(\text{different}) = 1 - P(\text{same})$$

$$= 1 - \frac{3}{11}$$

$$= \frac{8}{11}$$

(d)

$$f(x) \times f(-x) = (1 + e^x)(1 + e^{-x})$$

$$= 1 + e^x + e^{-x} + e^0$$

$$= 2 + e^x + e^{-x}$$

$$\text{(since } e^0 = 1)$$

$$f(x) + f(-x) = (1 + e^x) + (1 + e^{-x})$$

$$= 1 + e^x + 1 + e^{-x}$$

$$= 2 + e^x + e^{-x}$$

$$\therefore f(x) \times f(-x) = f(x) + f(-x)$$

Question 5

$$(a) (i) \quad V = Ah$$

$$V = \pi r^2 h$$

$$10 = \pi \times r^2 \times h$$

$$h = \frac{10}{\pi r^2}$$

$$A = 2\pi r^2 + 2\pi r h$$

$$= 2\pi r^2 + 2\pi r \left(\frac{10}{\pi r^2} \right)$$

$$= 2\pi r^2 + \frac{20}{r}$$

$$(ii) \quad A = 2\pi r^2 + 20r^{-1}$$

$$\frac{dA}{dr} = 4\pi r - 20r^{-2}$$

$$\text{let } \frac{dA}{dr} = 0$$

$$0 = 4\pi r - \frac{20}{r^2}$$

$$4\pi r = \frac{20}{r^2}$$

$$\pi r^3 = 5$$

$$r = \sqrt[3]{\frac{5}{\pi}}$$

$$\frac{d^2A}{dr^2} = 4\pi + \frac{40}{r^3}$$

Since $r > 0$ for all r

$$\frac{d^2A}{dr^2} > 0 \quad \therefore \text{a minimum value}$$

∴ the minimum value is $\sqrt[3]{\frac{5}{\pi}}$ m.

(b) (i) $LHS = \sec^2 x + \sec x \tan x$
 $= \frac{1}{\cos^2 x} + \frac{1}{\cos x} \times \frac{\sin x}{\cos x}$
 $= \frac{1}{\cos^2 x} + \frac{\sin x}{\cos^2 x}$
 $= \frac{1 + \sin x}{\cos^2 x}$
 $= RHS$

(ii) $LHS = \sec^2 x + \sec x \tan x$
 $= \frac{1 + \sin x}{\cos^2 x}$ from part (i)
 $= \frac{1 + \sin x}{1 - \sin^2 x}$
 $= \frac{1 + \sin x}{(1 + \sin x)(1 - \sin x)}$
 $= \frac{1}{1 - \sin x}$
 $= RHS$
 $\therefore \sec^2 x + \sec x \tan x = \frac{1}{1 - \sin x}$

(iii) $\int_0^{\frac{\pi}{4}} \frac{1}{1 - \sin x} dx$
 $= \int_0^{\frac{\pi}{4}} \sec^2 x + \sec x \tan x dx$
 $= [\tan x + \sec x]_0^{\frac{\pi}{4}}$
 $= \left[\left(\tan \frac{\pi}{4} + \sec \frac{\pi}{4} \right) - (\tan 0 + \sec 0) \right]$
 $= 1 + \sqrt{2} - (0 + 1)$
 $= \sqrt{2}$

(c) $1 = \int_a^1 \frac{1}{x} dx$
 $= [\ln x]_a^1$
 $= \ln 1 - \ln a$
 $= 0 - \ln a$
 $= -\ln a$
 $-1 = \ln a$
 $a = e^{-1}$

$1 = \int_1^b \frac{1}{x} dx$
 $= [\ln x]_1^b$
 $= [\ln b - \ln 1]$
 $= \ln b - 0$
 $= \ln b$
 $b = e^1$
 $= e$
 $\therefore a = e^{-1}$ and $b = e$

Question 6



(a) (i) let $u = x + 2$ and $v = x^2 + 4$
 $u' = 1$ $v' = 2x$
 $f'(x) = u'v + uv'$
 $= 1 \times (x^2 + 4) + (x + 2) \times 2x$
 $= x^2 + 4 + 2x^2 + 4x$
 $= 3x^2 + 4x + 4$
 $a = 3, b = 4, c = 4$
 $\Delta = b^2 - 4ac$
 $= 4^2 - 4 \times 3 \times 4$
 $= -32$
 $\Delta < 0 \therefore$ no real roots
 $\therefore f'(x) \neq 0$
 $\therefore f(x)$ has no stationary points.

(ii) $f'(x) = 3x^2 + 4x + 4$
 $f''(x) = 6x + 4$
 $f''(x) < 0$ when $6x + 4 < 0$
 $x < -\frac{2}{3}$
 $f''(x) > 0$ when $6x + 4 > 0$
 $x > -\frac{2}{3}$
 $\therefore f(x)$ is concave down when $x < -\frac{2}{3}$

$\therefore f(x)$ is concave up when $x > -\frac{2}{3}$

(iii) $f(x) = (x + 2)(x^2 + 4)$
 when $x = 0$,
 $y = (0 + 2)(0^2 + 4)$
 $y = 8$
 when $y = 0$
 $0 = (x + 2)(x^2 + 4)$
 $x + 2 = 0$ or $x^2 + 4 = 0$
 $x = -2$ no solutions

From (ii), change in concavity
 when $x = -\frac{2}{3}$ ($y \approx 5.9$)



$l = r\theta$
 $9 = 5\theta$
 $\theta = \frac{9}{5} = 1.8$ radians

(ii) Aim: Prove $\triangle OPT \cong \triangle OQT$
 Data: arc $PQ = 9$ cm
 $OP = OQ = 5$ cm
 $\angle OPT = \angle OQT = 90^\circ$
 Construction: Line OT
 Proof: $OP = OQ$ (radii)
 $\angle OPT = \angle OQT$ (given)
 OT (common)
 $\therefore \triangle OPT \cong \triangle OQT$ (RHS)

(iii) $\angle POQ = 1.8$ radians
 $\therefore \angle TOP = 0.9$ radians
 since OT bisects $\angle POQ$
 $\tan 0.9^\circ = \frac{PT}{5}$
 $PT = 5 \tan 0.9^\circ$
 $PT = 6.30079\dots$
 $PT = 6.3$ cm (1 dp)

(iv) Area = $2 \times$ Area $\triangle OPT$ - Area sector
 $= 2 \times \frac{1}{2}bh - \frac{1}{2}r^2\theta$
 $= 2 \times \frac{1}{2} \times 5 \times 6.3 - \frac{1}{2} \times 5^2 \times 1.8^\circ$
 $= 31.5 - 22.5$
 $= 9$ cm² (0 dp)

Question 7


(a) (i) $\ddot{x} = 4 \cos 2t$
 $\dot{x} = \int 4 \cos 2t dt$
 $= \frac{4 \sin 2t}{2} + C$
 $= 2 \sin 2t + C$
 when $t = 0, \dot{x} = 1,$
 $1 = 2 \sin 0 + C$
 $\therefore C = 1$
 $\therefore \dot{x} = 2 \sin 2t + 1$

(ii) when $\dot{x} = 0,$
 $2 \sin 2t + 1 = 0$
 $2 \sin 2t = -1$
 $\sin 2t = -\frac{1}{2}$
 $2t = \sin^{-1}\left(-\frac{1}{2}\right)$
 $2t = \pi + \frac{\pi}{6}$
 $= \frac{7\pi}{6}$
 $t = \frac{7\pi}{12}$

\therefore The particle first comes to rest after $\frac{7\pi}{12}$ seconds.

(iii) $x = \int (2 \sin 2t + 1) dt$
 $= -\frac{2 \cos 2t}{2} + t + C$
 $= t - \cos 2t + C$
 when $t = 0, x = 0:$
 $0 = 0 - \cos 0 + C$
 $\therefore C = 1$
 $\therefore x = t - \cos 2t + 1$

(b)



(i) $y = x^2$
 $\frac{dy}{dx} = 2x$
 when $x = -1, \frac{dy}{dx} = -2$
 equation of tangent at A:
 $y - y_1 = m(x - x_1)$
 $y - 1 = -2(x - (-1))$
 $y = -2x - 1$
 $2x + y + 1 = 0.$

(ii) $M = \left(\frac{2 + (-1)}{2}, \frac{4 + 1}{2}\right)$
 $= \left(\frac{1}{2}, \frac{5}{2}\right)$
 $m_{AB} = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{4 - 1}{2 - (-1)}$
 $= 1$

The gradient of tangent at C = m_{AB}
 For the tangent at C:

$\frac{dy}{dx} = 2x$ and $2x = 1$
 $\therefore x = \frac{1}{2}$

The line MC is vertical since M and C have the same x-coordinates.

(iii) From (i), the equation of the tangent at A is $y = -2x - 1$

At T, $x = \frac{1}{2}$
 $y = -2x - 1$
 $= -2\left(\frac{1}{2}\right) - 1$
 $= -2$
 $\therefore T\left(\frac{1}{2}, -2\right).$

$m_{BT} = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{4 - (-2)}{2 - \frac{1}{2}}$
 $= 4$

$\frac{dy}{dx} = 2x$
 at B(2, 4):
 $\frac{dy}{dx} = 2 \times 2$
 $= 4$
 $= m_{BT}$

$\therefore BT$ is a tangent to the parabola.

Question 8

(a) $\frac{dP}{dt} = kP$
 $P = P_0 e^{kt}$
 when $t = 0, P = 102:$
 $102 = P_0 e^0$
 $\therefore P_0 = 102$
 $\therefore P = 102e^{kt}$

when $t = 75, P = 200 \times 10^6,$
 $200 \times 10^6 = 102e^{75k}$
 $e^{75k} = \frac{200 \times 10^6}{102}$


$75k = \ln\left(\frac{200 \times 10^6}{102}\right)$
 $\therefore k = \frac{\ln\left(\frac{200 \times 10^6}{102}\right)}{75}$
 $\approx 0.19318\dots$

when $t = 100,$
 $P = 102e^{100k}$
 $= 102e^{100 \times 0.19318\dots}$
 $\approx 2.503263\dots \times 10^{10}$
 $= 2.5 \times 10^{10}$ (2 sf)

There will be approximately 2.5×10^{10} cane toads in 2035.

(b) $P(HH) = P(H) \times P(H)$
 $= 0.36$
 $P(H) = 0.6$
 $\therefore P(T) = 1 - 0.6$
 $= 0.4$
 $\therefore P(TT) = 0.4 \times 0.4$
 $= 0.16$

(c) (i) $A = 4.$
 (ii) $b = 2.$



- (d) $f(x) = x^3 - 3x^2 + kx + 8$
 $f(x)$ is increasing when $f'(x) > 0$
 $f'(x) = 3x^2 - 6x + k > 0$
 For positive definite, $a > 0$ and $\Delta < 0$:
 $a = 3$ ($a > 0$)
 $\Delta < 0$
 $b^2 - 4ac < 0$
 $6^2 - 4 \times 3 \times k < 0$
 $36 - 12k < 0$
 $12k > 36$
 $\therefore k > 3$

Question 9

- (a) (i) First amount $A_1 = 500(1.005)^{240}$
 Second amount $A_2 = 500(1.005)^{239}$
 Third amount $A_3 = 500(1.005)^{238}$
 \vdots
 240th amount $A_{240} = 500(1.005)^1$
 $P = A_1 + A_2 + A_3 + \dots + A_{240}$
 $= 500(1.005^{240} + 1.005^{239} + \dots + 1.005^1)$
 $= 500 \underbrace{(1.005^1 + 1.005^2 + \dots + 1.005^{240})}$
 This is a geometric series
 where $a = 1.005$, $r = 1.005$, $n = 240$
 $S_n = \frac{a(r^n - 1)}{r - 1}$
 $S_{240} = \frac{1.005(1.005^{240} - 1)}{1.005 - 1}$
 Thus $P = 500 \times \frac{1.005(1.005^{240} - 1)}{1.005 - 1}$
 $= \$232\,175.55$ (nearest cent)
- (ii) $A_1 = P \times 1.005 - 2000$
 (1) $A_2 = A_1 \times 1.005 - 2000$
 $= (P \times 1.005 - 2000) \times 1.005 - 2000$
 $= P(1.005)^2 - 2000(1 + 1.005)$

$$A_n = P(1.005)^n - 2000(1 + 1.005 + \dots + 1.005^{n-1})$$

$$= P(1.005)^n - 2000 \times \left(\frac{1.005^n - 1}{1.005 - 1} \right)$$

$$= P(1.005)^n - 2000 \times \left(\frac{1.005^n - 1}{0.005} \right)$$


$$= P(1.005)^n - 400\,000 \times [(1.005)^n - 1]$$

$$= (P - 400\,000)(1.005)^n + 400\,000$$

This is the required result.

- (ii) $A_n = 0$ when
 (2) $(P - 400\,000)(1.005)^n + 400\,000 = 0$
 $1.005^n = \frac{-400\,000}{P - 400\,000}$
 $= \frac{400\,000}{400\,000 - P}$
 $n \ln 1.005 = \ln \left(\frac{400\,000}{400\,000 - P} \right)$
 $n = \ln \left(\frac{400\,000}{400\,000 - P} \right) \div \ln 1.005$
 $= \ln \left(\frac{400\,000}{400\,000 - 232\,175.55} \right) \div \ln 1.005$
 $\approx 174.143 \dots$
 \therefore There will be money in the account for 175 months.

- (b) (i) $f(x)$ is increasing when $f'(x) > 0$.
 From the graph, this is when $0 \leq x < 2$.
- (ii) Maximum at $x = 2$ since $f'(x) = 0$ and the curve changes from increasing to decreasing at this point. $A_1 = 4 \therefore f(2) = 4$.
 \therefore the maximum value is 4 units.
- (iii) $f(4) = 0$ (from symmetry about $x = 2$)
 $f(6) = f(4) + 2 \times -3$
 $= 0 - 6$
 $= -6$.



Question 10

- (a) (i) $\triangle ABC$ is isosceles ($AC = BC$, given)
 $\therefore \angle ABC = \angle BAC$ (base \angle 's of isosceles \triangle)
 $\triangle ACD$ is isosceles ($AD = CD$, given)
 $\therefore \angle ACD = \angle CAD$ (base \angle 's of isosceles \triangle)
 But $\angle BAC = \angle CAD$ (same angle)
 $\therefore \angle ABC = \angle BAC = \angle ACD = \angle CAD$
 In \triangle 's ABC and ACD :
 $\angle ABC = \angle ACD$ (shown above)
 $\angle BAC = \angle CAD$ (shown above)
 $\therefore \triangle ABC \parallel \triangle ACD$ (equi-angular)
- (ii) $\frac{AB}{AC} = \frac{AC}{AD}$ (corres. sides of similar \triangle s are in proportion)
 $\therefore \frac{a+y}{x} = \frac{x}{a}$
 $\therefore x^2 = a(a+y)$
 $= a^2 + ay$
- (iii) Using the cosine rule:
 $x^2 = a^2 + a^2 - 2a^2 \cos \theta$
 $\therefore a^2 + ay = 2a^2 - 2a^2 \cos \theta$
 $ay = a^2 - 2a^2 \cos \theta$
 $[\div a] \rightarrow y = a - 2a \cos \theta$
 $\therefore y = a(1 - 2 \cos \theta)$

- (iv) $y = a - 2a \cos \theta$ from (iii)
 $y - a = -2a \cos \theta$
 $\cos \theta = \frac{y-a}{2a}$
 $-1 \leq \cos \theta \leq 1$
 $-1 \leq \frac{y-a}{2a} \leq 1$
 $-2a \leq y - a \leq 2a$
 $-a \leq y \leq 3a$
 $y > 0$ as DB is a length
 $\therefore y \leq 3a$

- (b) (i) $x^2 + y^2 = r^2$
 $\therefore y^2 = r^2 - x^2$
 Also, from $\triangle OPA$:
 $OA = r \sin \theta$
 $V = \pi \int y^2 dx$
 $= \pi \int_{r \sin \theta}^r (r^2 - x^2) dx$
 $= \pi \left[r^2 x - \frac{x^3}{3} \right]_{r \sin \theta}^r$
 $= \pi \left[r^3 - \frac{r^3}{3} - \left(r^3 \sin \theta - \frac{r^3 \sin^3 \theta}{3} \right) \right]$
 $= \frac{\pi r^3}{3} (3 - 1 - 3 \sin \theta + \sin^3 \theta)$
 $= \frac{\pi r^3}{3} (2 - 3 \sin \theta + \sin^3 \theta)$
- (ii) $OA = r \sin \theta = \frac{r}{2}$
 (1) $\therefore \sin \theta = \frac{1}{2}$
 $\theta = \sin^{-1} \left(\frac{1}{2} \right)$
 $= \frac{\pi}{6}$ (or 30°)

$$\begin{aligned} \text{(ii) Original volume} &= \frac{1}{2} \times \frac{4}{3} \pi r^3 \\ \text{(2)} &= \frac{2\pi r^3}{3} \end{aligned}$$

When $\theta = \frac{\pi}{6}$, from (i) :

$$\begin{aligned} V &= \frac{\pi r^3}{3} (2 - 3 \sin \theta + \sin^3 \theta) \\ &= \frac{\pi r^3}{3} \left(2 - 3 \sin \frac{\pi}{6} + \sin^3 \frac{\pi}{6} \right) \\ &= \frac{\pi r^3}{3} \left(2 - 3 \times \frac{1}{2} + \left(\frac{1}{2} \right)^3 \right) \\ &= \frac{5\pi r^3}{24} \end{aligned}$$

$$\begin{aligned} \therefore \text{Fraction} &= \frac{5\pi r^3}{24} \div \frac{2\pi r^3}{3} \\ &= \frac{5\pi r^3}{24} \times \frac{3}{2\pi r^3} \\ &= \frac{5}{16} \end{aligned}$$

End of Mathematics solutions
