

BOARD OF STUDIES
NEW SOUTH WALES

2011

HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 1

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
Black pen is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

Total marks – 84

- Attempt Questions 1–7
- All questions are of equal value

Total marks – 84

Attempt Questions 1–7

All questions are of equal value

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

Question 1 (12 marks) Use a SEPARATE writing booklet.

- (a) The point P divides the interval joining $A(-1, -2)$ to $B(9, 3)$ internally in the ratio $4 : 1$. Find the coordinates of P . 2
- (b) Differentiate $\frac{\sin^2 x}{x}$ with respect to x . 2
- (c) Solve $\frac{4-x}{x} < 1$. 3
- (d) Using the substitution $u = \sqrt{x}$, evaluate $\int_1^4 \frac{e^{\sqrt{x}}}{\sqrt{x}} dx$. 3
- (e) Find the exact value of $\cos^{-1}\left(-\frac{1}{2}\right)$. 1
- (f) What is the range of the function $f(x) = \ln(x^2 + e)$? 1

Question 2 (12 marks) Use a SEPARATE writing booklet.

- (a) Let $P(x) = x^3 - ax^2 + x$ be a polynomial, where a is a real number. 3

When $P(x)$ is divided by $x - 3$ the remainder is 12.

Find the remainder when $P(x)$ is divided by $x + 1$.

- (b) The function $f(x) = \cos 2x - x$ has a zero near $x = \frac{1}{2}$. 3

Use one application of Newton's method to obtain another approximation to this zero. Give your answer correct to two decimal places.

- (c) Find an expression for the coefficient of x^2 in the expansion of $\left(3x - \frac{4}{x}\right)^8$. 2

- (d) Sketch the graph of the function $f(x) = 2\cos^{-1}x$. Clearly indicate the domain and range of the function. 2

- (e) Alex's playlist consists of 40 different songs that can be arranged in any order.

- (i) How many arrangements are there for the 40 songs? 1

- (ii) Alex decides that she wants to play her three favourite songs first, in any order. 1

How many arrangements of the 40 songs are now possible?

Question 3 (12 marks) Use a SEPARATE writing booklet.

- (a) The equation of motion for a particle undergoing simple harmonic motion is

$$\frac{d^2x}{dt^2} = -n^2x,$$

where x is the displacement of the particle from the origin at time t , and n is a positive constant.

- (i) Verify that $x = A \cos nt + B \sin nt$, where A and B are constants, is a solution of the equation of motion. 1

- (ii) The particle is initially at the origin and moving with velocity $2n$. 2

Find the values of A and B in the solution $x = A \cos nt + B \sin nt$.

- (iii) When is the particle first at its greatest distance from the origin? 1


- (iv) What is the total distance the particle travels between $t = 0$ and 1

$$t = \frac{2\pi}{n}?$$

Question 3 continues on page 5

Question 3 (continued)

- (b) The diagram shows two distinct points $P(t, t^2)$ and $Q(1-t, (1-t)^2)$ on the parabola $y = x^2$. The point R is the intersection of the tangents to the parabola at P and Q .



- (i) Show that the equation of the tangent to the parabola at P is $y = 2tx - t^2$. 2
- (ii) Using part (i), write down the equation of the tangent to the parabola at Q . 1
- (iii) Show that the tangents at P and Q intersect at $R\left(\frac{1}{2}, t - t^2\right)$. 2
- (iv) Describe the locus of R as t varies, stating any restriction on the y -coordinate. 2

End of Question 3


Question 4 (12 marks) Use a SEPARATE writing booklet.

- (a) Consider the function $f(x) = e^{-x} - 2e^{-2x}$.
- (i) Find $f'(x)$. 1
- (ii) The graph $y = f(x)$ has one maximum turning point. 2
Find the coordinates of the maximum turning point.
- (iii) Evaluate $f(\ln 2)$. 1
- (iv) Describe the behaviour of $f(x)$ as $x \rightarrow \infty$. 1
- (v) Find the y -intercept of the graph $y = f(x)$. 1
- (vi) Sketch the graph $y = f(x)$ showing the features from parts (ii)–(v). 2
You are not required to find any points of inflexion.

Question 4 continues on page 7

Question 4 (continued)

- (b) In the diagram, the vertices of $\triangle ABC$ lie on the circle with centre O . The point D lies on BC such that $\triangle ABD$ is isosceles and $\angle ABC = x$.



Copy or trace the diagram into your writing booklet.


- (i) Explain why $\angle AOC = 2x$. 1
- (ii) Prove that $ACDO$ is a cyclic quadrilateral. 2
- (iii) Let M be the midpoint of AC and P the centre of the circle through A , C , D and O . 1

Show that P , M and O are collinear.

End of Question 4

Question 5 (12 marks) Use a SEPARATE writing booklet.

- (a) In the diagram, $Q(x_0, y_0)$ is a point on the unit circle $x^2 + y^2 = 1$ at an angle θ from the positive x -axis, where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. The line through $N(0, 1)$ and Q intersects the line $y = -1$ at P . The points $T(0, y_0)$ and $S(0, -1)$ are on the y -axis.



- (i) Use the fact that $\triangle TQN$ and $\triangle SPN$ are similar to show that $SP = \frac{2 \cos \theta}{1 - \sin \theta}$. 2
- (ii) Show that $\frac{\cos \theta}{1 - \sin \theta} = \sec \theta + \tan \theta$. 1
- (iii) Show that $\angle SNP = \frac{\theta}{2} + \frac{\pi}{4}$. 1
- (iv) Hence, or otherwise, show that $\sec \theta + \tan \theta = \tan\left(\frac{\theta}{2} + \frac{\pi}{4}\right)$. 1
- (v) Hence, or otherwise, solve $\sec \theta + \tan \theta = \sqrt{3}$, where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. 2

Question 5 continues on page 9

Question 5 (continued)

(b) To test some forensic science students, an object has been left in a park. At 10 am the temperature of the object is measured to be 30°C. The temperature in the park is a constant 22°C. The object is moved immediately to a room where the temperature is a constant 5°C.

(i) The temperature of the object in the room can be modelled by the equation

$$T = 5 + 25e^{-kt},$$

where T is the temperature of the object in degrees Celsius, t is the time in hours since the object was placed in the room and k is a constant.

After one hour in the room the temperature of the object is 20°C.

Show that $k = \ln\left(\frac{5}{3}\right)$.

(ii) In a similar manner, the temperature of the object in the park before it was discovered can be modelled by an equation of the form

$$T = A + Be^{-kt},$$

with the same constant $k = \ln\left(\frac{5}{3}\right)$.

Find the time of day when the object had a temperature of 37°C.


End of Question 5

Question 6 (12 marks) Use a SEPARATE writing booklet.

(a) Use mathematical induction to prove that, for $n \geq 1$,

$$1 \times 5 + 2 \times 6 + 3 \times 7 + \dots + n(n+4) = \frac{1}{6}n(n+1)(2n+13).$$

(b) The diagram shows the trajectory of a ball thrown horizontally, at speed $v \text{ m s}^{-1}$, from the top of a tower h metres above ground level.



The ball strikes the ground at an angle of 45°, d metres from the base of the tower, as shown in the diagram. The equations describing the trajectory of the ball are

$$x = vt \text{ and } y = h - \frac{1}{2}gt^2, \text{ (Do NOT prove this.)}$$

where g is the acceleration due to gravity, and t is time in seconds.

(i) Prove that the ball strikes the ground at time $t = \sqrt{\frac{2h}{g}}$ seconds.

(ii) Hence, or otherwise, show that $d = 2h$.

Question 6 continues on page 11

Question 6 (continued)

- (c) A game is played by throwing darts at a target. A player can choose to throw two or three darts.

Darcy plays two games. In Game 1, he chooses to throw two darts, and wins if he hits the target at least once. In Game 2, he chooses to throw three darts, and wins if he hits the target at least twice.


The probability that Darcy hits the target on any throw is p , where $0 < p < 1$.

- (i) Show that the probability that Darcy wins Game 1 is $2p - p^2$. 1
- (ii) Show that the probability that Darcy wins Game 2 is $3p^2 - 2p^3$. 1
- (iii) Prove that Darcy is more likely to win Game 1 than Game 2. 2
- (iv) Find the value of p for which Darcy is twice as likely to win Game 1 as he is to win Game 2. 2

End of Question 6

Question 7 (12 marks) Use a SEPARATE writing booklet.

- (a) The diagram shows two identical circular cones with a common vertical axis. Each cone has height h cm and semi-vertical angle 45° .



NOT TO SCALE

The lower cone is completely filled with water. The upper cone is lowered vertically into the water as shown in the diagram. The rate at which it is lowered is given by

$$\frac{d\ell}{dt} = 10,$$

where ℓ cm is the distance the upper cone has descended into the water after t seconds.

As the upper cone is lowered, water spills from the lower cone. The volume of water remaining in the lower cone at time t is V cm³.

- (i) Show that 1

$$V = \frac{\pi}{3}(h^3 - \ell^3).$$

- (ii) Find the rate at which V is changing with respect to time when $\ell = 2$. 2
- (iii) Find the rate at which V is changing with respect to time when the lower cone has lost $\frac{1}{8}$ of its water. Give your answer in terms of h . 2

Question 7 continues on page 13

Question 7 (continued)

(b) The binomial theorem states that

$$(1+x)^n = \sum_{r=0}^n \binom{n}{r} x^r$$

for all integers $n \geq 1$.

(i) Show that 2

$$\sum_{r=1}^n \binom{n}{r} r x^r = n x (1+x)^{n-1}.$$

(ii) By differentiating the result from part (i), or otherwise, show that 2

$$\sum_{r=1}^n \binom{n}{r} r^2 = n(n+1)2^{n-2}.$$

(iii) Assume now that n is even. Show that, for $n \geq 4$, 3

$$\binom{n}{2} 2^2 + \binom{n}{4} 4^2 + \binom{n}{6} 6^2 + \dots + \binom{n}{n} n^2 = n(n+1)2^{n-3}.$$

End of paper

2011 Higher School Certificate Solutions Mathematics Extension 1

Question 1

(a) $A(-1, -2), B(9, 3)$, ratio 4:1

$$P = \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right)$$

$$= \left(\frac{1 \times -1 + 9 \times 4}{4+1}, \frac{1 \times -2 + 3 \times 4}{4+1} \right)$$

$$= \left(\frac{35}{5}, \frac{10}{5} \right)$$

$$= (7, 2)$$

(b) Using the quotient rule:

$$\frac{d}{dx} \left(\frac{\sin^2 x}{x} \right) = \frac{2 \sin x \cos x (x) - \sin^2 x}{x^2}$$

$$= \frac{2x \sin x \cos x - \sin^2 x}{x^2}$$

(c) **METHOD 1** – Algebraic:

$$\frac{4-x}{x} < 1, x \neq 0$$

$$\frac{4-x}{x} \times x^2 < 1 \times x^2$$


$$x(4-x) < x^2$$

$$4x - x^2 - x^2 < 0$$

$$4x - 2x^2 < 0$$

$$2x^2 - 4x > 0$$

$$2x(x-2) > 0$$



$\therefore x < 0$ or $x > 2$.

METHOD 2 – Critical Point:

Consider $\frac{4-x}{x} = 1$ (noting $x \neq 0$)

$$4-x = x$$

$$2x = 4$$


$$x = 2$$

$\therefore x = 0$ and $x = 2$ are critical points.

Test $x = -1$ $\frac{5}{-1} < 1$ True

$x = 1$ $\frac{3}{1} < 1$ False

$x = 3$ $\frac{1}{3} < 1$ True



$\therefore x < 0$ or $x > 2$.

(d)

$$u = \sqrt{x} = x^{\frac{1}{2}}$$

$$\frac{du}{dx} = \frac{1}{2} x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}}$$

$$2du = \frac{1}{\sqrt{x}} dx$$

when $x = 4, u = 2$

when $x = 1, u = 1$

$$\int_1^4 \frac{e^{\sqrt{x}}}{\sqrt{x}} dx = 2 \int_1^2 e^u du$$

$$= 2[e^u]_1^2$$

$$= 2(e^2 - e)$$

(e) $\cos 60^\circ = \frac{1}{2} \Rightarrow \cos^{-1} \left(\frac{1}{2} \right) = 60^\circ = \frac{\pi}{3}$

$$\cos^{-1} \left(-\frac{1}{2} \right) = \pi - \frac{\pi}{3}$$

$$= \frac{2\pi}{3}$$

(f) $x^2 \geq 0$ and $e > 0$

The minimum value of x^2 is 0.

The minimum value of $x^2 + e$ is e .

The minimum value of:

$$\ln(x^2 + e) \text{ is } \ln(e) = 1.$$

Thus the range is $f(x) \geq 1$.

Question 2

(a) $P(x) = x^3 - ax^2 + x$

$$P(3) = 12$$

$$12 = (3)^3 - a(3)^2 + 3$$

$$12 = 27 - 9a + 3$$

$$12 = 30 - 9a$$

$$9a = 18$$

$$a = 2$$

$$P(x) = x^3 - 2x^2 + x$$

$$P(-1) = (-1)^3 - 2(-1)^2 + (-1)$$

$$= -1 - 2 - 1$$

$$= -4$$

\therefore The remainder is -4 .

(b) $f(x) = \cos 2x - x$

$$f'(x) = -2 \sin 2x - 1$$

$$x_2 = x_1 - \frac{f(x)}{f'(x)}$$

$$x_2 = \frac{1}{2} - \frac{f\left(\frac{1}{2}\right)}{f'\left(\frac{1}{2}\right)}$$

$$= \frac{1}{2} - \left(\frac{\cos(1) - \frac{1}{2}}{-2 \sin(1) - 1} \right)$$

$$= 0.515021683\dots$$

$$= 0.52 \text{ (2dp)}$$

(c) $T_k = {}^8C_k (3x)^{8-k} \left(\frac{-4}{x} \right)^k$

$$= {}^8C_k \times 3^{8-k} \times x^{8-k} \times \frac{(-4)^k}{x^k}$$

$$= {}^8C_k \times 3^{8-k} \times (-4)^k \times \frac{x^{8-k}}{x^k}$$

$$= {}^8C_k \times 3^{8-k} \times (-4)^k \times x^{8-2k}$$

Indices of x^{8-2k} and x^2 :

$$8 - 2k = 2$$

$$2k = 6$$

$$k = 3$$


$$T_3 = {}^8C_3 \times 3^{8-3} \times (-4)^3 \times x^2$$

$$= 56 \times 243 \times (-64) \times x^2$$

$$= -870,912x^2$$

\therefore The coefficient is $-870,912$.

(d)



Domain: $-1 \leq x \leq 1$

Range: $0 \leq y \leq 2\pi$

(e) (i) 40!

(ii) $3! \times 37!$

Question 3

(a) (i) $x = A \cos nt + B \sin nt$

$$\dot{x} = -An \sin nt + Bn \cos nt$$

$$\ddot{x} = -An^2 \cos nt - Bn^2 \sin nt$$

$$= -n^2 (A \cos nt + B \sin nt)$$


$$= -n^2 x$$

Verified as required.

(ii) $t=0, x=0, v=2n$
 $x = A \cos nt + B \sin nt$
 $0 = A \cos(n \times 0) + B \sin(n \times 0)$
 $0 = A \times 1 + B \times 0$
 $A = 0$
 $\dot{x} = -A n \sin nt + B n \cos nt$
 $2n = -(0 \times n) \sin(n \times 0) + B n \cos(n \times 0)$
 $2n = 0 + B n \times 1$
 $2n = B n$
 $B = 2$
 $\therefore A = 0, B = 2.$

(iii) **METHOD 1**
 Greatest distance when $\dot{x} = 0$
 $\dot{x} = -A n \sin nt + B n \cos nt$
 $0 = 2n \cos nt$ (since $A = 0, B = 2$)
 $0 = \cos nt$
 $\cos^{-1} 0 = nt$
 $\frac{\pi}{2} = nt$
 $t = \frac{\pi}{2n}.$

METHOD 2
 Max distance is the amplitude of x .
 $x = 2 \sin nt$
 Thus max distance is at $x = 2$.
 $2 = 2 \sin nt$
 $1 = \sin nt$
 $\sin^{-1} 1 = nt$
 $nt = \frac{\pi}{2}$
 $t = \frac{\pi}{2n}.$



(b) (i) $y = x^2$
 $\frac{dy}{dx} = 2x$
 At $P(t, t^2), x = t$
 $m_p = 2t$
 $y - y_1 = m(x - x_1)$
 $y - t^2 = 2t(x - t)$
 $y - t^2 = 2tx - 2t^2$
 $y = 2tx - t^2.$

(ii) Sub $1-t$ for t into $y = 2tx - t^2$
 $y = 2(1-t)x - (1-t)^2.$

(iii) $y = 2tx - t^2$
 $y = 2(1-t)x - (1-t)^2$
 Solving simultaneously:
 $2tx - t^2 = 2(1-t)x - (1-t)^2$
 $2tx - 2(1-t)x = t^2 - (1-t)^2$
 $2x[t - (1-t)] = [t - (1-t)][t + (1-t)]$
 $2x(2t - 1) = (t - 1 + t)(t + 1 - t)$
 $2x(2t - 1) = (2t - 1) \cdot 1$
 $2x = 1$
 $x = \frac{1}{2}$
 When $x = \frac{1}{2}$:
 $y = 2t\left(\frac{1}{2}\right) - t^2$
 $= t - t^2$
 $\therefore R$ is $\left(\frac{1}{2}, t - t^2\right).$

(iv) The locus is the vertical line, $x = \frac{1}{2}$
 The tangents cannot meet inside the parabola and the points P and Q cannot be the same point, so y must be below the curve.

When $x = \frac{1}{2}: y = \left(\frac{1}{2}\right)^2$
 $= \frac{1}{4}$
 So the restriction on y is $y < \frac{1}{4}.$

Question 4

(a) (i) $f(x) = e^{-x} - 2e^{-2x}$
 $f'(x) = -e^{-x} + 4e^{-2x}$

(ii) Stationary points when $f'(x) = 0$
 $-e^{-x} + 4e^{-2x} = 0$
 $4e^{-2x} = e^{-x}$
 $4 = e^x$
 $x = \ln 4$
 $f(\ln 4) = e^{-\ln 4} - 2e^{-2 \ln 4}$
 $= e^{-\ln 4} - 2e^{-\ln 16}$
 $= \frac{1}{4} - \frac{2}{16}$
 $= \frac{1}{4} - \frac{1}{8}$
 $= \frac{1}{8}$

The stationary point occurs at $(\ln 4, \frac{1}{8}).$



$f''(x) = e^{-x} - 8e^{-2x}$
 $f''(\ln 4) = e^{-\ln 4} - 8e^{-2 \ln 4}$
 $= \frac{1}{4} - 8\left(\frac{1}{16}\right)$
 $= \frac{1}{4} - \frac{1}{2}$
 $= -\frac{1}{4} < 0$

So $(\ln 4, \frac{1}{8})$ is a maximum turning point.

(iii) $f(\ln 2) = e^{-\ln 2} - 2e^{-2 \ln 2}$
 $= \frac{1}{e^{\ln 2}} - \frac{2}{e^{\ln 4}}$
 $= \frac{1}{2} - \frac{2}{4}$
 $= 0.$

(iv) as $x \rightarrow \infty$
 $e^{-x} \rightarrow 0$ and $e^{-2x} \rightarrow 0$
 so $f(x) \rightarrow 0.$

(v) When $x = 0,$
 $y = e^0 - 2e^0$
 $= 1 - 2$
 $= -1.$



$\angle AOC = 2x$ (angle at the centre is twice the angle at the circumference subtended by the same arc AC).

(ii) $\angle DAB = \angle DBA$ (isosceles triangle)
 $\angle CDA = 2x$ (exterior angle of $\triangle ADB$)
 so $\angle AOC = \angle CDA = 2x$
 $\therefore ACDO$ is a cyclic quadrilateral (angles subtended on the same arc AC at the points D and O are equal).

(iii) M is the midpoint of AC (given)
 $OM \perp AC$ (The line joining the midpoint of a chord and the centre of the circle is perpendicular to the chord).

$OP \perp AC$ (The line joining the midpoint of a chord and the centre of the circle is perpendicular to the chord).

$\therefore M, P$ and O are collinear.

Question 5

- (a) (i) From the definitions of $\sin \theta$ and $\cos \theta$ in the unit circle:

$$TQ = \cos \theta$$

$$OT = \sin \theta$$

$$TN = 1 - \sin \theta$$

$$SN = 2$$

From the similar triangles:

$$\triangle TQN \sim \triangle SPN$$

$$\frac{SP}{SN} = \frac{TQ}{TN}$$

$$\frac{SP}{2} = \frac{\cos \theta}{1 - \sin \theta}$$

$$SP = \frac{2 \cos \theta}{1 - \sin \theta}$$

(ii)
$$\begin{aligned} LHS &= \frac{\cos \theta}{1 - \sin \theta} \\ &= \frac{\cos \theta}{1 - \sin \theta} \times \frac{1 + \sin \theta}{1 + \sin \theta} \\ &= \frac{\cos \theta(1 + \sin \theta)}{1 - \sin^2 \theta} \\ &= \frac{\cos \theta(1 + \sin \theta)}{\cos^2 \theta} \\ &= \frac{(1 + \sin \theta)}{\cos \theta} \\ &= \frac{1}{\cos \theta} + \frac{\sin \theta}{\cos \theta} \\ &= \sec \theta + \tan \theta \\ &= RHS \end{aligned}$$

(iii) $\angle SOQ = \frac{\pi}{2} + \theta$

$$\angle SNQ = \frac{1}{2} \angle SOQ$$

(Angle at circumference of circle is half the angle at centre standing on same arc SQ)

$$\therefore \angle SNP = \frac{1}{2} \left(\frac{\pi}{2} + \theta \right) = \frac{\theta}{2} + \frac{\pi}{4}$$

(iv)
$$\begin{aligned} RHS &= \tan \left(\frac{\theta}{2} + \frac{\pi}{4} \right) \\ &= \tan(\angle SNP) \\ &= \frac{SP}{SN} \\ &= \frac{\left(\frac{2 \cos \theta}{1 - \sin \theta} \right)}{2} \\ &= \frac{2 \cos \theta}{1 - \sin \theta} \times \frac{1}{2} \\ &= \frac{\cos \theta}{1 - \sin \theta} \\ &= \sec \theta + \tan \theta \quad (\text{from part (ii)}). \end{aligned}$$

(v)
$$\begin{aligned} \sec \theta + \tan \theta &= \sqrt{3} \\ \tan \left(\frac{\theta}{2} + \frac{\pi}{4} \right) &= \sqrt{3} \quad \text{from part (iv)} \\ \frac{\theta}{2} + \frac{\pi}{4} &= \frac{\pi}{3}, \frac{4\pi}{3}, \frac{7\pi}{3}, \dots \end{aligned}$$

From the limits on θ :

$$-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$-\frac{\pi}{4} \leq \frac{\theta}{2} \leq \frac{\pi}{4}$$

$$0 \leq \frac{\theta}{2} + \frac{\pi}{4} \leq \frac{\pi}{2}$$

So the only solution to

$$\tan \left(\frac{\theta}{2} + \frac{\pi}{4} \right) = \sqrt{3}$$

$$\frac{\theta}{2} + \frac{\pi}{4} = \frac{\pi}{3}$$

$$\frac{\theta}{2} = \frac{\pi}{12}$$

$$\theta = \frac{\pi}{6}$$

(b) (i) $T = 5 + 25e^{-kt}$
 when $t = 1$, $T = 20$
 $20 = 5 + 25e^{-k \times 1}$
 $\frac{15}{25} = e^{-k}$
 $\ln \left(\frac{3}{5} \right) = -k$
 $k = -\ln \left(\frac{3}{5} \right)$
 $= \ln \left(\frac{5}{3} \right)$

- (ii) Similarly to (i), A will be the surrounding temperature, so $A = 22$. Let t be the time after 10 am in hours. We would expect a negative answer for t . Also, when $t = 0$, $T = 30$, so

$$T = A + Be^{-kt}$$

$$30 = 22 + Be^{-k \times 0}$$

$$8 = B$$

The time when $T = 37$ will be given by solving:

$$37 = 22 + 8e^{-kt}$$

$$15 = 8e^{-kt}$$

$$\frac{15}{8} = e^{-kt}$$

$$-kt = \ln \left(\frac{15}{8} \right)$$

$$kt = \ln \left(\frac{8}{15} \right)$$

$$t = \ln \left(\frac{8}{15} \right) \div k$$

Now, the constant is the same as in (i),

$$\text{so } k = \ln \left(\frac{5}{3} \right)$$

$$t = \ln \left(\frac{8}{15} \right) \div \ln \left(\frac{5}{3} \right)$$

$$\approx -1.23$$

$$\approx -1 \text{ hr } 14 \text{ min}$$

Therefore, the object had a temperature of 37°C at 8:46 am.

Question 6

- (a) Prove for $n = 1$:

$$LHS = 1 \times 5$$

$$= 5$$

$$RHS = \frac{1}{6} \times 1 \times (1+1) \times (2 \times 1 + 1)$$

$$= \frac{1}{6} \times 1 \times 2 \times 15$$

$$= 5$$

$$= LHS$$

Assume true for $n = k$:

$$1 \times 5 + 2 \times 6 + \dots + k(k+4) = \frac{1}{6} k(k+1)(2k+13)$$

Prove for $n = k + 1$:

$$1 \times 5 + 2 \times 6 + \dots + (k+1)(k+5) = \frac{1}{6} (k+1)(k+2)(2k+15)$$

$$LHS = 1 \times 5 + 2 \times 6 + \dots + k(k+4) + (k+1)(k+5)$$

$$= \frac{1}{6} k(k+1)(2k+13) + (k+1)(k+5)$$

$$= \frac{k(k+1)(2k+13) + 6(k+1)(k+5)}{6}$$

$$= \frac{(k+1)[k(2k+13) + 6(k+5)]}{6}$$

$$= \frac{(k+1)(2k^2 + 13k + 6k + 30)}{6}$$

$$= \frac{(k+1)(2k^2 + 19k + 30)}{6}$$

$$= \frac{1}{6} (k+1)(k+2)(2k+15)$$

$$= RHS$$

Therefore by the principal of mathematical induction it is true for all integers $n \geq 1$.

- (b) (i) Ball strikes the ground when $y = 0$.

$$y = h - \frac{1}{2}gt^2$$

$$0 = h - \frac{1}{2}gt^2$$

$$\frac{1}{2}gt^2 = h$$

$$t^2 = \frac{2h}{g}$$

$$t = \sqrt{\frac{2h}{g}}$$

(ii) $x = vt$
 $\dot{x} = v$
 $y = h - \frac{1}{2}gt^2$
 $\dot{y} = -gt$
 $= -g\sqrt{\frac{2h}{g}}$ from part (i)

Since the gradient is negative, the gradient is $-\tan 45^\circ$ when the ball hits the ground.

$-\tan 45^\circ = \frac{\dot{y}}{\dot{x}}$
 $-1 = \frac{\dot{y}}{\dot{x}}$
 $\dot{x} = -\dot{y}$
 $\therefore v = g\sqrt{\frac{2h}{g}}$

But $x = d$ and $x = vt$ so $d = vt$

$= g\sqrt{\frac{2h}{g}} \times \sqrt{\frac{2h}{g}}$
 $= g \times \frac{2h}{g}$
 $= 2h.$

(c) (i) In Game 1:
 $P(\text{hit}) = p$
 $P(\text{miss}) = 1 - p$
 $P(\text{win}) = P(\text{hit on first throw}) + P(\text{miss then hit})$
 $= p + (1 - p)p$
 $= p + p - p^2$
 $= 2p - p^2.$

(ii) In Game 2:
 $P(\text{win}) = P(3 \text{ hits}) + P(2 \text{ hits from } 3)$
 $= {}^3C_3 p^3 + {}^3C_2 p^2(1 - p)$
 $= p^3 + 3p^2(1 - p)$
 $= p^3 + 3p^2 - 3p^3$
 $= 3p^2 - 2p^3.$

(iii) From (i) and (ii) we need to show that $2p - p^2 > 3p^2 - 2p^3$
 $2p - p^2 - (3p^2 - 2p^3) > 0$
 So, start with
 $2p - p^2 - (3p^2 - 2p^3) = 2p - p^2 - 3p^2 + 2p^3$
 $= 2p^3 - 4p^2 + 2p$
 $= 2p(p^2 - 2p + 1)$
 $= 2p(p - 1)^2$
 > 0 since $p > 0$

\therefore Darcy is more likely to win Game 1 than Game 2.

(iv) $P(\text{win Game 1}) = 2 \times P(\text{win Game 2})$
 $2p - p^2 = 2(3p^2 - 2p^3)$
 $2p - p^2 = 6p^2 - 4p^3$
 $4p^3 - 7p^2 + 2p = 0$
 $p(4p^2 - 7p + 2) = 0$
 Since $p > 0$, the solutions will be given by:
 $p = \frac{7 \pm \sqrt{7^2 - 4 \times 4 \times 2}}{2 \times 4}$
 $= \frac{7 \pm \sqrt{17}}{8}$

Given $0 < p < 1$, and $\frac{7 + \sqrt{17}}{8} > 1$, the only solution is $p = \frac{7 - \sqrt{17}}{8}$.

Question 7

(a) (i) Since the semi-vertical angle is 45° , the radius and the height will be equal.
 $V = V_L - V_U$
 $= \frac{1}{3}\pi R^2 h - \frac{1}{3}\pi r^2 \ell$
 $= \frac{\pi}{3}(R^2 h - r^2 \ell)$ but $R = h, r = \ell$
 $= \frac{\pi}{3}(h^3 - \ell^3).$

(ii) $\frac{dV}{dt} = \frac{dV}{d\ell} \times \frac{d\ell}{dt}$
 $= \left(-\frac{\pi}{3} \times 3\ell^2\right) \times 10$
 $= -10\pi\ell^2$
 when $\ell = 2$:
 $\frac{dV}{dt} = -10 \times \pi \times 2^2$
 $= -40\pi \text{ cm}^3/\text{s econd.}$

(iii) Substitute $V = \frac{7}{8} \times \frac{\pi}{3} h^3$
 in $V = \frac{\pi}{3}(h^3 - \ell^3)$,
 $\frac{7}{8} \times \frac{\pi}{3} h^3 = \frac{\pi}{3}(h^3 - \ell^3)$
 $\frac{7}{8} \times h^3 = h^3 - \ell^3$
 $\ell^3 = \frac{1}{8} h^3$
 $\ell = \frac{h}{2}$
 $\frac{dV}{dt}$ at $\ell = \frac{h}{2}$ is:
 $\frac{dV}{dt} = -10\pi\ell^2$
 $= -10\pi \times \left(\frac{h}{2}\right)^2$
 $= -\frac{5}{2}\pi h^2 \text{ cm}^3/\text{second.}$

(b) (i) expand $(1 + x)^n = \sum_{r=0}^n \binom{n}{r} x^r$:
 $(1 + x)^n = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{n}x^n$
 $\frac{d}{dx}[(1 + x)^n] = \binom{n}{1} + \binom{n}{2}2x + \dots + \binom{n}{n}nx^{n-1}$
 $n(1 + x)^{n-1} = \binom{n}{1} + \binom{n}{2}2x + \dots + \binom{n}{n}nx^{n-1}$
 $n(1 + x)^{n-1} \times x = \binom{n}{1}x + \binom{n}{2}2x^2 + \dots + \binom{n}{n}nx^n$
 $nx(1 + x)^{n-1} = \binom{n}{1}x + \binom{n}{2}2x^2 + \dots + \binom{n}{n}nx^n$
 $\therefore \sum_{r=1}^n \binom{n}{r} r x^r = nx(1 + x)^{n-1}.$

(ii) Differentiate both sides in (i) with respect to x (product rule on RHS):
 $\sum_{r=1}^n \binom{n}{r} r \times x^{r-1} = n(1 + x)^{n-1} + nx(n-1)(1 + x)^{n-2}$
 Substitute in $x = 1$:
 $\sum_{r=1}^n \binom{n}{r} r^2 = n2^{n-1} + n(n-1)2^{n-2}$
 $= n.2.2^{n-2} + n(n-1)2^{n-2}$
 $= 2^{n-2} [2n + n^2 - n]$
 $= n(n+1)2^{n-2}.$

(iii) From part (ii):
 $\sum_{r=1}^n \binom{n}{r} r^2 x^{r-1} = n(1 + x)^{n-1} + nx(n-1)(1 + x)^{n-2}$
 Substitute $x = -1$:
 $\sum_{r=1}^n \binom{n}{r} r^2 (-1)^{r-1} = n(0)^{n-1} + n(-1)(n-1)(0)^{n-2}$
 $\sum_{r=1}^n \binom{n}{r} r^2 (-1)^{r-1} = 0$
 Expand $\sum_{r=1}^n \binom{n}{r} r^2 (-1)^{r-1} = 0$
 $\binom{n}{1} - \binom{n}{2}2^2 + \binom{n}{3}3^2 - \dots - \binom{n}{n}n^2 = 0$

Expand (ii)
 $\binom{n}{1} + \binom{n}{2}2^2 + \binom{n}{3}3^2 + \dots + \binom{n}{n}n^2 = n(n+1)2^{n-2}$

Subtract the above two expansions:
 $2 \times \left\{ \binom{n}{2}2^2 + \binom{n}{4}4^2 + \binom{n}{6}6^2 + \dots + \binom{n}{n}n^2 \right\} = n(n+1)2^{n-2} - 0$
 $\binom{n}{2}2^2 + \binom{n}{4}4^2 + \binom{n}{6}6^2 + \dots + \binom{n}{n}n^2 = n(n+1)2^{n-3}$
 $\binom{n}{2}2^2 + \binom{n}{4}4^2 + \binom{n}{6}6^2 + \dots + \binom{n}{n}n^2 = n(n+1)2^{n-3}.$