

NAME :

# CENTRE OF EXCELLENCE IN MATHS TUTORIAL

MOBILE 0412880475



PHONE 96969696

## MATHEMATICS SPECIMEN PAPER 1

### EXPONENTIALS & LOGARITHMS REVIEW

1. Using the substitution  $u = 7^x$  in the equation  
$$7^x = 5 + 24 \times 7^{-x}$$
find the value of  $x$  correct to 4 decimal places [4]

2. Solve the equation  
$$2^{2y} - 2^{y+3} - 2^{y+2} + 32 = 0$$
by forming a quadratic equation in  $x$  where  $x = 2^y$ . [5]

3.

(a) Solve the equation

$$2 \ln x = \ln 4 + \ln(2x + 5) \quad [2]$$

(b) Solve the equation

$$e^{2x} - 3e^x = 54,$$

giving your answers correct to 3 decimal places. [3]

4. Solve the equation

$$2^y + \frac{16}{2^y} = 17 \quad [4]$$

5.

(a) Assuming that  $x = e^{\ln x}$  where  $x > 0$ , show that  $\ln x^m = m \ln x$  [3]

(b) If  $4^{(x+1)} = 7 \times 8^{(x-2)}$   
find  $x$  to 3 decimal places [6]

**Extension 1: Q6 - 7**

6.

(a) Factorise the expression  $3x^3 - 4x^2 - 5x + 2$  [4]

(b) Hence, using a suitable substitution, solve the equation  
 $3e^{3y} - 4e^{2y} - 5e^y + 2 = 0$   
giving your values of  $y$  correct to 4 decimal places [3]

7.

(a) Using the factor theorem, factorise the expression

$$2u^3 - 3u^2 - 8u + 12$$

[4]

(b) By using the substitution  $u = e^x$ , solve the equation

$$2e^x - 8e^{-x} + 12e^{-2x} = 3$$

giving your answers correct to 3 decimal places.

[3]

**SOLUTIONS:**

1.

$$7^x = 5 + 24 \times 7^{-x}$$

$$\Rightarrow 7^x = 5 + 24 \times \frac{1}{7^x}$$

Let  $u = 7^x$

$$\Rightarrow u = 5 + 24 \times \frac{1}{u}$$

$$\times u \Rightarrow u^2 = 5u + 24$$

$$\Rightarrow u^2 - 5u - 24 = 0$$

$$(u - 8)(u + 3) = 0$$

$$\Rightarrow u - 8 = 0 \text{ or } u + 3 = 0$$

$$\Rightarrow u = 8 \text{ or } u = -3$$

$$\Rightarrow 7^x = 8 \text{ or } 7^x = -3$$

Take logarithms of both sides

$$\ln 7^x = \ln 8 \quad \text{no solution}$$

$$x \ln 7 = \ln 8$$

$$x = \frac{\ln 8}{\ln 7}$$

$$x = 1.0686 \text{ to 4 decimal places}$$


---

2.

$$2^{2y} - 2^{y+3} - 2^{y+2} + 32 = 0$$

$$(2^y)^2 - 2^y \times 2^3 - 2^y \times 2^2 + 32 = 0$$

$$(2^y)^2 - 8 \times 2^y - 4 \times 2^y + 32 = 0$$

$$(2^y)^2 - 12 \times 2^y + 32 = 0$$

Let  $x = 2^y$

$$x^2 - 12x + 32 = 0$$

$$(x - 8)(x - 4) = 0$$

$$x - 8 = 0 \quad \text{or} \quad x - 4 = 0$$

$$x = 8 \quad \text{or} \quad x = 4$$

$$\Rightarrow 2^y = 8 \quad \text{or} \quad 2^y = 4$$

$$2^y = 2^3 \quad \text{or} \quad 2^y = 2^2$$

$$\Rightarrow y = 3 \quad \text{or} \quad y = 2$$


---

3.

(a)  $2 \ln x = \ln 4 + \ln(2x + 5)$

$$\Rightarrow \ln x^2 = \ln(4(2x + 5))$$

$$\Rightarrow x^2 = 4(2x + 5)$$

$$\Rightarrow x^2 = 8x + 20$$

$$x^2 - 8x - 20 = 0$$

$$(x - 10)(x + 2) = 0$$

$$\Rightarrow x - 10 = 0 \text{ or } x + 2 = 0$$

$$x = 10 \text{ or } x = -2$$

As  $x > 0$  then  $x = 10$

(b)

$$e^{2x} - 3e^x = 54$$

$$(e^x)^2 - 3(e^x) = 54$$

Let  $u = e^x$

$$\Rightarrow u^2 - 3u = 54$$

$$u^2 - 3u - 54 = 0$$

$$(u - 9)(u + 6) = 0$$

$$\Rightarrow u - 9 = 0 \text{ or } u + 6 = 0$$

$$u = 9 \text{ or } u = -6$$

$$\Rightarrow e^x = 9 \text{ or } e^x = -6$$

$$x = \ln 9 \quad \text{No solution}$$

$$x = 2.197 \text{ to 3 decimal places}$$


---

4.

$$2^y + \frac{16}{2^y} = 17$$

$$\times 2^y \quad (2^y)^2 + 16 = 17 \times 2^y$$

$$(2^y)^2 - 17 \times 2^y + 16 = 0$$

Let  $x = 2^y$

$$x^2 - 17x + 16 = 0$$

$$(x - 1)(x - 16) = 0$$

$x - 1 = 0$	or	$x - 16 = 0$
$x = 1$	or	$x = 16$
$2^y = 1$	or	$2^y = 16$
$2^y = 2^0$	or	$2^y = 2^4$
$\Rightarrow y = 0$	or	$y = 4$

---

5.

(a)

$$x = e^{\ln x} \quad \text{-\{*\}}$$

$$\Rightarrow x^m = (e^{\ln x})^m$$

$$\Rightarrow x^m = e^{(m \ln x)}$$

But, using \{\*\}

$$x^m = e^{\ln x^m}$$

$$\Rightarrow \ln x^m = m \ln x$$


---

(b)

$$4^{(x+1)} = 7 \times 8^{(x-2)}$$

Take logarithms of both sides

$$\ln 4^{(x+1)} = \ln (7 \times 8^{(x-2)})$$

$$\Rightarrow (x+1) \ln 4 = \ln 7 + \ln 8^{(x-2)}$$

$$\Rightarrow (x+1) \ln 4 = \ln 7 + (x-2) \ln 8$$

$$\Rightarrow x \ln 4 + \ln 4 = \ln 7 + x \ln 8 - 2 \ln 8$$

Re-arranging

$$2 \ln 8 + \ln 4 - \ln 7 = x \ln 8 - x \ln 4$$

$$\ln 8^2 + \ln 4 - \ln 7 = x(\ln 8 - \ln 4)$$

$$\Rightarrow \ln \left( \frac{64 \times 4}{7} \right) = x \ln \left( \frac{8}{4} \right)$$

$$\begin{aligned} \Rightarrow \ln\left(\frac{256}{7}\right) &= x \ln 2 \\ \Rightarrow x &= \frac{\ln\left(\frac{256}{7}\right)}{\ln 2} \\ \Rightarrow x &= 5.193 \text{ to 3 decimal places} \end{aligned}$$


---

6.

(a) Let  $f(x) = 3x^3 - 4x^2 - 5x + 2$   
Using the factor theorem  
 $x = 1 \Rightarrow f(1) = 3(1)^3 - 4(1)^2 - 5(1) + 2$   
 $= -4$

 $\Rightarrow (x - 1)$  is not a factor

$$x = -1 \Rightarrow f(-1) = 3(-1)^3 - 4(-1)^2 - 5(-1) + 2$$

$$= 0$$

 $\Rightarrow (x + 1)$  is a factor of  $f(x)$ To find the other factor divide  $(x + 1)$  into  $f(x)$ 

$$\begin{array}{r} \phantom{(x+1)} \overline{3x^2 - 7x + 2} \\ (x+1) \overline{) 3x^3 - 4x^2 - 5x + 2} \\ \underline{3x^3 + 3x^2} \phantom{+ 2} \\ \phantom{(x+1)} \underline{-7x^2 - 5x} \phantom{+ 2} \\ \phantom{(x+1)} \underline{-7x^2 - 7x} \phantom{+ 2} \\ \phantom{(x+1)} \phantom{-7x^2} \underline{2x + 2} \\ \phantom{(x+1)} \phantom{-7x^2} \underline{2x + 2} \\ \phantom{(x+1)} \phantom{-7x^2} \phantom{2x} \underline{-} \end{array}$$

$$\begin{aligned} \Rightarrow f(x) &= (x + 1)(3x^2 - 7x + 2) \\ &= (x + 1)(3x - 1)(x - 2) \\ \Rightarrow 3x^3 - 4x^2 - 5x + 2 &= (x + 1)(3x - 1)(x - 2) \end{aligned}$$


---

(b)  $3e^{3y} - 4e^{2y} - 5e^y + 2 = 0$

$$\Rightarrow 3(e^y)^3 - 4(e^y)^2 - 5(e^y) + 2 = 0$$

Let  $x = e^y$ 

$$3x^3 - 4x^2 - 5x + 2 = 0$$

$$\Rightarrow (x + 1)(3x - 1)(x - 2) = 0$$

$$\Rightarrow (x + 1) = 0 \text{ or } (3x - 1) = 0 \text{ or } (x - 2) = 0$$

$$x = -1 \text{ or } x = \frac{1}{3} \text{ or } x = 2$$

$$\Rightarrow e^y = -1 \quad e^y = \frac{1}{3} \quad e^y = 2$$

No solution  $y = \ln\left(\frac{1}{3}\right)$  or  $y = \ln 2$   
 $y = -1.0986$  or  $y = 0.6931$  to 4 decimal places

---

7.

(a) Let  $f(u) = 2u^3 - 3u^2 - 8u + 12$

$$u = 1 \Rightarrow f(1) = 2(1)^3 - 3(1)^2 - 8(1) + 12 = 3$$

$\Rightarrow (u - 1)$  is not a factor

$$u = -1 \Rightarrow f(-1) = 2(-1)^3 - 3(-1)^2 - 8(-1) + 12 = 15$$

$\Rightarrow (u + 1)$  is not a factor

$$u = 2 \Rightarrow f(2) = 2(2)^3 - 3(2)^2 - 8(2) + 12 = 0$$

$\Rightarrow (u - 2)$  is a factor

---

To find the other factor, divide  $(u - 2)$  into  $f(u)$

$$\begin{array}{r} \phantom{(u-2)} \overline{2u^2 + u - 6} \\ (u-2) \overline{) 2u^3 - 3u^2 - 8u + 12} \\ \underline{2u^3 - 4u^2} \phantom{+ 12} \\ \phantom{2u^3 - } u^2 - 8u \phantom{+ 12} \\ \underline{u^2 - 2u} \phantom{+ 12} \\ \phantom{2u^3 - } \phantom{u^2 - } -6u + 12 \\ \underline{-6u + 12} \\ \phantom{2u^3 - } \phantom{u^2 - } \phantom{-6u + } 0 \end{array}$$

$$\Rightarrow \begin{aligned} f(u) &= (u - 2)(2u^2 + u - 6) \\ f(u) &= (u - 2)(2u - 3)(u + 2) \end{aligned}$$


---

(b)  $2e^x - 8e^{-x} + 12e^{-2x} = 3$

$$\Rightarrow 2e^x - \frac{8}{e^x} + \frac{12}{e^{2x}} = 3$$

$$\Rightarrow 2e^x - \frac{8}{e^x} + \frac{12}{(e^x)^2} = 3$$

Let  $u = e^x$

$$\Rightarrow 2u - \frac{8}{u} + \frac{12}{u^2} = 3$$

$$\times u^2 \quad 2u^3 - 8u + 12 = 3u^2$$

$$\Rightarrow 2u^3 - 3u^2 - 8u + 12 = 0$$

$$\Rightarrow (u + 2)(2u - 3)(u - 2) = 0$$

$$\Rightarrow (u + 2) = 0 \text{ or } (2u - 3) = 0 \text{ or } (u - 2) = 0$$

$$u = -2 \text{ or } u = \frac{3}{2} \text{ or } u = 2$$

$$e^x = -2 \quad e^x = \frac{3}{2} \quad e^x = 2$$

$$\text{No solution} \quad x = \ln\left(\frac{3}{2}\right) \text{ or } x = \ln 2$$

$$x = 0.405, 0.693 \text{ to 3 decimal places}$$

---