

# C.E.M. TUITION

**Student Name :** \_\_\_\_\_


**Review Topic : Trapezoidal and Simpson's Rule**

**(HSC Course - Paper 1)**

**Year 12 - 2 Unit**

**1995**

1.



The diagram represents a scale drawing of an area of parkland bounded by a chasm on one side, a fence, and a road perpendicular to the fence on the other side.

$x$	0	20	40	60	80
$f(x)$					

Perpendicular distances have been found from the road to the chasm at 20 m intervals.

All distances are in metres.

- (a) Read the distances from the diagram and complete the table.
- (b) Estimate the area of the parkland using Simpson's Rule with five function values. Answer to the nearest  $m^2$ .



2. Use the Trapezoidal Rule with five equal subintervals to estimate

$$\int_1^6 \log_e \left( \frac{1}{x} \right) dy \text{ correct to two decimal places.}$$

3. The function  $f(t)$  is defined as  $f(t) = te^{-t}$ .

- (a) Complete this table of values for  $f(t)$ , writing values of  $f(t)$  correct to 3 decimal places.


$t$	0	1	2	3	4	5
$f(t)$						

- (b) Use the Trapezoidal Rule and 6 function values to calculate an approximate area under the curve  $f(t) = te^{-t}$  between  $t = 0$  and  $t = 5$ .

- 
4. The curve  $y = \log x$  is rotated about the  $x$  axis between  $x = 1$  and  $x = 3$ . By using Simpson's Rule and three function values, calculate the volume of the solid so formed.



5.



A section of grass between a path (AB) and a curved garden bed is to be removed, and concrete to a depth of 10 cm is to be laid.

- (a) Use Simpson's Rule with 2 subintervals to calculate the area to be concreted. Answer correct to three significant figures.
- (b) Calculate the volume of concrete required to the nearest 0.1 m<sup>3</sup>.



6. Some values of a function  $\phi(x)$ , continuous over  $5 \leq x \leq 25$ , are listed in this table. Using the Trapezoidal Rule and four subintervals, estimate

$x$	5	10	15	20	25
$\phi(x)$	0.03	0.24	0.76	1.8	0.87

the value of  $\int_5^{25} \phi(x) dx$ .

$$\int_5^{25} \phi(x) dx.$$







4.  $y = \log x \quad \therefore y^2 = [\log_e x]^2$

Volume =  $\pi \int_1^3 y^2 dx = \pi \int_1^3 (\log_e x)^2 dx$

Put  $f(x) = (\log_e x)^2$

$x$	1	2	3
$f(x)$	0	0.480 453	1.206 949

Evaluate  $\int_1^3 (\log_e x)^2 dx$

Now  $\int_1^3 (\log_e x)^2 dx = \frac{1}{6}(b-a) \left[ f(a) + f(b) + 4f\left(\frac{a+b}{2}\right) \right]$

where  $a = 1, \therefore f(a) = f(1)$

$b = 3, \therefore f(b) = f(3)$

$\frac{a+b}{2} = \frac{3+1}{2} = 2$

$\therefore f\left(\frac{a+b}{2}\right) = f(2) = 0.480 453$


$(b-a) = 3 - 1 = 2$

$\therefore \int_1^3 (\log_e x)^2 dx = \frac{1}{6}(3-1)[f(1) + f(3) + 4f(2)]$   
 $= \frac{1}{3}[0 + 1.206 949 + 4(0.480 453)]$   
 $= 1.042 920 3.$

Then  $V = \pi \int_1^3 (\log_e x)^2 dx = \pi \times 1.042 920 3$   
 $= 3.276 430 9$   
 $= 3.28 \text{ (2 dec. places).}$

Volume is  $3.28 \text{ units}^3$ .

5.



Call curve  $y = f(x)$ .  
 Set up table of values.

$x$	0	4	8	12	16
$f(x)$	0	3.5	4.6	4.6	0

(a) Subintervals will be  $0 - 8$ , and  $8 - 16$ .  
 Midpoints are thus 4 and 12, i.e., if  $a = 0$  and  $b = 8$ ,

then  $\frac{a+b}{2} = \frac{0+8}{2} = 4.$

Area =  $\frac{1}{6}(8-0)[f(0) + f(8) + 4f(4)]$   
 $+ \frac{1}{6}(16-8)[f(8) + f(16) + 4f(12)]$   
 $= \frac{4}{3}[0 + 4.6 + 4(3.5)] + \frac{4}{3}[4.6 + 0 + 4(4.6)]$   
 $= 24.8 + 30.666 67$   
 $= 55.466 67.$

Area to be concreted is  $55.5 \text{ m}^2$ .

OR TODD Simpson method

$x$	0	4	8	12	16
$f(x)$	0	3.5	4.6	4.6	0

$$h = 4 - 0 = 8 - 4, \text{ etc.} \\ = 4$$

$\uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow$   
*F*            *ODD*            *L*  
*EVEN*    *EVEN*

$$\begin{aligned} \text{Area} &= \frac{h}{3} [f(F) + f(L) + 2(ODD) + 4(EVEN)] \quad (FLOE) \\ &= \frac{4}{3} [0 + 0 + 2(4.6) + 4(3.5 + 4.6)] \\ &= \frac{4}{3} [41.6] \\ &= 55.46667 \\ &= 55.5 \text{ m}^2. \end{aligned}$$

(b) Volume = Area  $\times$  depth =  $55.46667 \times 0.1$

$$= 5.546667$$

$$V = Ah$$

$$\approx 5.5 \text{ (1 dec. place).}$$

Volume of concrete required is  $5.5 \text{ m}^3$ .

6.

$x$	5	10	15	20	25
$\phi(x)$	0.03	0.24	0.76	1.8	0.87

(TREMAINING)

$$h = 10 - 5 \\ = 5$$

$$\begin{aligned} \int_5^{25} \phi(x) dx &= \frac{h}{2} [FIRST + LAST + 2(REMAINING)] \quad (FLR) \\ &= \frac{5}{2} [f(5) + f(25) + 2(f(10) + f(15) + f(20))] \\ &= 2.5 [0.03 + 0.87 + 2(0.24 + 0.76 + 1.8)] \\ &= 16.25. \end{aligned}$$

Value of integral  $\int_5^{25} \phi(x) dx$  is 16.25.