



CATHOLIC SECONDARY SCHOOLS
ASSOCIATION OF NEW SOUTH WALES

--	--	--	--	--

Centre Number

--	--	--	--	--	--	--	--	--	--

Student Number

2007
TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics

Extension 1

Afternoon Session
Tuesday 14 August 2007

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using blue or black pen
- Board-approved calculators may be used
- A table of standard integrals is provided separately
- All necessary working should be shown in every question

Total marks – 84

- Attempt Questions 1-7
- All questions are of equal value


Disclaimer
Every effort has been made to prepare these 'Trial' Higher School Certificate Examinations in accordance with the Board of Studies documents, *Principles for Setting HSC Examinations in a Standards-Referenced Framework* (BOS Bulletin, Vol 8, No 9, Nov/Dec 1999), and *Principles for Developing Marking Guidelines Examinations in a Standards Referenced Framework* (BOS Bulletin, Vol 9, No 3, May 2000). No guarantee or warranty is made or implied that the 'Trial' Examination papers mirror in every respect the actual HSC Examination question paper in any or all courses to be examined. These papers do not constitute 'advice' nor can they be construed as authoritative interpretations of Board of Studies intentions. The CSSA accepts no liability for any reliance use or purpose related to these 'Trial' question papers. Advice on HSC examination issues is only to be obtained from the NSW Board of Studies.

3201 - 1

Question 1 (12 Marks) Use a SEPARATE writing booklet

- (a) If $y = \sin^2 3x$ find $\frac{dy}{dx}$. 2
- (b) $A(-3, 1)$, $B(8, -2)$ and $C(4, 16)$ are the vertices of a triangle. AM is a median of the triangle.
- (i) Show that M has coordinates $(6, 7)$. 1
- (ii) Hence find the coordinates of the point which divides the interval AM internally in the ratio 2:1. 2
- (c)(i) Express $x^3 - 3x^2 + 4$ as a product of three linear factors by first showing that $(x-2)$ is a factor of this polynomial. 2
- (ii) Hence solve the inequality $x^3 - 3x^2 + 4 \geq 0$. 1

(d)



In the diagram, $ABCD$ is a cyclic quadrilateral. E is a point on CD produced.

- (i) Give a reason why $\angle ADE = \angle ABC$. 1
- (ii) If $\triangle ADE \parallel \triangle CBA$, show that $AE \parallel BD$. 3

Marks

Question 2 (12 Marks) Use a SEPARATE writing booklet

(a) If $y = \log_{\frac{1}{a}} \left(\frac{1}{N} \right)$, where $a > 0$ and $N > 0$, show that $y = \log_a N$. 2

(b) The lines $y = mx$ and $y = 2mx$, where $m > 0$, are inclined to each other at an angle θ such that $\tan \theta = \frac{1}{3}$.

(i) Show that $2m^2 - 3m + 1 = 0$. 1

(ii) Hence find the possible values of m . 2

(c)(i) Show that $\tan 2x + \tan x = \frac{\sin 3x}{\cos 2x \cos x}$. 2

(ii) Hence solve the equation $\tan 2x + \tan x = 0$ for $0 < x < \frac{\pi}{2}$. 1

(d) $P(2ap, ap^2)$, $Q(2aq, aq^2)$ and $R(2ar, ar^2)$, where $p < q < r$, are three points on the parabola $x^2 = 4ay$.

(i) Use differentiation to show that the tangent to the parabola at Q has gradient q . 1

(ii) If the chord PR is parallel to the tangent at Q , show that p , q and r are consecutive terms in an arithmetic sequence. 3

Marks

Question 3 (12 Marks) Use a SEPARATE writing booklet

(a) Find the number of ways in which the letters of the word EPSILON can be arranged in a straight line so that the three vowels are all next to each other. 2

(b) Use Mathematical induction to show that $5^n > 3^n + 4^n$ for all positive integers $n \geq 3$. 4

(c) Consider the function $f(x) = x + e^{-x}$, $x \geq 0$.

(i) Show that for all values of $x > 0$, the function $f(x)$ is increasing and the curve $y = f(x)$ is concave up. 2

(ii) Sketch the graph of $y = f(x)$ showing clearly the coordinates of the endpoint and the equation of the asymptote. 2

(iii) On the same diagram, sketch the graph of the inverse function $y = f^{-1}(x)$. 1

(iv) Find the domain of the function $g(x) = f(x) + f^{-1}(x)$. 1

Question 4 (12 Marks) Use a SEPARATE writing booklet


Marks

- (a)(i) Show that the equation $x^3 - 2x - 5 = 0$ has a root α such that $2 < \alpha < 3$. 2
- (ii) Use one application of Newton's method and an initial approximation of 2 to find the next approximation for α . 2
- (b)(i) Sketch the graph of $y = 2\sin^{-1}2x$ showing clearly the coordinates of the endpoints. 2
- (ii) Find the exact area of the region in the first quadrant bounded by the curve $y = 2\sin^{-1}2x$, the y -axis and the line $y = \pi$. 2
- (c)(i) Show that $\frac{u^2}{1+u^2} = 1 - \frac{1}{1+u^2}$. 1
- (ii) Use the substitution $x = u^2$, $u \geq 0$, to find $\int \frac{\sqrt{x}}{1+x} dx$. 3

Question 5 (12 Marks) Use a SEPARATE writing booklet

Marks

(a)




An egg timer in the shape of an inverted right circular cone of semi vertical angle $\frac{\pi}{6}$ contains sand to a depth of h cm. The sand flows out of the apex of the cone at a constant rate of $0.5 \text{ cm}^3/\text{s}$.

- (i) Show that the volume $V \text{ cm}^3$ of sand in the cone is given by $V = \frac{1}{9}\pi h^3$. 1
- (ii) Find the value of h when the depth of sand in the egg timer is decreasing at a rate of 0.05 cm/s , giving your answer correct to 2 decimal places. 3
- (b) A particle is moving in a straight line. At time t seconds it has displacement x metres from a fixed point O on the line, velocity $v \text{ ms}^{-1}$ given by $v = -\frac{1}{8}x^3$, and acceleration $a \text{ ms}^{-2}$. The particle is initially 2 metres to the right of O .
- (i) Show that $a = \frac{3}{64}x^5$. 1
- (ii) Find an expression for x in terms of t . 2
- (iii) Describe the limiting motion of the particle. 1
- (c) A game is played by throwing three fair coins. The game is considered a failure if all three coins show heads or all three coins show tails. Otherwise the game is considered a success.
- (i) Show that the probability of success in any play of the game is $\frac{3}{4}$. 2
- (ii) Find the probability of exactly two successes in four plays of the game. 2

Question 6 (12 Marks) Use a SEPARATE writing booklet

- (a) At time t years the area A km² controlled by a colony of animals is given by $A = 300 - 200e^{-kt}$ for some $k > 0$.
- (i) Sketch the graph of A as a function of t showing clearly the initial and limiting areas controlled by the colony. 2
- (ii) Find the value of k if the area controlled by the colony is increasing at a rate of 10 km² per year when this area is twice its initial value. 2



A particle is projected from a point O with speed V ms⁻¹ at an angle θ above the horizontal, where $\frac{\pi}{4} < \theta < \frac{\pi}{2}$. It moves in a vertical plane under gravity where the acceleration due to gravity is g ms⁻². At time t seconds its horizontal and vertical displacements from O are x metres and y metres respectively.

- (i) Use integration to show that $x = Vt \cos \theta$ and $y = Vt \sin \theta - \frac{1}{2}gt^2$. 2
- (ii) The particle takes T seconds to reach its greatest height H metres. Find expressions for T and H in terms of g , V and θ . 2
- (iii) The path of the particle is inclined at an angle $\frac{\pi}{4}$ above the horizontal at time $\frac{1}{4}T$. Show that $\theta = \tan^{-1} \frac{4}{3}$. 2
- (iv) What fraction of its maximum height has the particle attained at time $\frac{1}{4}T$? 2

Marks

Question 7 (12 Marks) Use a SEPARATE writing booklet

- (a) A particle is moving in a straight line with Simple Harmonic motion. At time t seconds it has displacement x metres from a fixed point O on the line, where $x = (\cos t + \sin t)^2$, velocity v ms⁻¹ and acceleration a ms⁻².
- (i) Show that $a = -4(x-1)$. 2
- (ii) Find the extreme positions of the particle during its motion. 1
- (iii) Find the time taken by the particle to move from one extreme position of its motion to the other extreme position. 1
- (iv) A second particle moves along a parallel straight line with Simple Harmonic motion so that $x = 1 - \cos t$, where x metres is its displacement from a fixed point level with O . During the first complete oscillation of the particle with the slower average speed, find the number of times the particles pass each other, and state their relative directions of travel on each occasion. 3

- (b) (i) Considering the identity $(1-t)^n(1+t)^n = (1-t^2)^n$, where n is a positive integer, show that for integer values of r , 2
- $$\sum_{k=0}^{2r} (-1)^k {}^nC_k {}^nC_{2r-k} = (-1)^r {}^nC_r \text{ provided } 0 \leq r \leq \frac{1}{2}n.$$
- (ii) Hence show that $\sum_{k=0}^r (-1)^k {}^nC_k {}^nC_{2r-k} = \frac{1}{2}(-1)^r {}^nC_r \{1 + {}^nC_r\}$ for $0 \leq r \leq \frac{1}{2}n$. 2
- (iii) Hence evaluate $\sum_{k=0}^{10} (-1)^k ({}^nC_k)^2$ as a basic numeral. 1

Examiners

Graham Arnold
Denise Arnold
Dianne Williams
Franklin Samyia

Terra Sancta College, Quakers Hill
Patrician Brothers' College, Blacktown
St Andrews College, Marayong
Holy Cross College, Ryde

Marks

But $\sum_{k=0}^{2r} (-1)^k {}^nC_k {}^nC_{2r-k} = (-1)^r {}^nC_r$ for $0 \leq r \leq \frac{1}{2}n$.

$\therefore (-1)^r {}^nC_r = 2 \sum_{k=0}^r (-1)^k {}^nC_k {}^nC_{2r-k} - (-1)^r {}^nC_r {}^nC_r$

$(-1)^r {}^nC_r + (-1)^r {}^nC_r {}^nC_r = 2 \sum_{k=0}^r (-1)^k {}^nC_k {}^nC_{2r-k}$

$\therefore \sum_{k=0}^r (-1)^k {}^nC_k {}^nC_{2r-k} = \frac{1}{2} (-1)^r {}^nC_r \{1 + {}^nC_r\}$ for $0 \leq r \leq \frac{1}{2}n$.

iii. Substituting $n=20$ and $r=10$ gives $\sum_{k=0}^{10} (-1)^k {}^{20}C_k {}^{20}C_{20-k} = \frac{1}{2} (-1)^{10} {}^{20}C_{10} \{1 + {}^{20}C_{10}\}$ since

$10 \leq \frac{1}{2} \times 20$.

But ${}^{20}C_k = {}^{20}C_{20-k}$ for $k=1, 2, \dots, 20$.

Hence

$\sum_{k=0}^{10} (-1)^k ({}^{20}C_k)^2 = \frac{1}{2} (-1)^{10} {}^{20}C_{10} \{1 + {}^{20}C_{10}\}$

$= \frac{1}{2} \times 184756 \times 184757$

$= 17067482146$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.



CATHOLIC SECONDARY SCHOOLS ASSOCIATION
2007 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION
MATHEMATICS EXTENSION 1

Question 1

a. Outcomes assessed : H5, PE5

Marking Guidelines

Criteria	Marks
• applies the chain rule, writing one factor of the derivative as $2\sin 3x$	1
• obtains the second factor $3\cos 3x$ (even if final simplification is not carried out)	1

Answer

$y = \sin^2 3x \quad \therefore \frac{dy}{dx} = 2\sin 3x \cdot 3\cos 3x = 3(2\sin 3x \cos 3x) = 3\sin 6x$

b. Outcomes assessed : H5


Marking Guidelines

Criteria	Marks
i • identifies M as the midpoint of BC and obtains its coordinates	1
ii • finds the x coordinate of the division point	1
• finds the y coordinate of the division point	1

Answer

i. M is the midpoint of BC . Hence M has coordinates $(\frac{8+4}{2}, \frac{-2+16}{2}) = (6, 7)$

ii.



Required point has coordinates $(3, 5)$

$(\frac{2 \times 6 + 1 \times (-3)}{2+1}, \frac{2 \times 7 + 1 \times 1}{2+1})$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

c. Outcomes assessed : PE3

Marking Guidelines

Criteria	Marks
i • shows that $(x-2)$ is a factor	1
• completes the factorisation	1
ii • solves the inequality	1

Answer

i. Let $P(x) = x^3 - 3x^2 + 4$. Then $P(2) = 8 - 12 + 4 = 0 \therefore (x-2)$ is a factor of $P(x)$

Then by inspection (or by division) $x^3 - 3x^2 + 4 = (x-2)(x^2 - x - 2) = (x-2)^2(x+1)$

ii. Since $(x-2)^2 \geq 0$ for all real x , $x^3 - 3x^2 + 4 \geq 0$ whenever $x+1 \geq 0 \therefore x \geq -1$

d. Outcomes assessed : PE2, PE3

Marking Guidelines

Criteria	Marks
i • quotes an appropriate geometrical property as a reason	1
ii • uses the similarity to identify a pair of equal angles	1
• identifies equal alternate or corresponding angles using an appropriate circle property	1
• quotes an appropriate test for parallel lines to support the required deduction	1

Answer

i. In cyclic quadrilateral $ABCD$, the exterior angle ADE is equal to the interior opposite angle ABC .

ii. If $\triangle ADE \parallel \triangle CBA$, then $\angle AED = \angle CAB$ (corresponding \angle 's in similar triangles are equal)

But $\angle CAB = \angle CDB$ (\angle 's subtended at the circumference by the same arc BC are equal)

$\therefore \angle AED = \angle BDC$ (both equal to $\angle CAB$)

$\therefore AE \parallel BD$ (equal corresponding angles on transversal CDE)

Question 2

a. Outcomes assessed : H3

Marking Guidelines

Criteria	Marks
• converts the logarithm statement to an equivalent index statement	1
• after taking reciprocals, converts the index statement to an equivalent logarithm statement	1

Answer

The statement $y = \log_{\frac{1}{a}} \left(\frac{1}{N} \right)$ is equivalent to $\left(\frac{1}{a} \right)^y = \frac{1}{N}$.

Taking reciprocals, $a^y = N \therefore y = \log_a N$

DISCLAIMER
The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

b. Outcomes assessed : P4, H5

Marking Guidelines

Criteria	Marks
i • uses the formula for the angle between two lines to obtain result	1
ii • factors quadratic expression	1
• writes down both possible values of m	1

Answer

i. θ is the acute angle between lines with gradients m and $2m$, where $m > 0$.

$$\therefore \frac{|2m - m|}{|1 + 2m^2|} = \tan \theta \Rightarrow \frac{m}{1 + 2m^2} = \frac{1}{3}. \quad \text{Hence } 2m^2 - 3m + 1 = 0.$$

ii. $(2m-1)(m-1) = 0 \therefore m = \frac{1}{2}, 1$

c. Outcomes assessed : H5

Marking Guidelines

Criteria	Marks
i • writes expressions for \tan in terms of \sin and \cos	1
• takes a common denominator and recognises the expression for sine of an angle sum	1
ii • solves $\sin 3x = 0$ in the required domain	1

Answer

i.

$$\begin{aligned} \tan 2x + \tan x &= \frac{\sin 2x}{\cos 2x} + \frac{\sin x}{\cos x} \\ &= \frac{\sin 2x \cos x + \cos 2x \sin x}{\cos 2x \cos x} \\ &= \frac{\sin 3x}{\cos 2x \cos x} \end{aligned}$$

ii. $\sin 3x = 0, 0 < 3x < \frac{3\pi}{2} \Rightarrow 3x = \pi$
Hence $\tan 2x + \tan x = 0, 0 < x < \frac{\pi}{2}$
has solution $x = \frac{\pi}{3}$

d. Outcomes assessed : PE3, PE4, H5

Marking Guidelines

Criteria	Marks
i • uses direct or parametric differentiation to establish result	1
ii • finds simplified expression for gradient of PR	1
• equates gradients of parallel lines	1
• rearranges relation to show p, q, r in arithmetic progression	1

Answer

i. $y = \frac{1}{4a} x^2$

$\therefore \frac{dy}{dx} = \frac{1}{2a} x = q$ at Q

Hence tangent at Q has gradient q .

DISCLAIMER
The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

$$\text{ii. gradient } PR = \frac{a(r^2 - p^2)}{2a(r-p)} = \frac{r+p}{2}$$

If PR is parallel to tangent at Q

$$\frac{r+p}{2} = q$$

$$r+p = 2q$$

$$r-q = q-p$$

$\therefore p, q, r$ are in arithmetic progression

Question 3

a. Outcomes assessed : PE3

Marking Guidelines

Criteria	Marks
• groups vowels and arranges 5 objects	1
• multiplies by number of arrangements of the 3 vowels	1

Answer

(EIO), P, S, L, N arranged in $5!$ ways, then E, I, O in $3!$ ways. Hence $5! \times 3! = 720$ arrangements.

b. Outcomes assessed : HE2

Marking Guidelines

Criteria	Marks
• defines a sequence of statements and establishes the truth of the first $5^3 > 3^3 + 4^3$	1
• writes an inequality for 5^{k+1} in terms of $3^k + 4^k$, conditional on the truth of $S(k)$	1
• works with this inequality to show that $5^{k+1} > 3^{k+1} + 4^{k+1}$	1
• deduces the required result from the previous steps	1

Answer

Define the sequence of statements $S(n): 5^n > 3^n + 4^n$, $n = 3, 4, 5, \dots$

Consider $S(3)$: $5^3 = 125$, $3^3 + 4^3 = 91 \Rightarrow 5^3 > 3^3 + 4^3 \quad \therefore S(3)$ is true.

If $S(k)$ is true: $5^k > 3^k + 4^k$ **

Consider $S(k+1)$: $5^{k+1} = 5 \cdot 5^k$

$$> 5(3^k + 4^k) \quad \text{if } S(k) \text{ is true, using **}$$

$$= 5 \cdot 3^k + 5 \cdot 4^k$$

$$> 3 \cdot 3^k + 4 \cdot 4^k$$

$$= 3^{k+1} + 4^{k+1}$$

Hence if $S(k)$ is true, then $S(k+1)$ is true. But $S(3)$ is true, hence $S(4)$ is true, and then $S(5)$ is true and so on. Hence by Mathematical induction, $5^n > 3^n + 4^n$ for all positive integers $n \geq 3$.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

c. Outcomes assessed : H5, HE4

Marking Guidelines

Criteria	Marks
i • uses the first derivative to show the function is increasing	1
• uses the second derivative to show the curve is concave up	1
ii • sketches curve with correct shape showing the endpoint $(0, 1)$	1
• shows the oblique asymptote with equation $y = x$	1
iii • sketches the inverse as the reflection in the line $y = x$	1
iv • writes the domain of the function g	1

Answer

ii., iii.


$$\text{i. } f(x) = x + e^{-x}, x \geq 0$$

$$f'(x) = 1 - e^{-x} > 0 \text{ for } x > 0$$

Hence function is increasing for $x > 0$.

$$f''(x) = e^{-x} > 0 \text{ for } x > 0$$

Hence the curve is concave up for $x > 0$.



iv. Domain of g is the intersection of the domains of f and f^{-1} .

$$\therefore g \text{ has domain } \{x : x \geq 1\}.$$

Question 4

a. Outcomes assessed : PE3

Marking Guidelines

Criteria	Marks
i • shows that expression is negative for $x = 2$ and positive for $x = 3$.	1
• uses the continuity of the function to deduce equation has a root between 2 and 3	1
ii • calculates the value of the derivative at $x = 2$	1
• applies Newton's method once to calculate the next approximation	1

Answer

$$\text{i. Let } f(x) = x^3 - 2x - 5$$

$$\text{Then } f(2) = 8 - 4 - 5 = -1 < 0$$

$$\text{and } f(3) = 27 - 6 - 5 = 16 > 0$$

But f is continuous. Hence there exists some real number α , $2 < \alpha < 3$, such that $f(\alpha) = 0$.

$$\text{ii. } f'(x) = 3x^2 - 2 \Rightarrow f'(2) = 10$$

Using Newton's method, next approximation for the root α is

$$2 - \frac{f(2)}{f'(2)} = 2 - \frac{-1}{10} = 2.1$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.


b. Outcomes assessed : HE4

Marking Guidelines

Criteria	Marks
i • sketches curve with correct shape and position	1
• shows the coordinates of the endpoints	1
ii • writes an expression for the area as the definite integral of a function of y	1
• finds the primitive then evaluates the definite integral	1

Answer

i.



ii. Area is A square units where $A = \int_0^\pi x \, dy$.

$$\begin{aligned} A &= \int_0^\pi \frac{1}{2} \sin\left(\frac{1}{2}y\right) \, dy \\ &= -\left[\cos\left(\frac{1}{2}y\right)\right]_0^\pi \\ &= -(\cos\frac{\pi}{2} - \cos 0) \\ &= 1 \end{aligned}$$

Hence area is 1 square unit.

c. Outcomes assessed : P4, HE6

Marking Guidelines

Criteria	Marks
i • shows the required result	1
ii • writes du in terms of dx and writes integrand as a function of u	1
• finds primitive as a function of u	1
• substitutes for u in terms of x	1

Answer

i.

$$\begin{aligned} \frac{u^2}{1+u^2} &= \frac{(1+u^2)-1}{1+u^2} \\ &= 1 - \frac{1}{1+u^2} \end{aligned}$$

ii. $x = u^2 \quad u > 0$
 $dx = 2u \, du$

$$\begin{aligned} \int \frac{\sqrt{x}}{1+x} \, dx &= \int \frac{u}{1+u^2} \cdot 2u \, du \\ &= 2 \int \frac{u^2}{1+u^2} \, du \\ &= 2 \int \left(1 - \frac{1}{1+u^2}\right) \, du \\ &= 2(u - \tan^{-1}u) + c \\ &= 2(\sqrt{x} - \tan^{-1}\sqrt{x}) + c \end{aligned}$$

DISCLAIMER
 The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 5

a. Outcomes assessed : P4, HE5, HE7

Marking Guidelines

Criteria	Marks
i • expresses the radius of the cone of sand in terms of h then uses $V = \frac{1}{3}\pi r^2 h$	1
ii • expresses $\frac{dV}{dt}$ in terms of $\frac{dh}{dt}$	1
• substitutes appropriate negative values for both of these derivatives	1
• calculates depth to required accuracy	1

Answer

i. The cone of sand has radius $h \tan \frac{\pi}{6}$

$$\begin{aligned} \therefore V &= \frac{1}{3}\pi \left(\frac{1}{\sqrt{3}}h\right)^2 h \\ &= \frac{1}{9}\pi h^3 \end{aligned}$$

ii. $\frac{dV}{dt} = \frac{1}{9}\pi \cdot 3h^2 \frac{dh}{dt}$
 $-0.5 = \frac{1}{3}\pi h^2 (-0.05)$

$$\begin{aligned} \frac{3 \times 0.5}{0.05 \times \pi} &= h^2 \\ h &= \sqrt{\frac{30}{\pi}} \end{aligned}$$

Depth is 3.09 cm (to 2 dec. pl.)

b. Outcomes assessed : HE5

Marking Guidelines

Criteria	Marks
i • writes a in terms of either $\frac{dv}{dx}$ or $\frac{dv^2}{dx}$ to obtain required result	1
ii • integrates $\frac{dt}{dx}$ and evaluates the constant to find t in terms of x	1
• rearranges to find x in terms of t , choosing the appropriate square root	1
iii • describes the limiting position, speed and acceleration as $t \rightarrow \infty$	1

Answer

i. $a = v \frac{dv}{dx} = -\frac{1}{8}x^3 \cdot \left(-\frac{3}{8}x^2\right)$
 $\therefore a = \frac{3}{64}x^5$

ii. $\frac{dx}{dt} = -\frac{1}{8}x^3 \quad t=0 \Rightarrow c=-1$
 $\frac{dx}{dt} = -8x^{-3} \quad x=2 \Rightarrow t = \frac{4}{x^2} - 1$
 $t+1 = \frac{4}{x^2}$
 $\therefore x^2 = \frac{4}{t+1}$ and $x > 0$ for $t > 0$
 $\therefore x = \frac{2}{\sqrt{t+1}}$

iii. As $t \rightarrow \infty$, the particle is moving left, approaching O and slowing down at an ever decreasing rate with speed approaching 0.

DISCLAIMER
 The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

c. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • writes an expression for the probability of success in terms of outcomes	1
• shows how this probability is calculated	1
ii • recognises this as a binomial probability and writes corresponding numerical expression	1
• calculates this probability	1

Answer

$$i. P(\text{success}) = 1 - \{P(H, H, H) + P(T, T, T)\}$$

$$= 1 - \left\{ \left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^3 \right\}$$

$$= \frac{3}{4}$$

$$ii. P(\text{exactly 2 successes}) = {}^4C_2 \left(\frac{3}{4}\right)^2 \left(\frac{1}{4}\right)$$

$$= \frac{27}{128}$$


Question 6

a. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • sketches curve with correct shape and vertical intercept 100	1
• shows horizontal asymptote $A = 300$	1
ii • expresses $\frac{dA}{dt}$ in terms of A and k	1
• writes and solves an equation for k	1

Answer



DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

b. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • uses integration to show result for x	1
• uses integration to show result for y	1
ii • finds expression for T	1
• find expression for H	1
iii • finds \dot{x} and \dot{y} when $t = \frac{1}{4}T$	1
• deduces equality of horizontal and vertical components of velocity at this time to find θ	1
iv • finds y when $t = \frac{1}{4}T$	1
• expresses this y value as a fraction of H	1

Answer

i. $\ddot{x} = 0$ $\ddot{y} = -g$

x is constant $\therefore \dot{x} = V \cos \theta$ $\dot{y} = -gt + c_2$

$$x = Vt \cos \theta + c_1$$

$$t = 0 \Rightarrow c_1 = 0 \quad \therefore x = Vt \cos \theta$$

$$y = Vt \sin \theta - \frac{1}{2}gt^2 + c_3$$

$$t = 0 \Rightarrow c_3 = 0 \quad \therefore y = Vt \sin \theta - \frac{1}{2}gt^2$$

ii. At greatest height, $\dot{y} = 0$, $y = H$

$$\therefore gT = V \sin \theta$$

$$T = \frac{V \sin \theta}{g}$$


$$H = V \left(\frac{V \sin \theta}{g} \right) \sin \theta - \frac{1}{2}g \left(\frac{V \sin \theta}{g} \right)^2$$

$$= \frac{V^2 \sin^2 \theta}{2g}$$

iii. When $t = \frac{1}{4}T$,

$$\dot{y} = \frac{3}{4}V \sin \theta$$

$$\dot{x} = V \cos \theta$$



$$\therefore \dot{y} = \dot{x}$$

$$\frac{3}{4} \sin \theta = \cos \theta$$

$$\tan \theta = \frac{4}{3}$$

$$\therefore \theta = \tan^{-1} \frac{4}{3}$$

iv. When $t = \frac{1}{4}T$, $y = V \left(\frac{V \sin \theta}{4g} \right) \sin \theta - \frac{1}{2}g \left(\frac{V \sin \theta}{4g} \right)^2 = \frac{7V^2 \sin^2 \theta}{32g} = \frac{7}{16}H$

Hence particle has attained $\frac{7}{16}$ of its maximum height.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

Question 7

a. Outcomes assessed : HE3

Marking Guidelines

Criteria	Marks
i • uses trigonometric identities to simplify expression for x	1
• finds second derivative then rearranges to obtain required result	1
ii • states both extreme positions	1
iii • states time taken	1
iv • realises that the second particle with period 2π seconds has the slower average speed	1
• graphs x as a function of t for both particles on the same axes and counts the intersections (or solves simultaneous trigonometric equations) for $0 \leq t \leq 2\pi$	1
• investigates the signs of the gradients of the graphs at the intersection points (or the values of v at the solutions of the corresponding trigonometric equation) and relates these to the relative directions of travel	1

Answer

i. $x = (\cos t + \sin t)^2$
 $= \cos^2 t + \sin^2 t + 2 \sin t \cos t$
 $= 1 + \sin 2t$

$v = 2 \cos 2t$


$a = -4 \sin 2t$

$= -4(x - 1)$

ii. $-1 \leq \sin 2t \leq 1$
 $\therefore 0 \leq x \leq 2$, where x attains both extremes.
 Extreme positions are O and the point 2 m to right of O .

iii. Period of motion is $\frac{2\pi}{2} = \pi$ seconds.
 Hence time taken is $\frac{\pi}{2}$ seconds.

iv. The second particle moves between extreme positions where $x = 0$ and $x = 2$, starting at $x = 0$ and taking 2π seconds for each complete oscillation, and hence has the slower average speed.



The 4 intersection points correspond to 4 times when the particles pass each other while the second particle completes its first oscillation.

The gradient of the curve gives the velocity of the corresponding particle, and hence the sign of the gradient gives the direction of the particle.

Hence on the first, second and fourth occasions that they pass each other they are travelling in opposite directions, but on the third occasion they are travelling in the same direction.

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.

b. Outcomes assessed : PE6, HE3

Marking Guidelines

Criteria	Marks
i • applies Binomial expansions to $(1-t)^n, (1+t)^n$ and $(1-t^2)^n$	1
• equates coefficients of t^{2r} on both sides of the identity to obtain required result	1
ii • realises that in the summation in (i), the terms given by $k = r - j$ and $k = r + j$ are the same for $j = 1, 2, 3, \dots, r$.	1
• uses this fact with appropriate rearrangement and regrouping to obtain required result	1
iii • substitutes appropriate values for n and r then uses (ii) to calculate given sum.	1

Answer

i. $(1-t)^n(1+t)^n \equiv (1-t^2)^n$ ** Using the binomial expansion for each factor on the LHS,
 $(1-t)^n = 1 - {}^n C_1 t + {}^n C_2 t^2 - {}^n C_3 t^3 + \dots + (-1)^k {}^n C_k t^k + \dots + (-1)^n {}^n C_n t^n$
 $(1+t)^n = 1 + {}^n C_1 t + {}^n C_2 t^2 + {}^n C_3 t^3 + \dots + {}^n C_k t^k + \dots + {}^n C_n t^n$

The term in t^{2r} on the LHS of ** is formed by adding the products of terms $(-1)^k {}^n C_k t^k$ and ${}^n C_{2r-k} t^{2r-k}$ taken from the first and second expansions respectively in all possible ways. Provided $2r \leq n$, k can take all the values $0, 1, 2, \dots, 2r$ when forming this product,

giving the coefficient of t^{2r} as $\sum_{k=0}^{2r} (-1)^k {}^n C_k {}^n C_{2r-k}$.

The binomial expansion of the RHS of ** gives the coefficient of t^{2r} as $(-1)^r {}^n C_r$.

Hence, provided $0 \leq r \leq \frac{1}{2}n$, equating coefficients of t^{2r} on both sides of the identity ** gives

$$\sum_{k=0}^{2r} (-1)^k {}^n C_k {}^n C_{2r-k} = (-1)^r {}^n C_r.$$

ii. For $j = 1, 2, 3, \dots, r$, $(-1)^{r-j} {}^n C_{r-j} {}^n C_{2r-(r-j)} = (-1)^r (-1)^{-j} {}^n C_{r-j} {}^n C_{r+j}$
 $= (-1)^r (-1)^j {}^n C_{r-j} {}^n C_{r+j}$
 $= (-1)^{r+j} {}^n C_{r+j} {}^n C_{r-j}$
 $= (-1)^{r+j} {}^n C_{r+j} {}^n C_{2r-(r+j)}$

Hence the terms in the sum obtained by putting $k = r - j$ and $k = r + j$ are equal.

$$\begin{aligned} \therefore \sum_{k=0}^{2r} (-1)^k {}^n C_k {}^n C_{2r-k} &= \sum_{k=0}^{r-1} (-1)^k {}^n C_k {}^n C_{2r-k} + (-1)^r {}^n C_r {}^n C_r + \sum_{k=r+1}^{2r} (-1)^k {}^n C_k {}^n C_{2r-k} \\ &= 2 \sum_{k=0}^{r-1} (-1)^k {}^n C_k {}^n C_{2r-k} + (-1)^r {}^n C_r {}^n C_r \\ &= 2 \sum_{k=0}^{r-1} (-1)^k {}^n C_k {}^n C_{2r-k} - (-1)^r {}^n C_r {}^n C_r \end{aligned}$$

DISCLAIMER

The information contained in this document is intended for the professional assistance of teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible Trial HSC answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Board of Studies. No guarantee nor warranty is made or implied with respect to the application or use of CSSA Marking Guidelines in relation to any specific trial exam question or answer. The CSSA assumes no liability nor responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the Trial HSC papers.