

Name: _____

Teacher: _____



DANE BANK

An Anglican School for Girls

Year 12 Mathematics Extension 1 Task # 2 June, 2006

Time Allowed – 50 minutes

(This examination paper does not necessarily reflect the content or format of the final Higher School Certificate Examination Paper for this subject)

Weight: 30%

Outcomes examined: PE3, PE5, PE6, H5, H8, H9, HE4, HE6, HE7

DIRECTIONS TO CANDIDATE:

- Attempt all questions.
- Start each new question on a new page, using the supplied paper.
- Show all necessary working otherwise full marks may not be awarded
- Marks may be deducted for careless or badly arranged work.
- All questions are NOT of equal value.
- Board approved calculators may be used.
- Write your name on this paper.
- A table of standard integrals is provided on the last page.

Question 1 (15 Marks)

Marks

- (a) Find the exact value of $\cos[\sin^{-1}(\frac{12}{13})]$

2

- (b) Differentiate: $\cos^{-1}\left(\frac{1}{\sqrt{x}}\right)$

3

- (c) Find: $\int \frac{dx}{\sqrt{9-x^2}}$


2

- (d) Evaluate: $\int_0^4 \frac{dx}{16+x^2}$

3

- (e) Write down the equation for the following inverse trig function:

2



- (f) Find the value of "a" if

$$\int_{-a}^a \sec^2 2x \, dx = 1$$

3

Question 2 (11 Marks)

Marks

- (a) $\int \frac{e^x}{1+e^{2x}} \, dx$ is equal to


1

A. $\log_e(1+e^{2x}) + c$ B. $\frac{1}{2} \log_e(1+e^{2x}) + c$

C. $\tan^{-1} e^{2x} + c$ D. $\tan^{-1} e^x + c$

- (b) Draw the inverse function of the function shown on the SEPARATE ANSWER SHEET on the same graph.

2



Use Simpson's Rule with 3 function values to find an approximation to the shaded area.

3

- (d) Find the general solution to $\sin 2x = \frac{1}{2}$.

2

- (e) Given the equation $x - \cos x = 0$:

2

- (i) explain how you would find your first approximation to the root of this equation.

- (ii) If you take $x_1 = \frac{1}{2}$ as your first approximation, use one application of Newton's Method, to find a better approximation, correct to 3 decimal places.

1

ANSWER SHEET FOR Question 2(b)

(2 Marks)

Question 3 (14 Marks)

- (a) Show that the equation

$$x^3 + x - 3 = 0$$

has a root near 1.2 and, by 'halving the interval' twice, find a better approximation to this root correct to 3 decimal places.

3

1

2

2

1

1

3


1

- (b) Find the remainder when $P(x) = x^3 + 4x^2 - 3$ is divided by $x + 1$.
- (c) If α, β and γ are the roots of the equation $2x^3 - x^2 - 5x + 4 = 0$, find

(i) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$

(ii) $\alpha^2 + \beta^2 + \gamma^2$

(d)




The diagram shows a sketch of the function $y = \tan^{-1} x$.

- (i) Write down the domain and range of the function. 1
 (ii) What are the coordinates of the point B? 1
 (iii) Show that the exact size of the shaded on the diagram is $\frac{1}{2} \log_e 2$. 3
 (iv) Hence, or otherwise, find the area bounded by $y = \tan^{-1} x$, the interval BC and the x -axis. 1

End of Paper.

3




Draw the inverse function of the function shown on the same graph.

(Attach this sheet to your written answers.)

4

Q1

a) $\cos[\sin^{-1} \frac{12}{13}]$



$$\therefore \cos \theta = \frac{5}{13}$$

b) $\frac{d}{dx} \left\{ \cos^{-1} \frac{1}{\sqrt{2x}} \right\} = \frac{-1}{\sqrt{1 - (\frac{1}{\sqrt{2x}})^2}} \times -\frac{1}{2} x^{-\frac{3}{2}}$

$$= \frac{1}{2\sqrt{2x^3} \sqrt{1 - \frac{1}{2x}}}$$

$$= \frac{1}{2\sqrt{2x^3 - x^2}}$$

$$= \frac{1}{2x\sqrt{x-1}}$$

c) $\int \frac{dx}{\sqrt{9-x^2}} = \sin^{-1} \frac{x}{3} + C$

d) $\int_0^4 \frac{dx}{16+x^2} = \left[\frac{1}{4} \tan^{-1} \frac{x}{4} \right]_0^4$
 $= \frac{1}{4} \tan^{-1} 1 - \frac{1}{4} \tan^{-1} 0$

$$= \frac{1}{4} \times \frac{\pi}{4}$$

$$= \frac{\pi}{16}$$

e) range -3 to 3
domain $0 < 2x < \pi$
Inverse function has

$$\therefore y = 3 \cos \frac{x}{2}$$

$$\therefore \text{inverse}$$

$$x = 3 \cos \frac{y}{2}$$

$$\frac{x}{3} = \cos \frac{y}{2}$$

$$y = 2 \cos^{-1} \frac{x}{3}$$

f) $\int_{-a}^a \sec^2 2x dx = \left[\frac{1}{2} \tan 2x \right]_{-a}^a$

$$= \frac{1}{2} \tan 2a - \frac{1}{2} \tan(-2a)$$

$$= \frac{1}{2} \tan 2a + \frac{1}{2} \tan 2a$$

$$= \tan 2a$$

$$\therefore \tan 2a = 1$$


$$2a = \frac{\pi}{4}$$

$$a = \frac{\pi}{8}$$

Q2

a) $\int \frac{e^x}{1+e^{2x}} dx = \tan^{-1} e^x + C$

D



(e) $h = \frac{1-0}{2}$
 $= \frac{1}{2}$

x	f(x)	
0	$\cos^{-1} 0$	$\frac{\pi}{2}$
$\frac{1}{2}$	$\cos^{-1} \frac{1}{2}$	$\times 4$
1	$\cos^{-1} 1$	$\pi \times 4$

$$\text{Total} = \frac{11\pi}{6}$$

$$\therefore \text{Area} = \frac{11/2}{3} \times \frac{11\pi}{6}$$

$$= \frac{11\pi}{36} \text{ units}^2$$

d) $\sin 2x = \frac{1}{2}$

$$2x = \pi n + (-1)^n \sin^{-1} \frac{1}{2}$$

$$= \pi n + (-1)^n \times \frac{\pi}{6}$$

$$x = \frac{\pi n}{2} + \frac{(-1)^n \pi}{12}$$

e) $xc - \cos xc = 0$

i) let $f(x) = xc - \cos xc$, try substituting different values of x until $f(x)$ is +ve for one value and -ve for another. Take x_1 to be the midpoint of these values.

ii) $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$

$$f(\frac{1}{2}) = \frac{1}{2} - \cos \frac{1}{2}$$

$$f'(x) = 1 + \sin x$$

$$f'(\frac{1}{2}) = 1 + \sin \frac{1}{2}$$

$$\therefore x_2 = \frac{1}{2} - \frac{\frac{1}{2} - \cos \frac{1}{2}}{1 + \sin \frac{1}{2}}$$

$$= 0.75522$$

$$= 0.755$$

(3)
 $x^3 + x - 3 = 0$

$$f(1) = 1^3 + 1 - 3 \\ = -1$$

$$f(1.5) = 1.5^3 + 1.5 - 3 \\ = 1.875$$

∴ there is a root between
1 & 1.5 i.e. near 1.2.

$$f\left(\frac{1+1.5}{2}\right) = f(1.25) \\ = 1.25^3 + 1.25 - 3 \\ = 0.203$$

∴ the root lies between 1 & 1.25

$$f\left(\frac{1+1.25}{2}\right) = f(1.125) \\ = 1.125^3 + 1.125 - 3 \\ = -0.45$$

∴ the root lies between 1.125
and 1.25.

$$\text{i.e. } x = \frac{1.125 + 1.25}{2} \\ = 1.1875 \\ = 1.188$$

(b) $P(-1) = (-1)^3 + 4x(-1)^2 - 3$
 $= -1 + 4 - 3$
 $= 0$

c) $2x^3 - x^2 - 5x + 4 = 0$

$$\alpha + \beta + \gamma = \frac{1}{2}$$

$$\alpha\beta + \alpha\gamma + \gamma\beta = -\frac{5}{2}$$

$$\alpha\beta\gamma = -\frac{4}{2} \\ = -2$$

$$\text{i) } \frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma} \\ = \frac{-5/2}{-2} \\ = \frac{5}{4}$$

ii) $\alpha^2 + \beta^2 + \gamma^2 =$

$$\text{Now } (\alpha + \beta + \gamma)^2 = \alpha^2 + \beta^2 + \gamma^2 \\ + 2\alpha\beta + 2\alpha\gamma + 2\beta\gamma$$

$$\therefore \alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma) \\ = \left(\frac{1}{2}\right)^2 - 2 \times -\frac{5}{2}$$

$$= \frac{1}{4} + 5 \\ = 5\frac{1}{4}$$

d) $y = \tan x$

i) domain: all real x
range: $-\frac{\pi}{2} < y < \frac{\pi}{2}$

ii) $\frac{\pi}{4} = \tan^{-1} x$

$$x = \tan \frac{\pi}{4} \\ = 1$$

$$\therefore B(1, \frac{\pi}{4})$$

iii) Area = $\int_0^{\pi/4} f(y) dy$

$$y = \tan^{-1} x$$

$$\therefore x = \tan y$$

$$= \int_0^{\pi/4} \tan y dy$$

$$= \int_0^{\pi/4} \frac{\sin y}{\cos y} dy$$

$$= \left[-\ln(\cos y) \right]_0^{\pi/4}$$

$$= -\ln\left(\cos \frac{\pi}{4}\right) - -\ln(\cos 0)$$

$$= -\ln \frac{1}{\sqrt{2}} + \ln 1$$

$$= -\ln 2^{1/2} + 0$$

$$= \frac{1}{2} \ln 2.$$

iv)

$$\text{Area} = 1 \times \frac{\pi}{4} - \frac{1}{2} \ln 2$$

$$= \frac{\pi}{4} - \frac{1}{2} \ln 2 \text{ units}^2$$