

Question One: (15 Marks) Start a new sheet of paper.

a) Find $\int \frac{x}{\sqrt{2-x^2}} dx$ using the substitution $x = \sqrt{2} \sin \theta$. [2]

b) Show that $\sin(A+B) + \sin(A-B) = 2 \sin A \cos B$, and hence find $\int \sin 5x \cos 3x dx$. [3]

c) Use Integration by Parts to show that $\int_0^1 \tan^{-1} x dx = \frac{\pi}{4} - \frac{1}{2} \ln 2$. [3]

d) Given that $J_n = \int_0^{\frac{\pi}{2}} \cos^n x dx$:

i) Prove that $J_n = \frac{(n-1)}{n} J_{n-2}$, where n is an integer and $n \geq 2$. [4]

ii) Hence evaluate $\int_0^{\frac{\pi}{2}} \cos^6 x dx$. [3]

Question Two: (15 Marks) Start a new sheet of paper.

a) Given that $z = 2+i$ and $\omega = 2-3i$, find, in the form $a+ib$

i) $\left(\frac{\omega}{z}\right)^2$ [1]

ii) $\left(\frac{z}{\omega}\right)$ [1]

b) On the Argand diagram, shade the region where the inequalities


$|z| < 1$ and $\frac{\pi}{4} < \arg(z) < \frac{3\pi}{4}$ both hold. [3]

c) Find the complex square roots of $7+6i\sqrt{2}$, giving your answer in the form $a+ib$, where a and b are real. [3]

(Question 2 continued over)

d) Given the two complex numbers $z_1 = r_1 \text{cis} \theta$ and $z_2 = r_2 \text{cis} \phi$,

i) Show that, if z_1 and z_2 are parallel, $z_1 = kz_2$, for k real. [1]



ABCD is a quadrilateral with vertices A, B, C and D represented by the complex numbers (vectors) z_1, z_2, z_3 and z_4 , as shown in the sketch opposite.

ii) Give two possible vectors (in terms of z_1, z_2) for side AB. [1]

iii) If ABCD is a parallelogram, show that $z_1 - z_2 - z_3 + z_4 = 0$. [3]

e) Explain the fallacy in the following argument: [2]

$$-1 = \sqrt{-1} \times \sqrt{-1} = \sqrt{(-1)(-1)} = \sqrt{1} = 1. \text{ Hence } 1 = -1.$$

Question Three: (15 Marks) Start a new sheet of paper.

a) $F(x)$ is defined by the equation $f(x) = x^2 \left(x - \frac{3}{2}\right)$, on the domain $-2 \leq x \leq 2$.

Note: each sketch below should take about one third of a page.

i) Draw a neat sketch of $F(x)$, labelling all intersections with coordinate axes and turning points. [2]

ii) Sketch $y = \frac{1}{F(x)}$ [2]


iii) Sketch $y = \sqrt{F(x)}$ [2]

iv) Sketch $y = \ln(F(|x|))$ [2]

- b) The Hyperbola \mathcal{H} has the equation $\frac{x^2}{25} - \frac{y^2}{9} = 1$.
- Find the eccentricity of \mathcal{H} . [1]
 - Find the coordinates of the foci of \mathcal{H} . [1]
 - Draw a neat one third of a page size sketch of \mathcal{H} . [2]
 - The line $x = 6$ cuts \mathcal{H} at A and B. Find the coordinates of A and B if A is in the first quadrant. [1]
 - Derive the equation of the tangent to \mathcal{H} at A. [2]

Question Four: (15 Marks) Start a new sheet of paper.

- a) The shaded region bounded by $y = 3 - x^2$, $y = x^2 - x$ and $x = 1$ is rotated about the line $x = 1$. The point P is the intersection of $y = 3 - x^2$ and $y = x^2 - x$ in the second quadrant.



- Find the x coordinate of P. [1]
- Use the method of cylindrical shells to express the volume of the resulting solid of revolution as an integral. [3]
- Evaluate the integral in part (ii) above. [2]

(Question 4 continued over)

- b) Find real numbers A, B and C such that
- $$\frac{x}{(x-1)^2(x-2)} = \frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(x-2)}. \quad [2]$$

Hence show that $\int_0^{\frac{1}{2}} \frac{x}{(x-1)^2(x-2)} dx = 2 \ln\left(\frac{3}{2}\right) - 1$. [2]

- c) Find all x such that $\cos 2x = \sin 3x$, if $0 \leq x \leq \frac{\pi}{2}$. [2]


- d) Solve for x : $\tan^{-1}(3x) - \tan^{-1}(2x) = \tan^{-1}\left(\frac{1}{5}\right)$ [3]

Question Five: (15 Marks) Start a new sheet of paper.

- a) For the polynomial equation $x^3 + 4x^2 + 2x - 3 = 0$ with roots α, β and γ , find:
- The value of $\alpha^2 + \beta^2 + \gamma^2$ [1]
 - The equation whose roots are $(1-\alpha), (1-\beta), (1-\gamma)$. [2]
 - The equation whose roots are $\frac{1}{\alpha}, \frac{1}{\beta}, \frac{1}{\gamma}$. [3]
- b) Determine all the roots of $8x^4 - 25x^3 + 27x^2 - 11x + 1 = 0$ given that it has a root of multiplicity 3. [4]
- c) The equation $x^4 + 4x^3 + 5x^2 + 2x - 20 = 0$ has roots α, β, γ and δ over the complex field.
- Show that the equation whose roots are $\alpha + 1, \beta + 1, \gamma + 1$ and $\delta + 1$ is given by $x^4 - x^2 - 20 = 0$. [2]
 - Hence solve the equation $x^4 + 4x^3 + 5x^2 + 2x - 20 = 0$. [3]

Question Six: (15 Marks) Start a new sheet of paper.

a)



The circles above have centres at O and radii of 5 units and 3 units respectively.

A ray from O making an angle θ with the positive x -axis, cuts the circles at the points M and N as shown.

NS is drawn parallel to the y -axis and MT parallel to the x -axis.

NS and MT intersect at P.

i) Show that the parametric equations of the locus of P in terms of θ are given by $x = 5 \cos \theta$ and $y = 3 \sin \theta$. [2]

ii) By eliminating θ , find the Cartesian equation of this locus. [1]

iii) Find the equation of the normal (in general form) at the point P when $\theta = \frac{\pi}{3}$. [2]

b) The functions $S(x)$ and $C(x)$ are defined by the formulae

$$S(x) = \frac{1}{2}(e^x - e^{-x}) \text{ and } C(x) = \frac{1}{2}(e^x + e^{-x}).$$

i) Verify that $S'(x) = C(x)$. [1]

ii) Show that $S(x)$ is an increasing function for all real x . [1]

iii) Prove $[C(x)]^2 = 1 + [S(x)]^2$ [2]


iv) $S(x)$ has an inverse function, $S^{-1}(x)$, for all real values of x . Briefly justify this statement. [1]

v) Let $y = S^{-1}(x)$. Prove that $\frac{dy}{dx} = \frac{1}{\sqrt{1+x^2}}$. [2]

vi) Hence, or otherwise, show that $S^{-1}(x) = \ln(x + \sqrt{1+x^2})$. [3]

Question Seven: (15 Marks) Start a new sheet of paper.

a) Let OAB be an isosceles triangle, $OA = OB = r$, $AB = b$.




Let OABD be a triangular pyramid with height $OD = h$ and OD perpendicular to the plane of OAB as in the diagram above.

Consider a slice S of the pyramid of width δa as shown at EFGH in the diagram. The slice S is perpendicular to the plane of OAB at FG with $FG \parallel AB$ and $BG = a$. Note that $GH \parallel OD$.

i) Show that the volume of S is $\left(\frac{r-a}{r}\right)b\left(\frac{ah}{r}\right)\delta a$ when δa is small. (You may assume the slice is approximately a rectangular prism of base EFGH and height δa). [3]

ii) Hence show that the pyramid DOAB has a volume of $\frac{1}{6}hbr$. [2]

- c) The diagram below shows a mound of height H . At height h above the horizontal base, the horizontal cross-section of the mound is elliptical in shape, with equation $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \lambda^2$, where $\lambda = 1 - \frac{h^2}{H^2}$, and x, y are appropriate coordinates in the plane of the cross-section.



Show that the volume of the mound is $\frac{8\pi abH}{15}$. [3]

- d) The quadratic equation $x^2 - (2\cos\theta)x + 1 = 0$ has roots α and β .

- i) Find expressions for α and β . [1]
- ii) Show that $\alpha^{10} + \beta^{10} = 2\cos(10\theta)$. [3]

- iii) Suppose now that $\angle AOB = \frac{2\pi}{n}$ and that n identical pyramids DOAB are arranged about O as the centre with common vertical axis OD to form a solid C. Show that the volume V_n of C is given by $V_n = \frac{1}{3}r^2hn\sin\frac{\pi}{n}$. [2]
- iv) Note that when n is large, C approximates a right circular cone. Hence, find $\lim_{n \rightarrow \infty} V_n$ and verify a right circular cone of radius r and height h has a volume of $\frac{1}{3}\pi r^2h$. [2]
- b) On the hyperbola $xy = c^2$, three points P, Q and R are on the same branch, with parameters p, q and r respectively. The tangents at P and Q intersect at U. If O, U and R are collinear, find the relationship between p, q and r . [6]

Question Eight: (15 Marks) Start a new sheet of paper.

- a)
- i) Use the substitution $x = \frac{2}{3}\sin\theta$ to prove that $\int_0^{\frac{2}{3}} \sqrt{4-9x^2} dx = \frac{\pi}{3}$. [3]
- ii) Hence, or otherwise, find the area enclosed by the ellipse $9x^2 + y^2 = 4$. [1]
- b)
- i) Use an appropriate substitution to verify that $\int_0^a \sqrt{a^2 - x^2} dx = \frac{\pi a^2}{4}$. [2]
- ii) Deduce that the area enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is given by πab . [2]


SOLUTIONS: Yr 12 TRIAL HSC EXTN II: 2003

MARKING COMMENTS P1

QUESTION ONE:

a) $x = \sqrt{2} \sin \theta$ so $dx = \sqrt{2} \cos \theta \cdot d\theta$

$$\begin{aligned} & \int \frac{x}{\sqrt{2-x^2}} dx \\ &= \int \frac{\sqrt{2} \sin \theta \cdot \sqrt{2} \cos \theta \cdot d\theta}{\sqrt{2 - (2 \sin^2 \theta)}} \\ &= \int \frac{2 \sin \theta \cos \theta \cdot d\theta}{\sqrt{2(1 - \sin^2 \theta)}} \\ &= \int \frac{2 \sin \theta \cos \theta \cdot d\theta}{2 \cos \theta} \\ &= \int \sin \theta \cdot d\theta \\ &= -\cos \theta + c \\ &= -\sqrt{2} \cdot \frac{\sqrt{2-x^2}}{\sqrt{2}} + c \\ &= -\sqrt{2-x^2} + c \end{aligned}$$



① converting from x to θ . + resolving for $\cos \theta$

• some got $dx = \sqrt{2} \cos \theta \cdot d\theta$, but then left it out of the substitution when changing to θ !!
• very common error was not expressing the indefinite integral back in terms of x ;
• $-\sqrt{2} \cos \theta + c$ got 1 mark.
• $-\sqrt{2} \cos(\sin^{-1}(\frac{x}{\sqrt{2}})) + c$ was also not sufficient for both marks (not simplest form).

① answer

b) $\sin(A+B) = \sin A \cos B + \sin B \cos A$ — ①
 $\sin(A-B) = \sin A \cos B - \sin B \cos A$ — ②
 $\sin(A+B) + \sin(A-B) = 2 \sin A \cos B$ ①+②

① showing relationship

• well done

$$\begin{aligned} & \int \sin 5x \cos 3x \cdot dx \\ &= \int \frac{1}{2} (\sin(5x+3x) + \sin(5x-3x)) \cdot dx \\ &= \int \frac{1}{2} \sin 8x + \frac{1}{2} \sin 2x \cdot dx \\ &= \frac{1}{16} \cos 8x - \frac{1}{4} \cos 2x + c \end{aligned}$$

① correct use of formula

① Answer

c) $\int_0^1 \tan^{-1} x \cdot dx$
 $= \int_0^1 \frac{d(x)}{dx} \cdot \tan^{-1} x \cdot dx$
 $= [x \tan^{-1} x]_0^1 - \int_0^1 x \cdot \frac{1}{1+x^2} dx$
 $= (\frac{\pi}{4} - 0) - \frac{1}{2} \int_0^1 \frac{2x}{1+x^2} dx$
 $= \frac{\pi}{4} - \frac{1}{2} [\ln(1+x^2)]_0^1$
 $= \frac{\pi}{4} - \frac{1}{2} (\ln 2 - \ln 1)$
 $= \frac{\pi}{4} - \frac{1}{2} \ln 2$

① correct integration by parts

• well done

① correct log integration
① correct algebra

SOLUTIONS: Yr 12 TRIAL HSC EXTN II: 2003

MARKING COMMENTS P2

d) i) $J_n = \int_0^{\frac{\pi}{2}} \cos^n x \cdot dx$
 $= \int_0^{\frac{\pi}{2}} \cos x \cos^{n-1} x \cdot dx$
 $= [\sin x \cos^{n-1} x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} (n-1) \cos^{n-2} x \cdot \sin x \cdot \sin x \cdot dx$
 $= (0-0) + (n-1) \int_0^{\frac{\pi}{2}} \cos^{n-2} x \sin^2 x \cdot dx$
 $= (n-1) \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) \cos^{n-2} x \cdot dx$
 $= (n-1) \int_0^{\frac{\pi}{2}} \cos^{n-2} x \cdot dx - (n-1) \int_0^{\frac{\pi}{2}} \cos^n x \cdot dx$
 $J_n = (n-1) J_{n-2} - (n-1) J_n$
 $J_n + (n-1) J_n = (n-1) J_{n-2}$
 $n J_n = (n-1) J_{n-2}$
 $\therefore J_n = \frac{(n-1)}{n} J_{n-2}$

① correct method for splitting cos

• many tried the approach $\int \frac{d(\cos)}{dx} \cdot \cos^n x \cdot dx$ and got lost. These need to be known.

① resolving to integral

① resolving to J_{n-2}

① correct algebra to solution

ii) $\therefore \int_0^{\frac{\pi}{2}} \cos^6 x \cdot dx = J_6$
 $J_6 = \frac{5}{6} \cdot J_4$
 $= \frac{5}{6} \cdot \frac{3}{4} \cdot J_2$
 $= \frac{5}{6} \cdot \frac{3}{4} \cdot \frac{1}{2} \cdot J_0$
 $= \frac{15}{48} \cdot \int_0^{\frac{\pi}{2}} dx$
 $= \frac{15\pi}{96}$

① correctness of formula

• well done.

① evaluates J_0
① answer

QUESTION TWO:


a) (z) $(2-i)^2$
 $= 4 - 4i + i^2$
 $= 3 - 4i$

① answer

ii) $(\frac{z}{w})$
 $= \frac{(2+i)(2+3i)}{(2-3i)(2+3i)}$
 $= \frac{4+6i+2i-3}{4+6i+2i-3}$
 $= \frac{1+3i}{1+8i}$

① answer

• generally good



① boundaries

① correct $|z|$

• generally good.

① correct arg limits.

c) $(a+ib) = \sqrt{7+6i\sqrt{2}}$ a, b real
 $\therefore (a+ib)^2 = 7+6i\sqrt{2}$
 $a^2 - b^2 + 2abi = 7 + 6\sqrt{2}i$
 equating real and imaginary parts.
 $a^2 - b^2 = 7$ — (1) $2ab = 6\sqrt{2}$
 $\therefore a = \frac{6\sqrt{2}}{2b} = \frac{3\sqrt{2}}{b}$ — (2)

substituting (2) in (1):
 $(\frac{3\sqrt{2}}{b})^2 - b^2 = 7$
 $\frac{18}{b^2} - b^2 = 7$
 $18 - b^4 = 7b^2$
 $b^4 + 7b^2 - 18 = 0$
 $\therefore (b^2 + 9)(b^2 - 2) = 0$
 $b^2 = 2, -9$

reject $b^2 = -9$ as b is real.
 $\therefore b = \pm\sqrt{2}$ in (2):
 $b = \sqrt{2} \quad a = \frac{3\sqrt{2}}{\sqrt{2}} = 3$
 $b = -\sqrt{2} \quad a = \frac{3\sqrt{2}}{-\sqrt{2}} = -3$

\therefore roots are $3 + \sqrt{2}i, -3 - \sqrt{2}i$

i) for $z_1 \parallel z_2, \theta = \phi, \therefore z_2 = r_2 \cos \theta$
 $\therefore \cos \theta = \frac{z_2}{r_2}$

\therefore from $z_1 = r_1 \cos \theta = r_1 \cdot \frac{z_2}{r_2}$
 $\therefore z_1 = k z_2$ where $k = \frac{r_1}{r_2}$

ii) Side AB is either \vec{AB} or \vec{BA}
 $\vec{A} + \vec{AB} = \vec{OB} \quad \vec{OB} + \vec{BA} = \vec{OA}$
 $\therefore \vec{AB} = \vec{OB} - \vec{OA} \quad \therefore \vec{BA} = \vec{OA} - \vec{OB}$
 $= z_2 - z_1 \quad = z_1 - z_2$
 $= -(z_2 - z_1)$

① setup a, b relationship
 mostly well done. Some students tried to use formulas for finding square roots of complex numbers (not very successfully)

① resolve for correct b values

① correct roots.

① deducing relationship.

① for both possibilities

generally well done

o.k

iii) side CD is given by $z_3 - z_4$ or $-(z_4 - z_3)$
 as $CD \parallel AB, (z_4 - z_3) = k(z_2 - z_1)$
 but opposite sides of a parallelogram are equal,
 so $|z_4 - z_3| = |z_2 - z_1| \Rightarrow k = \pm 1$
 $k=1: \therefore z_4 - z_3 = z_2 - z_1$
 or $z_1 - z_2 - z_3 + z_4 = 0$
 $k=-1: \therefore z_4 - z_3 = -(z_2 - z_1)$
 but from (ii), AB (or BA) can be either $z_2 - z_1$ or $-(z_2 - z_1)$
 $\therefore z_4 - z_3 = z_2 - z_1$
 $\therefore z_1 - z_2 - z_3 + z_4 = 0$

① side CD

① deriving k

① both cases for k.

not well done.


iv) the argument uses only real numbers, not complex numbers, thus
 $-1 = (a+ib)^2$, where a, b are real.
 $= a^2 - b^2 + 2abi$
 $\therefore a^2 - b^2 = -1$ and $2ab = 0$
 $b = 0 \Rightarrow a^2 = -1$, which cannot happen as a is real
 so $a = 0$ and $-b^2 = -1 \Rightarrow b = \pm 1$
 \therefore the roots are $-i$ and i
 $i^2 = -1$ and $(-i)^2 = (-1)^2 i^2 = -1$
 thus $\sqrt{-1} \times \sqrt{-1} = \sqrt{i^2} \times \sqrt{i^2} = i \times i = i^2 \neq \sqrt{(-1) \times (-1)}$

① identifies use of real nos to try and solve a complex no problem
 ① demonstrates correct procedure (in some way).

Some good answers, most realized something was wrong, but couldn't articulate what it was

QUESTION 3:

a) i) $y = f(x)$
 $f(x) = 2$
 $f(x) = -14$
 $f(x) = \frac{1}{2}$




① axis intercepts

① turning pt. @ $(1, -\frac{1}{2})$

SOLUTIONS: Yr12 TRIAL HSC. EXTN II: 2003

$y = \frac{1}{F(x)}$




MARKING

- ① asymptotes
- ① correct shape in areas.

COMMENTS


p5

i) $y = \sqrt{F(x)}$



- ① correct intercept
- ① correct shape in areas

no penalty for (0,0) not plotted



- ① asymptotes
- ① correct shape in area.

SOLUTIONS: Yr12 TRIAL HSC. EXTN II: 2003

MARKING

COMMENTS

P6


i) $a=5, b=3$ and for hyperbola: $b^2 = a^2(e^2 - 1)$
 $\therefore 9 = 25(e^2 - 1)$
 $e^2 - 1 = \frac{9}{25}$
 $e^2 = \frac{34}{25}$
 $\therefore e = \frac{\sqrt{34}}{5}$

- ① correct value of e

ii) Foci are $S(ae, 0)$ and $S'(-ae, 0)$
 $\therefore (\sqrt{34}, 0)$ and $(-\sqrt{34}, 0)$

- ① correct foci.

iii)



- ① asymptotes
- ① correct shape + intercepts.

could also get this mark if shape is reasonable as scale indicated on y-axis

i) $x=6: \frac{36}{25} - \frac{y^2}{9} = 1$
 $\frac{36}{25} - 1 = \frac{y^2}{9}$
 $y^2 = \frac{9 \times 11}{25}$
 $\therefore y = \pm \frac{\sqrt{99}}{5}$
 $\therefore A$ is $(6, \frac{\sqrt{99}}{5})$ and B is $(6, -\frac{\sqrt{99}}{5})$

- ① correct A and B

ii) $\frac{x^2}{25} - \frac{y^2}{9} = 1$
 $\frac{2x}{25} - \frac{2y}{9} \cdot \frac{dy}{dx} = 0$
 $\therefore \frac{dy}{dx} = \frac{2x}{25} \cdot \frac{9}{2y}$
 $\text{at } (6, \frac{\sqrt{99}}{5}), \frac{dy}{dx} = \frac{9 \cdot 6 \cdot 5}{25 \cdot \sqrt{99}}$
 $= \frac{54}{5\sqrt{99}}$

- ① correct differentiation for $\frac{dy}{dx}$
- ① correct subst to eqn.

ignored small arithmet errors

(cont) $\therefore y - \frac{\sqrt{99}}{5} = \frac{54}{5\sqrt{99}}(x-6)$
 $5\sqrt{99}y - 99 = 54x - 324$
 $0 = 54x - 5\sqrt{99}y - 225$

QUESTION 4:


a) i) at P: $3-x^2 = x^2-x$

$\therefore 0 = 2x^2 - x - 3$
 $= 2x^2 + 2x - 3x - 3$
 $= 2x(x+1) - 3(x+1)$
 $= (x+1)(2x-3)$

$\therefore x = -1, \frac{3}{2}$

$\therefore x$ co-ord of P is -1 (as P is in 2nd quadrant)

ii) typical shell:



inner radius: $r = 1-x$
 outer radius: $R = 1-(x+\delta x)$

\therefore Area of annulus:
 $SA = \pi R^2 - \pi r^2$
 $= \pi(1-(x+\delta x))^2 - \pi(1-x)^2$
 $= \pi[1-2(x+\delta x) + (x+\delta x)^2 - (1-2x+x^2)]$
 $= \pi[1-2x-2\delta x + x^2+2x\delta x + \delta x^2 - 1+2x-x^2]$
 $= \pi(2x\delta x - 2\delta x + \delta x^2)$
 $= 2\pi(x-1)\delta x$ (ignoring δx^2 as too small).

\therefore a small volume of shell is given by

$SV = SA \cdot h$ where $h = (3-x^2) - (x^2-x)$
 $= 3+x-2x^2$

$\therefore \delta V = 2\pi(x-1)(3+x-2x^2)\delta x$
 $= 2\pi(3x+x^2-2x^3-3-x+2x^2)\delta x$
 $= 2\pi(-3+2x+3x^2-2x^3)\delta x$

\therefore Volume of the solid is given by

$V = \sum \delta V$
 $= \lim_{\delta x \rightarrow 0} \sum_{x=1}^{\frac{3}{2}} 2\pi(-3+2x+3x^2-2x^3)\delta x$
 $= 2\pi \int_1^{\frac{3}{2}} (-3+2x+3x^2-2x^3) dx$

① correct value for x

① area of annulus SA

① correct h leading to δV

① correct summing leading to integral

alternatively:
 1 mark: volume of typical shell
 1 mark: correct limits
 1 mark: integration

i) $\therefore V = 2\pi \int_{-1}^{\frac{3}{2}} [-3x+x^2+x^3-\frac{1}{2}x^4] dx$
 $= 2\pi \left[-\frac{3}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 - \frac{1}{10}x^5 \right]_{-1}^{\frac{3}{2}}$
 $= 2\pi \left[\left(-\frac{3}{2} + \frac{1}{3} + \frac{1}{4} - \frac{1}{10}\right) - \left(-\frac{3}{2} + \frac{1}{3} + \frac{1}{4} - \frac{1}{10}\right) \right]$
 $= 2\pi \left[-\frac{1}{2} - 2\frac{1}{2} \right]$
 $= 8\pi$

$\therefore \frac{x}{(x-1)^2(x-2)} = \frac{A(x-1)(x-2) + B(x-2) + C(x-1)^2}{(x-1)^2(x-2)}$

$\therefore x = A(x-1)(x-2) + B(x-2) + C(x-1)^2$ — ①

in ①: $x=1$ $x=2$
 guess: $1 = B(1-2)$ guess: $2 = C(2-1)^2$
 ie $B = -1$ ie $C = 2$

also, from ①: $x = A(x^2-3x+2) + B(x-2) + C(x^2-2x+1)$

equating coefficients of x^2 :
 $0 = A + C$
 $\therefore A = -2$

$\therefore \frac{x}{(x-1)^2(x-2)} = \frac{-2}{(x-1)} - \frac{1}{(x-1)^2} + \frac{2}{x-2}$

$\therefore \int_0^{\frac{1}{2}} \frac{x}{(x-1)^2(x-2)} dx$
 $= \int_0^{\frac{1}{2}} \left[\frac{2}{x-2} - \frac{2}{x-1} - \frac{1}{(x-1)^2} \right] dx$
 $= \int_0^{\frac{1}{2}} \left[\frac{-2}{2-x} + \frac{2}{1-x} - \frac{1}{(x-1)^2} \right] dx$

$= \left[-2\ln(2-x)(-1) + 2\ln(1-x)(-1) + \frac{1}{x-1} \right]_0^{\frac{1}{2}}$
 $= \left[2\ln(2-x) - 2\ln(1-x) + \frac{1}{x-1} \right]_0^{\frac{1}{2}}$
 $= 2\ln\frac{3}{2} + 2\ln\frac{1}{2} - 2 - (2\ln 2 - 2\ln 1 - 1)$
 $= 2\ln\frac{3}{2} + 2\ln 2 - 2 - 2\ln 2 + 0 + 1$
 $= 2\ln\left(\frac{3}{2}\right) - 1$

c) $\cos 2x = c = \sin 3x$ c constant
 $\therefore \sin 3x = c$
 or $\cos\left(\frac{\pi}{2} - 3x\right) = c$
 $\frac{\pi}{2} - 3x = \cos^{-1}(c) + 2\pi n$ $n=0, \pm 1, \pm 2, \dots$

① correct integration
 ① correct answer

• Full marks only for correct solution.

① subst to find B, C (or any other method)

① equating coeffs to find A.

① correct rearrangement to get to integration

① correct subst to show answer.

1) (cont) $\therefore -3x = \frac{\pi}{2} + 2\pi n + \cos^{-1}(c)$
 or $3x = \frac{\pi}{2} - 2\pi n - \cos^{-1}(c)$
 but $\cos 2x = c$ also,
 so $2x = \cos^{-1}(c)$
 $\therefore 3x = \frac{\pi}{2} - 2\pi n - 2x$
 $5x = \left(\frac{1-4n}{2}\right)\pi$
 $\therefore x = \left(\frac{1-4n}{10}\right)\pi$
 for $0 \leq x \leq \frac{\pi}{2}$, we get (using $n=0, n=1$)
 $x = \frac{\pi}{10}, \frac{\pi}{2}$

① correct setup of problem (any method)
 ① correct solutions in range.

d) let $\tan^{-1} 3x = \theta$ and $\tan^{-1} 2x = \phi$
 $\therefore \tan \theta = 3x$ $\tan \phi = 2x$
 for $\tan^{-1}\left(\frac{1}{5}\right) = \tan^{-1}(3x) - \tan^{-1}(2x)$
 $= \theta - \phi$
 taking tan of both sides:
 $\tan(\tan^{-1}\frac{1}{5}) = \tan(\theta - \phi)$
 $\therefore \frac{1}{5} = \frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi}$
 $= \frac{3x - 2x}{1 + 3x \cdot 2x}$
 $1 + 6x^2 = 5x$
 or $0 = 6x^2 - 5x + 1$
 $= 6x^2 - 3x - 2x + 1$
 $= 3x(2x-1) - 1(2x-1)$
 $= (2x-1)(3x-1)$
 $\therefore x = \frac{1}{2}, \frac{1}{3}$

① correct use of tan
 ① forms quadratic
 ① correct answers

QUESTION 5:

a) $\alpha + \beta + \gamma = -4$
 $\alpha\beta + \alpha\gamma + \beta\gamma = 2$
 $\alpha\beta\gamma = 3$
 i) $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \alpha\gamma)$
 $= (-4)^2 - 2(2)$
 $= 12$

① answer

• some had an incorrect squares expansion!

ii) for roots $x = 1 - \alpha \Rightarrow \alpha = (1-x)$
 $\therefore (1-x)$ in eqn gives:
 $(1-x)^3 + 4(1-x)^2 + 2(1-x) - 3 = 0$
 $1 - 3x + 3x^2 - x^3 + 4 - 8x + 4x^2 + 2 - 2x - 3 = 0$
 $\therefore 4 - 13x + 7x^2 - x^3 = 0$
 or $x^3 - 7x^2 + 13x - 4 = 0$

① correct setup with roots
 ① correct eqn.

• many simple algebraic errors

iii) for roots $x = \frac{1}{\alpha} \Rightarrow \alpha = \frac{1}{x}$
 $\therefore \left(\frac{1}{x}\right)^3 + 4\left(\frac{1}{x}\right)^2 + 2\left(\frac{1}{x}\right) - 3 = 0$
 $x^3: 1 + 4x + 2x^2 - 3x^3 = 0$
 or $3x^3 - 2x^2 - 4x - 1 = 0$

① setup with roots
 ① correct eqn.

b) let α be the root of multiplicity 3,
 then $P(\alpha) = P'(\alpha) = P''(\alpha) = 0$.
 $\therefore P'(x) = 32x^3 - 75x^2 + 54x - 11$
 $P''(x) = 96x^2 - 150x + 54$
 if $P''(\alpha) = 0$, α is the soln to $0 = 96x^2 - 150x + 54$
 or $0 = 48x^2 - 75x + 27$
 $\therefore x = \frac{75 \pm \sqrt{2625 - 4 \cdot 48 \cdot 27}}{96}$
 $= \frac{75 \pm \sqrt{441}}{96}$
 $= \frac{75 \pm 21}{96}$
 $= 1, \frac{9}{16}$

① set up problem with $P''(\alpha) = 0$.

• several used $P''(\alpha) = 0$
 • many did not understand the implications for a root with multiplicity!

now, $P'(1) = 32 - 75 + 54 - 11 = 0$
 and $P(1) = 8 - 25 + 27 - 11 + 1 = 0$

① correct possibilities for triple root.

$\therefore (x-1)^3$ is a factor of $P(x)$
 so $\alpha = 1$ is the triple root.

Also, $\alpha^3 \beta = \frac{3}{8}$
 $\therefore \beta = \frac{3}{8}$ is the other root.


① correct triple root with reasons
 ① other root

• need to explicitly state the other root. $(3x-1)$ as a factor implies a root of $x = \frac{1}{3}$.


c) let α, β, γ and δ be the roots of the equation
 i) $\therefore x = \alpha + 1$ is a root of the req'd eqn
 so $\alpha = x - 1$
 $\therefore (x-1)^4 + 4(x-1)^3 + 5(x-1)^2 + 2(x-1) - 20 = 0$

① correct subst for root.

ANS: Yr 12 TRIAL H.S.C. EXTN II: 2003	MARKING	COMMENTS	p11
(cont): $(x^4 - 4x^3 + 6x^2 - 4x + 1) + (4x^3 - 12x^2 + 12x - 4) + (5x^2 - 10x + 5) + 2x - 2 = 0$ $\therefore x^4 - x^2 - 20 = 0$ as reqd.	① correct alg. to soln.	many errors expanding this	
i) now $x^4 - x^2 - 20 = 0$ $(x^2 - 5)(x^2 + 4) = 0$ $\therefore x^2 = 5, -4$ $\therefore x = \pm\sqrt{5}, \pm 2i$ \therefore the roots of $x^4 + 4x^3 + 5x^2 + 2x - 20 = 0$ are given by $x = x - 1$ \therefore roots are $-1 + \sqrt{5}, -1 - \sqrt{5}, -1 + 2i, -1 - 2i$	① correct roots for $x^2 - x^2 - 20 = 0$ ① correct roots for orig. eqn.	many had trouble linking the roots back to the original with $x = x - 1$	
QUESTION 6: a) i) $x_p = 0.5$ $y_p = 0.7$ $= 5 \cos \theta$ $= 3 \sin \theta$	① ① each answer	link back to the definition given in the diagram. This is the starting point, and many missed it.	
ii) $\therefore \frac{x}{5} = \cos \theta$ and $\frac{y}{3} = \sin \theta$ $\frac{x^2}{25} = \cos^2 \theta$ $\frac{y^2}{9} = \sin^2 \theta$ $\therefore \frac{x^2}{25} + \frac{y^2}{9} = \cos^2 \theta + \sin^2 \theta$ $\therefore \frac{x^2}{25} + \frac{y^2}{9} = 1$ is the cartesian eqn.	① correct eqn.	"eliminate θ " \Rightarrow show how this happens, don't just write the equation down!	
iii) normal to an ellipse is given by $\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$ where $a = 5, b = 3$ $\theta = \frac{\pi}{3}$ $\therefore \frac{5x}{\cos \frac{\pi}{3}} - \frac{3y}{\sin \frac{\pi}{3}} = 25 - 9$ $10x - \frac{6y}{\sqrt{3}} = 16$ $\therefore 10\sqrt{3}x - 6y - 16\sqrt{3} = 0$ is eqn.	① correct subst in formula ① correct eqn (any form)	many found tangent instead of normal. put your answer into one of the standard implicit forms - many left their answer unfinished	
b) i) $S(x) = \frac{d}{dx} (\frac{1}{2}(e^x - e^{-x}))$ $= \frac{1}{2}(e^x + e^{-x})$ $= C(x)$	① set out clearly.		
ii) $e^x > 0$ for all x $e^{-x} > 0$ for all x $\therefore e^x + e^{-x} > 0$ for all x $\therefore C(x) > 0$ for all x $\therefore S'(x) > 0$ for all x $\Rightarrow S(x)$ is monotonically increasing	① correct reasoning.	asked to show \Rightarrow give reasons why $S(x) > 0$. Just stating it earns no marks.	

SOLUTIONS: Yr 12 TRIAL H.S.C. EXTN II: 2003	MARKING	COMMENTS	p12
iii) $[C(x)]^2 = [\frac{1}{2}(e^x + e^{-x})]^2$ $= \frac{1}{4}(e^{2x} + 2e^x e^{-x} + e^{-2x})$ $= \frac{1}{4}(e^{2x} + e^{-2x} + 2)$ $1 + [S(x)]^2 = 1 + [\frac{1}{4}(e^{2x} - e^{-2x})]^2$ $= 1 + \frac{1}{4}(e^{2x} - 2e^x e^{-x} + e^{-2x})$ $= \frac{1}{4}(4 + e^{2x} - 2 + e^{-2x})$ $= \frac{1}{4}(e^{2x} + e^{-2x} + 2)$ $= [C(x)]^2$ from above	① expression for $[C(x)]^2$ correct ① reduction of $1 + [S(x)]^2$ correct (or equivalent)		
iv) as $S(x)$ is monotonically increasing, each x must produce a unique y value $\Rightarrow S(x)$ has a 1-1 correspondence $\therefore S^{-1}(x)$ exists for all values of x	① appropriate explanation	many attempted explanations revealed a lack of understanding of what inverse means.	
v) $y = S^{-1}(x)$ $\therefore S(y) = x$ $\therefore \frac{dy}{dx} = S'(y)$ $= C(y)$ $= \sqrt{1 + [S(y)]^2}$ $= \sqrt{1 + x^2}$ $\therefore \frac{dy}{dx} = \sqrt{1 + x^2}$	① inverse rule to give $\frac{dy}{dx}$ correctly ① correct subst to formula.	very few picked up the links to $S(x)$ and $C(x)$ in the previous parts, so many futile atten at a simple problem. look at linked parts like this one - th make the solution simpler!	
vi) $\therefore y = \int \frac{dx}{\sqrt{1+x^2}}$ let $x = \tan \theta$ $\therefore dx = \sec^2 \theta d\theta$ $= \int \frac{\sec^2 \theta d\theta}{\sqrt{1 + \tan^2 \theta}}$ $= \int \frac{\sec^2 \theta d\theta}{\sec \theta}$ $= \int \sec \theta d\theta$ $= \int \frac{\sec \theta (\sec \theta + \tan \theta)}{(\sec \theta + \tan \theta)} d\theta$ $= \ln(\sec \theta + \tan \theta) + C$ $y = \ln(x + \sqrt{1+x^2}) + C$	① reduction to $\int \sec \theta d\theta$ ① correct \int of $\sec \theta$ ① correct subst to give y in terms of x .		"The question is to show the relationship \Rightarrow not using the standard integral table. The integration is the question, its not part of something bigger.
QUESTION 7: a) i) In base ΔOAB :	① correct setup of variables		
	$GB = a$ $OB = r$ $NB = \frac{b}{2}$ $OQ = r - a$		

i) (cont) $\frac{MA}{OB} = \frac{NB}{OB}$
 or $MA = \frac{NB \cdot OB}{OB}$
 $= \frac{b \cdot (r-a)}{r}$
 $\therefore FG = \frac{2MA}{r}$
 $= \frac{2b(r-a)}{r}$



$\therefore V_s = AB \cdot FG \cdot \frac{1}{2} \cdot \frac{1}{n}$
 $= \left(\frac{r-a}{r}\right) b \cdot \left(\frac{ah}{r}\right) \frac{1}{2} \cdot \frac{1}{n}$

ii) $\therefore V = \int_0^r \left(\frac{r-a}{r}\right) b \cdot \left(\frac{ah}{r}\right) da$

$= \frac{bh}{r^2} \int_0^r a(r-a) da$
 $= \frac{bh}{r^2} \int_0^r ar - a^2 da$
 $= \frac{bh}{r^2} \left[\frac{1}{2} ar^2 - \frac{1}{3} a^3 \right]_0^r$
 $= \frac{bh}{r^2} \left[\frac{1}{2} r^3 - \frac{1}{3} r^3 - 0 \right]$
 $= \frac{bh}{r^2} \cdot \left(\frac{1}{2} - \frac{1}{3}\right) r^3$
 $= \frac{1}{6} bhr$ as reqd

ii) given $\angle AOB = \frac{2\pi}{n}$



ie $\theta = \frac{2\pi}{n}$
 $\therefore \frac{\theta}{2} = \frac{\pi}{n}$
 $\therefore \sin \frac{\theta}{2} = \frac{b}{2} \cdot \frac{1}{r}$
 or $b = 2r \sin \frac{\theta}{2}$
 $= 2r \sin \frac{\pi}{n}$

$\therefore V = \frac{1}{6} bhr$ from (ii) above
 so $V = \frac{1}{6} hr \cdot 2r \sin \frac{\pi}{n}$
 $= \frac{1}{3} hr^2 \sin \frac{\pi}{n}$
 ie $V_n = \frac{1}{3} hr^2 n \sin \frac{\pi}{n}$

① correct value for FG

① correct expression for FG


① reduction to correct \int

① correct subst to expression

① correct derivation for b

① correct expression for V

iii) (cont) $\lim_{n \rightarrow \infty} V_n = \lim_{n \rightarrow \infty} \frac{1}{3} nhr^2 \sin \frac{\pi}{n}$
 $= \lim_{n \rightarrow \infty} \frac{1}{3} hr^2 \pi \cdot \frac{2\pi n}{\pi}$
 let $x = \frac{\pi}{n}$; as $n \rightarrow \infty, \frac{\pi}{n} \rightarrow 0$
 $\therefore \lim_{n \rightarrow \infty} V_n = \lim_{x \rightarrow 0} \frac{1}{3} hr^2 \pi \cdot \lim_{x \rightarrow 0} \frac{\sin x}{x}$
 $= \frac{1}{3} \pi r^2 h$



tangent at P: $x + p^2y = 2cp$ — ①
 " " Q: $x + q^2y = 2cq$ — ②
 $(p^2 - q^2)y = 2c(p - q)$: ① - ②

$\therefore y_u = \frac{2c}{p+q}$

$\therefore x_u = 2cp - \frac{p^2 \cdot 2c}{p+q}$

Now $m_{OU} = \frac{2c}{p+q} \cdot \frac{p+q}{2cp}$
 $= \frac{1}{pq}$

\therefore eqn of OUR is $y = \frac{x}{pq}$ — ③

$xy = c^2$ — ④

subst ③ in ④: $\frac{x^2}{pq} = c^2$

$\therefore x^2 = pq c^2$

$\therefore x_R = c\sqrt{pq}$

but R is $(cr, \frac{c}{r})$

so $cr = c\sqrt{pq}$
 $r = \sqrt{pq}$
 or $r^2 = pq$

① limit expression
 ① correct use of limits to evaluate expression

① diagram showing relationships

① finding y coord of U

① finding x coord of U

① gradient of OU

① finding x_R (or y_R)

① correct relationships (either form)

many incorrect uses of the limit.

many students didn't draw correct diagrams!

not many students completed the correct relationship.

Some became lost after getting correct points of intersection, others made it much more complicated than it was.

QUESTION 8:

a) i) $\int_0^{\frac{2}{3}} \sqrt{4-9x^2} dx$

$\therefore = \int_0^{\frac{\pi}{2}} \sqrt{4-4\sin^2\theta} \cdot \frac{2}{3} \cos\theta d\theta$

$= \frac{4}{3} \int_0^{\frac{\pi}{2}} \sqrt{\cos^2\theta} \cdot \cos\theta d\theta$

$= \frac{4}{3} \int_0^{\frac{\pi}{2}} \cos^2\theta d\theta$

$= \frac{4}{3} \int_0^{\frac{\pi}{2}} \frac{1}{2}(\cos 2\theta + 1) d\theta$

$= \frac{2}{3} [\frac{1}{2} \sin 2\theta + \theta]_0^{\frac{\pi}{2}}$

$= \frac{2}{3} [(2 \cdot 0 + \frac{\pi}{2}) - 0]$

$x = \frac{2}{3} \sin\theta$
 $\therefore dx = \frac{2}{3} \cos\theta d\theta$
 when $x = \frac{2}{3}$, $\theta = \frac{\pi}{2}$
 $x = 0$, $\theta = 0$

① correct subst to θ , including limits

① correct reduction to $\int \cos^2$

① correct subst to soln.

ii) for $9x^2 + y^2 = 4$

$y^2 = 4 - 9x^2$

$\therefore y = \sqrt{4 - 9x^2}$

\therefore pt i) gives the area in the first quadrant so, from symmetry, this is $\frac{1}{4}$ the reqd. area.

$\therefore A = 4 \cdot \frac{\pi}{3}$

$= \frac{4\pi}{3}$ sq units

① answer.

b) i) $x = a \sin\theta$

$\therefore dx = a \cos\theta d\theta$

when $x = a$, $\theta = \frac{\pi}{2}$

$x = 0$, $\theta = 0$

$\therefore \int_0^a \sqrt{a^2 - x^2} dx$

$= \int_0^{\frac{\pi}{2}} \sqrt{a^2 - a^2 \sin^2\theta} \cdot a \cos\theta d\theta$

$= a^2 \int_0^{\frac{\pi}{2}} \sqrt{1 - \sin^2\theta} \cdot \cos\theta d\theta$

$= a^2 \int_0^{\frac{\pi}{2}} \cos^2\theta d\theta$

$= a^2 \int_0^{\frac{\pi}{2}} \frac{1}{2}(\cos 2\theta + 1) d\theta$

$= a^2 [\frac{1}{4} \sin 2\theta - \frac{\theta}{2}]_0^{\frac{\pi}{2}}$

$= a^2 [(0 + \frac{\pi}{4}) - 0]$

① correct subst inc limits

① correct integration to solution

ii) from $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$b^2x^2 + a^2y^2 = a^2b^2$

$a^2y^2 = a^2b^2 - b^2x^2$

$\therefore y^2 = b^2 - \frac{b^2}{a^2}x^2$

$\therefore y = \sqrt{b^2 - \frac{b^2}{a^2}x^2}$

$= \sqrt{\frac{b^2a^2 - b^2x^2}{a^2}}$

$= \frac{b}{a} \sqrt{a^2 - x^2}$

\therefore area of 1st quadrant is

$A_1 = \int_0^a \frac{b}{a} \sqrt{a^2 - x^2} dx$

$= \frac{b}{a} \cdot \frac{\pi a^2}{4}$ from (i)

$= \frac{\pi ab}{4}$

\therefore total area (from symmetry)

$A = 4 \cdot \frac{\pi ab}{4}$

$= \pi ab$

① reducing equation to stand form

① correct reasoning to soln.

c) $\Delta V = A \Delta h$ where A is the area of the ellipse at height h .

\therefore from (b) above:

$A = \pi ab \lambda^2$

(as $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \lambda^2$ becomes

$\frac{x^2}{a^2\lambda^2} + \frac{y^2}{b^2\lambda^2} = 1$)

$\therefore \Delta V = \pi ab \lambda^2 \Delta h$

$\therefore V = \int_0^H \pi ab (1 - \frac{h^2}{H^2})^2 dh$

$= \pi ab \int_0^H 1 - \frac{2h^2}{H^2} + \frac{h^4}{H^4} dh$

$= \pi ab [h - \frac{2}{3} \frac{h^3}{H^2} + \frac{h^5}{5H^4}]_0^H$

$= \pi ab [(H - \frac{2}{3} \frac{H^3}{H^2} + \frac{H^5}{5H^4}) - 0]$

$= \pi ab [\frac{15H - 10H + 3H}{15}]$

$= \frac{8\pi ab H}{15}$ as reqd.

① correct deduction of A

① correct expression for V in terms of h 's.

① correct \int leading to soln

wrong λ , but correct method, gained mark

SOLUTIONS: Yr 12 TRIAL HSC. EXTN II: 2003

MARKING

COMMENTS

P17

d) $x^2 - (2\cos\theta)x + 1 = 0$

i) $x^2 - (2\cos\theta)x + \cos^2\theta = -1 + \cos^2\theta$

$\therefore (x - \cos\theta)^2 = -\sin^2\theta$

$\therefore x - \cos\theta = \pm i \sin\theta$

$\therefore x = \cos\theta \pm i \sin\theta$

$\therefore \alpha = \cos\theta + i \sin\theta \quad \beta = \cos\theta - i \sin\theta$

ii) $\alpha = \cos\theta$

$\therefore \alpha^{10} = (\cos\theta)^{10}$

$= \cos 10\theta$ by de Moivre's Theorem

similarly $\beta = \overline{\cos\theta}$

so $\beta^{10} = (\overline{\cos\theta})^{10}$

$= \overline{\cos 10\theta}$

$\therefore \alpha^{10} + \beta^{10} = \cos 10\theta + \overline{\cos 10\theta}$

$= \cos 10\theta + i \sin 10\theta + \cos 10\theta - i \sin 10\theta$

$= 2 \cos 10\theta$ as reqd.

both
① answers

① correct use
of de Moivre's
Theorem

① correct
use of $\overline{\cos\theta}$

① correct
algebra to
soln.