

BOARD OF STUDIES
NEW SOUTH WALES

2009

HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

Total marks – 120

- Attempt Questions 1–10
- All questions are of equal value

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Total marks – 120

Attempt Questions 1–10

All questions are of equal value

Answer each question in the appropriate writing booklet. Extra writing booklets are available.

Question 1 (12 marks) Use the Question 1 Writing Booklet.


- (a) Sketch the graph of $y - 2x = 3$, showing the intercepts on both axes. 2
- (b) Solve $\frac{5x-4}{x} = 2$. 2
- (c) Solve $|x+1| = 5$. 2
- (d) Find the gradient of the tangent to the curve $y = x^4 - 3x$ at the point $(1, -2)$. 2
- (e) Find the exact value of θ such that $2 \cos \theta = 1$, where $0 \leq \theta \leq \frac{\pi}{2}$. 2
- (f) Solve the equation $\ln x = 2$. Give your answer correct to four decimal places. 2

Question 2 (12 marks) Use the Question 2 Writing Booklet.

- (a) Differentiate with respect to x :
- (i) $x \sin x$ 2
- (ii) $(e^x + 1)^2$. 2
- (b) (i) Find $\int 5 dx$. 1
- (ii) Find $\int \frac{3}{(x-6)^2} dx$. 2
- (iii) Evaluate $\int_1^4 x^2 + \sqrt{x} dx$. 3
- (c) Evaluate $\sum_{k=1}^4 (-1)^k k^2$. 2

Question 3 (12 marks) Use the Question 3 Writing Booklet.

- (a) An arithmetic series has 21 terms. The first term is 3 and the last term is 53. 2
Find the sum of the series.




The circle in the diagram has centre N . The line LM is tangent to the circle at P .

- (i) Find the equation of LM in the form $ax + by + c = 0$. 2
 (ii) Find the distance NP . 2
 (iii) Find the equation of the circle. 1
- (c) Shade the region in the plane defined by $y \geq 0$ and $y \leq 4 - x^2$. 2

Question 3 continues on page 5

Question 3 (continued)

- (d) The diagram shows a block of land and its dimensions, in metres. The block of land is bounded on one side by a river. Measurements are taken perpendicular to the line AB , from AB to the river, at equal intervals of 50 m. 3



Use Simpson's rule with six subintervals to find an approximation to the area of the block of land.

End of Question 3

Question 4 (12 marks) Use the Question 4 Writing Booklet.

- (a) A tree grows from ground level to a height of 1.2 metres in one year. In each subsequent year, it grows $\frac{9}{10}$ as much as it did in the previous year. 2


Find the limiting height of the tree.

- (b) Find the values of k for which the quadratic equation 3

$$x^2 - (k+4)x + (k+7) = 0$$

has equal roots.

- (c) In the diagram, $\triangle ABC$ is a right-angled triangle, with the right angle at C . The midpoint of AB is M , and $MP \perp AC$.



Copy or trace the diagram into your writing booklet.


- (i) Prove that $\triangle AMP$ is similar to $\triangle ABC$. 2
- (ii) What is the ratio of AP to AC ? 1
- (iii) Prove that $\triangle AMC$ is isosceles. 2
- (iv) Show that $\triangle ABC$ can be divided into two isosceles triangles. 1
- (v) 1



Copy or trace this triangle into your writing booklet and show how to divide it into four isosceles triangles.

Question 5 (12 marks) Use the Question 5 Writing Booklet.

- (a) In the diagram, the points A and C lie on the y -axis and the point B lies on the x -axis. The line AB has equation $y = \sqrt{3}x - 3$. The line BC is perpendicular to AB .




- (i) Find the equation of the line BC . 2
- (ii) Find the area of the triangle ABC . 2
- (b) On each working day James parks his car in a parking station which has three levels. He parks his car on a randomly chosen level. He always forgets where he has parked, so when he leaves work he chooses a level at random and searches for his car. If his car is not on that level, he chooses a different level and continues in this way until he finds his car.
- (i) What is the probability that his car is on the first level he searches? 1
- (ii) What is the probability that he must search all three levels before he finds his car? 1
- (iii) What is the probability that on every one of the five working days in a week, his car is not on the first level he searches? 1

Question 5 continues on page 8

Question 5 (continued)

- (c) The diagram shows a circle with centre O and radius 2 centimetres. The points A and B lie on the circumference of the circle and $\angle AOB = \theta$.




NOT TO SCALE

- (i) There are two possible values of θ for which the area of $\triangle AOB$ is $\sqrt{3}$ square centimetres. One value is $\frac{\pi}{3}$.
Find the other value. 2
- (ii) Suppose that $\theta = \frac{\pi}{3}$.
- (1) Find the area of the sector AOB . 1
- (2) Find the exact length of the perimeter of the minor segment bounded by the chord AB and the arc AB . 2

End of Question 5

Question 6 (12 marks) Use the Question 6 Writing Booklet.

- (a) The diagram shows the region bounded by the curve $y = \sec x$, the lines $x = \frac{\pi}{3}$ and $x = -\frac{\pi}{3}$, and the x -axis. 3




The region is rotated about the x -axis. Find the volume of the solid of revolution formed.

- (b) Radium decays at a rate proportional to the amount of radium present. That is, if $Q(t)$ is the amount of radium present at time t , then $Q = Ae^{-kt}$, where k is a positive constant and A is the amount present at $t = 0$. It takes 1600 years for an amount of radium to reduce by half.
- (i) Find the value of k . 2
- (ii) A factory site is contaminated with radium. The amount of radium on the site is currently three times the safe level. 2
- How many years will it be before the amount of radium reaches the safe level?

Question 6 continues on page 10

Question 6 (continued)

- (c) The diagram illustrates the design for part of a roller-coaster track. The section RO is a straight line with slope 1.2, and the section PQ is a straight line with slope -1.8 . The section OP is a parabola $y = ax^2 + bx$. The horizontal distance from the y -axis to P is 30 m.



In order that the ride is smooth, the straight line sections must be tangent to the parabola at O and at P .

- (i) Find the values of a and b so that the ride is smooth. 3
- (ii) Find the distance d , from the vertex of the parabola to the horizontal line through P , as shown on the diagram. 2

End of Question 6

Question 7 (12 marks) Use the Question 7 Writing Booklet.

- (a) The acceleration of a particle is given by

$$\ddot{x} = 8e^{-2t} + 3e^{-t},$$

where x is displacement in metres and t is time in seconds.

Initially its velocity is -6 m s^{-1} and its displacement is 5 m.

- (i) Show that the displacement of the particle is given by 2

$$x = 2e^{-2t} + 3e^{-t} + t.$$

- (ii) Find the time when the particle comes to rest. 3
- (iii) Find the displacement when the particle comes to rest. 1

- (b) Between 5 am and 5 pm on 3 March 2009, the height, h , of the tide in a harbour was given by

$$h = 1 + 0.7 \sin \frac{\pi}{6} t \text{ for } 0 \leq t \leq 12,$$


where h is in metres and t is in hours, with $t = 0$ at 5 am.

- (i) What is the period of the function h ? 1
- (ii) What was the value of h at low tide, and at what time did low tide occur? 2
- (iii) A ship is able to enter the harbour only if the height of the tide is at least 1.35 m. 3

Find all times between 5 am and 5 pm on 3 March 2009 during which the ship was able to enter the harbour.

Question 8 (12 marks) Use the Question 8 Writing Booklet.

(a)



The diagram shows the graph of a function $y = f(x)$.

- (i) For which values of x is the derivative, $f'(x)$, negative? 1
- (ii) What happens to $f'(x)$ for large values of x ? 1
- (iii) Sketch the graph $y = f'(x)$. 2

Question 8 continues on page 13

Question 8 (continued)

- (b) One year ago Daniel borrowed \$350 000 to buy a house. The interest rate was 9% per annum, compounded monthly. He agreed to repay the loan in 25 years with equal monthly repayments of \$2937.
- (i) Calculate how much Daniel owed after his first monthly repayment. 1
- (ii) Daniel has just made his 12th monthly repayment. He now owes \$346 095. The interest rate now decreases to 6% per annum, compounded monthly. 3
- The amount, $\$A_n$, owing on the loan after the n th monthly repayment is now calculated using the formula
- $$A_n = 346\,095 \times 1.005^n - 1.005^{n-1}M - \dots - 1.005M - M$$
- where $\$M$ is the monthly repayment, and $n = 1, 2, \dots, 288$. (Do NOT prove this formula.)
- Calculate the monthly repayment if the loan is to be repaid over the remaining 24 years (288 months).
- (iii) Daniel chooses to keep his monthly repayments at \$2937. Use the formula in part (ii) to calculate how long it will take him to repay the \$346 095. 3
- (iv) How much will Daniel save over the term of the loan by keeping his monthly repayments at \$2937, rather than reducing his repayments to the amount calculated in part (ii)? 1

End of Question 8

Question 9 (12 marks) Use the Question 9 Writing Booklet.


- (a) Each week Van and Marie take part in a raffle at their respective workplaces.
The probability that Van wins a prize in his raffle is $\frac{1}{9}$. The probability that Marie wins a prize in her raffle is $\frac{1}{16}$.

What is the probability that, during the next three weeks, at least one of them wins a prize?

- (b) An oil rig, S , is 3 km offshore. A power station, P , is on the shore. A cable is to be laid from P to S . It costs \$1000 per kilometre to lay the cable along the shore and \$2600 per kilometre to lay the cable underwater from the shore to S .

The point R is the point on the shore closest to S , and the distance PR is 5 km.

The point Q is on the shore, at a distance of x km from R , as shown in the diagram.



- (i) Find the total cost of laying the cable in a straight line from P to R and then in a straight line from R to S . 1
- (ii) Find the cost of laying the cable in a straight line from P to S . 1
- (iii) Let $\$C$ be the total cost of laying the cable in a straight line from P to Q , and then in a straight line from Q to S . 2
- Show that $C = 1000(5 - x + 2.6\sqrt{x^2 + 9})$.
- (iv) Find the minimum cost of laying the cable. 4
- (v) New technology means that the cost of laying the cable underwater can be reduced to \$1100 per kilometre. 2

Determine the path for laying the cable in order to minimise the cost in this case.

Question 10 (12 marks) Use the Question 10 Writing Booklet.

Let $f(x) = x - \frac{x^2}{2} + \frac{x^3}{3}$.

- (a) Show that the graph of $y = f(x)$ has no turning points. 2
- (b) Find the point of inflexion of $y = f(x)$. 1
- (c) (i) Show that $1 - x + x^2 - \frac{1}{1+x} = \frac{x^3}{1+x}$ for $x \neq -1$. 1
- (ii) Let $g(x) = \ln(1+x)$. 2

Use the result in part (c) (i) to show that $f'(x) \geq g'(x)$ for all $x \geq 0$.

- (d) On the same set of axes, sketch the graphs of $y = f(x)$ and $y = g(x)$ for $x \geq 0$. 2
- (e) Show that $\frac{d}{dx}[(1+x)\ln(1+x) - (1+x)] = \ln(1+x)$. 2
- (f) Find the area enclosed by the graphs of $y = f(x)$ and $y = g(x)$, and the straight line $x = 1$. 2


End of paper

2009 Higher School Certificate Solutions Mathematics

Question 1

(a) $y - 2x = 3$

x-intercept: y-intercept:
Let $y = 0$ Let $x = 0$
 $-2x = 3$ $y = 3$
 $x = -1.5$



(b) $\frac{5x-4}{x} = 2$

$5x - 4 = 2x$
 $3x = 4$

$x = \frac{4}{3}$ or $1\frac{1}{3}$.

(c) $|x+1| = 5$

$x+1=5$ $-(x+1)=5$
 $x=4$ $-x-1=5$
 $-x=6$
 $x=-6$

$\therefore x = 4$ or -6 .

(d) $y = x^4 - 3x$

$\frac{dy}{dx} = 4x^3 - 3$

When $x = 1$,

$m_{\text{tang}} = 4(1)^3 - 3$
 $= 1$.

(e) $2 \cos \theta = 1$

$\cos \theta = \frac{1}{2}$

$\theta = \frac{\pi}{3}$.

(f) $\ln x = 2$

i.e. $\log_e x = 2$

$x = e^2$
 $= 7.389\ 05\dots$
 $= 7.3891$ (to 4 decimal places).

Question 2

(a) (i) Let $y = x \sin x$

Using the product rule,

Let $u = x$, $u' = 1$

$v = \sin x$, $v' = \cos x$

$\frac{dy}{dx} = u \cdot \frac{dv}{dx} + v \cdot \frac{du}{dx}$

$\frac{dy}{dx} = x \cos x + \sin x$.

(ii) Let $y = (e^x + 1)^2$

$\frac{dy}{dx} = 2(e^x + 1)(e^x)$
 $= 2e^x(e^x + 1)$.

(b) (i) $\int 5 dx = 5x + c$.

(ii) $\int \frac{3}{(x-6)^2} dx$
 $= \int 3(x-6)^{-2} dx$
 $= -3(x-6)^{-1} + c$
 $= -\frac{3}{(x-6)} + c$.

(iii) $\int_1^4 x^2 + \sqrt{x} dx$
 $= \int_1^4 x^2 + x^{\frac{1}{2}} dx$
 $= \left[\frac{x^3}{3} + \frac{2x^{\frac{3}{2}}}{3} \right]_1^4$
 $= \left(\frac{4^3}{3} + \frac{2(4)^{\frac{3}{2}}}{3} \right) - \left(\frac{1^3}{3} + \frac{2(1)^{\frac{3}{2}}}{3} \right)$
 $= \frac{64}{3} + \frac{16}{3} - \frac{1}{3} - \frac{2}{3}$
 $= \frac{77}{3}$.

(c) $\sum_{k=1}^4 (-1)^k k^2$
 $= (-1) \cdot 1^2 + (-1)^2 \cdot 2^2 + (-1)^3 \cdot 3^2$
 $\quad + (-1)^4 \cdot 4^2$
 $= -1 + 4 - 9 + 16$
 $= 10$.

Question 3

(a) $a = 3$, $l = 53$, $n = 21$

$S_n = \frac{n}{2}(a+l)$

$S_{21} = \frac{21}{2}(3+53)$
 $= 588$.

(b) (i) $m = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{5-1}{5-2}$
 $= \frac{4}{3}$

Using $(2, 1)$ and $m = \frac{4}{3}$,

$y - y_1 = m(x - x_1)$

$y - 1 = \frac{4}{3}(x - 2)$

$3y - 3 = 4(x - 2)$

$3y - 3 = 4x - 8$

$0 = 4x - 3y - 5$

$\therefore 4x - 3y - 5 = 0$.

(ii) $d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$

Using $(1, 3)$ and $4x - 3y - 5 = 0$,

$NP = d = \frac{|4(1) - 3(3) - 5|}{\sqrt{4^2 + (-3)^2}}$

$= \frac{|-10|}{5}$


$= 2$ units.

(iii) Centre $(1, 3)$, $r = 2$

$(x-h)^2 + (y-k)^2 = r^2$

$(x-1)^2 + (y-3)^2 = 4$.

- (c) $y \geq 0$ and $y \leq 4 - x^2$
 Consider $y = 4 - x^2$.
 x-intercept: y-intercept:
 Let $y = 0$ Let $x = 0$
 $0 = 4 - x^2$ $y = 4$
 $x = \pm 2$.



(d) **METHOD 1** Function method:

x	0	50	100	150	200	250	300
y	210	220	200	190	210	240	240

$y_0 \quad y_1 \quad y_2 \quad y_3 \quad y_4 \quad y_5 \quad y_6$

$$\begin{aligned} \text{Area} &\doteq \frac{h}{3} [y_0 + 4(y_1 + y_3 + y_5) \\ &\quad + 2(y_2 + y_4) + y_6] \\ &= \frac{50}{3} [210 + 4(220 + 190 + 240) \\ &\quad + 2(200 + 210) + 240] \\ &= 64\,500 \text{ m}^2. \end{aligned}$$

METHOD 2 Weights method:

x	0	50	100	150	200	250	300
y	210	220	200	190	210	240	240
Weights	1	4	2	4	2	4	1
y × wts	210	880	400	760	420	960	240

$$\begin{aligned} \text{Area} &\doteq \frac{h}{3} \sum (y \times \text{wts}) \\ &= \frac{50}{3} \times (210 + 880 + 400 + 760 \\ &\quad + 420 + 960 + 240) \\ &= \frac{50}{3} \times 3870 \\ &= 64\,500 \text{ m}^2. \end{aligned}$$

Question 4

- (a) This is geometric series where

$$a = 1.2, \quad r = \frac{9}{10} = 0.9.$$

$$\begin{aligned} S_\infty &= \frac{a}{1-r}, \text{ since } |r| < 1 \\ &= \frac{1.2}{1-0.9} \\ &= 12 \end{aligned}$$

\therefore The limiting height is 12 m.

- (b) $x^2 - (k+4)x + (k+7) = 0$
 $a = 1, \quad b = -(k+4), \quad c = k+7$
 $\Delta = b^2 - 4ac$
 $= [-(k+4)]^2 - 4(1)(k+7)$
 $= k^2 + 8k + 16 - 4k - 28$
 $= k^2 + 4k - 12$


For equal roots, $\Delta = 0$

$$k^2 + 4k - 12 = 0$$

$$(k+6)(k-2) = 0$$


$\therefore k = -6$ or $k = 2$.

(c)



- (i) In triangles AMP and ABC ,
 $\angle APM = \angle ACB = 90^\circ$ ($MP \perp AC$ and $BC \perp AC$)
 $\angle MAP = \angle BAC$ (common angle)
 $\therefore \triangle AMP \sim \triangle ABC$ (equiangular).
- (ii) $AM = MB$ (M is midpoint of AB)
 $AM : MB = 1 : 2$
 $\therefore AP : AC = 1 : 2$ (corresponding sides in similar triangles $\triangle AMP$ and $\triangle ABC$ are in the same ratio).

- (iii) Construct CM .
 In triangles AMP and CMP ,
 $AP = PC$ (from (ii))
 In $\triangle AMP$ and $\triangle CMP$,
 $\angle APM = \angle CPM = 90^\circ$ ($MP \perp AC$)
 PM is common.
 $\therefore \triangle AMP \cong \triangle CMP$ (SAS)
 $\therefore AM = CM$ (corresponding sides of congruent Δ s)
 $\therefore \triangle AMC$ is isosceles.
- (iv) $AM = CM$ as proven in (iii) above.
 But $AM = MB$ (M is midpoint of AB)
 $\therefore AM = CM = MB$
 $\therefore \triangle MCB$ is isosceles.



(v)

Question 5

- (a) (i) $AB: y = \sqrt{3}x - 3$

$$\therefore m = \sqrt{3}$$

x-intercept:

$$\text{Let } y = 0$$

$$\sqrt{3}x = 3$$

$$x = \frac{3}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$$

$$= \sqrt{3}$$

$\therefore B$ is $(\sqrt{3}, 0)$.


Equation of BC :

$$\text{Using } B(\sqrt{3}, 0) \text{ and } m_{\text{perp}} = -\frac{1}{\sqrt{3}},$$

$$y - y_1 = m(x - x_1)$$

$$y - 0 = -\frac{1}{\sqrt{3}}(x - \sqrt{3})$$

$$y = -\frac{1}{\sqrt{3}}x + 1.$$



Coordinates of C : $(0, 1)$

Coordinates of A : $(0, -3)$

\therefore Length $AC = 4$

Also, length $OB = \sqrt{3}$


\therefore Area of $\triangle ABC = \frac{1}{2}bh$

$$= \frac{1}{2} \times 4 \times \sqrt{3}$$

$$= 2\sqrt{3} \text{ units}^2.$$

(b) (i) $P(\text{on 1st level searched}) = \frac{1}{3}$.

(ii) 1st level searched 2nd level searched 3rd level searched



$P(\text{search all 3 levels}) = \frac{2}{3} \times \frac{1}{2} \times 1 = \frac{1}{3}$.

(iii) $P\left(\text{not on 1st level searched for 5 days}\right) = \left(1 - \frac{1}{3}\right)^5 = \left(\frac{2}{3}\right)^5 = \frac{32}{243}$.

(c) (i) Area of $\triangle AOB = \frac{1}{2} ab \sin O$

$\sqrt{3} = \frac{1}{2} \times 2 \times 2 \times \sin \theta$

$\sqrt{3} = 2 \sin \theta$

$\therefore \sin \theta = \frac{\sqrt{3}}{2}$

$\theta = \frac{\pi}{3}, \pi - \frac{\pi}{3}$

$= \frac{\pi}{3}, \frac{2\pi}{3}$

\therefore Other value of $\theta = \frac{2\pi}{3}$.

(ii) (1) Area of sector $AOB = \frac{1}{2} r^2 \theta$

$= \frac{1}{2} \times 2^2 \times \frac{\pi}{3}$

$= \frac{2\pi}{3} \text{ cm}^2$.

(2) $\theta = \frac{\pi}{3} = 60^\circ$

$\therefore \angle A = \angle B = 60^\circ$

$\therefore AB = 2 \text{ cm}$

since $\triangle AOB$ is equilateral

Arc $AB: l = r\theta$

$= 2 \times \frac{\pi}{3} = \frac{2\pi}{3} \text{ cm}$

\therefore Perimeter $= \left(\frac{2\pi}{3} + 2\right) \text{ cm}$.

Question 6

(a) $V = \pi \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \sec^2 x \, dx$
 $= 2\pi \int_0^{\frac{\pi}{3}} \sec^2 x \, dx$ (by symmetry)

$= 2\pi \left[\tan x \right]_0^{\frac{\pi}{3}}$

$= 2\pi \left(\tan \frac{\pi}{3} - \tan 0 \right)$

$= 2\pi (\sqrt{3} - 0)$

$= 2\sqrt{3}\pi \text{ units}^3$.

(b) (i) $Q = Ae^{-kt}$

When half of 1 unit remains,

$\frac{1}{2} = e^{-k \times 1600}$

$\ln \frac{1}{2} = \ln e^{-k \times 1600}$

$-1600k = \ln \frac{1}{2}$

$\therefore k = \frac{\ln 2^{-1}}{-1600}$

$= \frac{-\ln 2}{-1600}$

$= \frac{\ln 2}{1600}$

$\approx 4.3322 \times 10^{-4}$.

(ii) Let the safe level be A .

When $t = 0, Q = 3A$

$Q = 3Ae^{-kt}$

When $Q = A, t = ?$

$A = 3Ae^{-kt}$

$\frac{1}{3} = e^{-kt}$

$\ln \frac{1}{3} = \ln e^{-kt}$

$-kt = -\ln 3$

$\therefore t = \frac{1}{k} \ln 3$

$= \frac{1600}{\ln 2} \times \ln 3$ (from (i))

$= 2535.94 \dots$

\therefore The amount of radium will reach the safe level near the end of the 2536th year.

(c) (i) $y = ax^2 + bx$

$\frac{dy}{dx} = 2ax + b$

Since the gradient of RO is 1.2,

then at O , when $x = 0$,

$\frac{dy}{dx} = 1.2$

$\therefore 1.2 = 2a(0) + b$

$b = 1.2$

Since the gradient of PQ is -1.8 ,

then at P , when $x = 30$,

$\frac{dy}{dx} = -1.8$

$\therefore -1.8 = 2a(30) + 1.2$

$-1.8 = 60a + 1.2$

$-3 = 60a$

$a = -0.05$.

(ii) From part (i),

$y = -0.05x^2 + 1.2x$

Find x at the vertex.

METHOD 1

$\frac{dy}{dx} = -0.1x + 1.2$

At the vertex, $\frac{dy}{dx} = 0$

$\therefore -0.1x + 1.2 = 0$

$-0.1x = -1.2$

$x = 12$

METHOD 2

Using the axis of symmetry,

$x = \frac{-b}{2a}$

$= \frac{-1.2}{2(-0.05)}$

$= 12$

Return to the parabola.

Substitute $x = 12$ into

$y = -0.05x^2 + 1.2x$.

$y = -0.05(12)^2 + 1.2(12)$

$= 7.2$

\therefore Vertex is $(12, 7.2)$.

At P , when $x = 30$,

$y = -0.05(30)^2 + 1.2(30)$

$= -9$

$\therefore d = 7.2 + 9$


$= 16.2 \text{ m}$

\therefore Distance from the vertex of the parabola to the horizontal line through P is 16.2 m.

Question 7

- (a) (i) $\ddot{x} = 8e^{-2t} + 3e^{-t}$
 $\dot{x} = \int (8e^{-2t} + 3e^{-t}) dt$
 $\dot{x} = -4e^{-2t} - 3e^{-t} + c_1$
 When $t = 0, \dot{x} = -6$
 $-6 = -4e^0 - 3e^0 + c_1$
 $-6 = -4 - 3 + c_1$
 $1 = c_1$
 $\therefore \dot{x} = -4e^{-2t} - 3e^{-t} + 1$
 $x = \int (-4e^{-2t} - 3e^{-t} + 1) dt$
 $x = 2e^{-2t} + 3e^{-t} + t + c_2$
 When $t = 0, x = 5$
 $5 = 2e^0 + 3e^0 + 0 + c_2$
 $0 = c_2$
 $\therefore x = 2e^{-2t} + 3e^{-t} + t.$
- (ii) At rest, when $\dot{x} = 0,$
 $-4e^{-2t} - 3e^{-t} + 1 = 0$
 $4e^{-2t} + 3e^{-t} - 1 = 0$
 $4(e^{-t})^2 + 3e^{-t} - 1 = 0$
 Let $m = e^{-t}$
 $\therefore 4m^2 + 3m - 1 = 0$
 $(4m - 1)(m + 1) = 0$
 $\therefore m = \frac{1}{4}$ or $m = -1$
 $\therefore e^{-t} = \frac{1}{4}$ or $e^{-t} = -1$
 But $e^{-t} > 0$ for all real $t.$
 $\therefore e^{-t} = \frac{1}{4}$ (only)
 $-t = \ln\left(\frac{1}{4}\right)$
 $\therefore t = -\ln 4^{-1}$
 $= -1 \times -\ln 4$
 $= \ln 4$
 $\doteq 1.39$ s.

- (iii) At rest, when $t = \ln 4,$
 $x = 2e^{-2\ln 4} + 3e^{-\ln 4} + \ln 4$
 $= 2.2612\dots$
 $\doteq 2.26$ m.
- (b) (i) $h = 1 + 0.7 \sin \frac{\pi}{6} t$
 Period = $\frac{2\pi}{n}$
 $= \frac{2\pi}{\frac{\pi}{6}}$
 $= 12$ hours.
- (ii) **METHOD 1** Algebraic:
 Consider:
 $-1 \leq \sin \frac{\pi}{6} t \leq 1$
 $\therefore -0.7 \leq 0.7 \sin \frac{\pi}{6} t \leq 0.7$
 $\therefore 1 - 0.7 \leq 1 + 0.7 \sin \frac{\pi}{6} t \leq 1 + 0.7$
 $\therefore 0.3 \leq h \leq 1.7$
 \therefore At low tide, $h = 0.3$ m.
 When $h = 0.3,$
 $0.3 = 1 + 0.7 \sin \frac{\pi}{6} t$
 $-0.7 = 0.7 \sin \frac{\pi}{6} t$
 $-1 = \sin \frac{\pi}{6} t$
 $\frac{\pi}{6} t = \frac{3\pi}{2}$
 $\therefore t = 9$ hours after 5 am
 \therefore Low tide occurs at 2 pm.
- METHOD 2** Graphical:
 $h = 1 + 0.7 \sin \frac{\pi}{6} t$
 Period = 12
 Amplitude = 0.7
 Shift: up by 1




\therefore Low tide is 0.3 metres at 2 pm.

- (iii) When $h = 1.35,$
 $1.35 = 1 + 0.7 \sin \frac{\pi}{6} t$
 $0.35 = 0.7 \sin \frac{\pi}{6} t$
 $\frac{1}{2} = \sin \frac{\pi}{6} t$
 $\frac{\pi}{6} t = \frac{\pi}{6}$ or $\left(\pi - \frac{\pi}{6}\right)$
 $\frac{\pi}{6} t = \frac{\pi}{6}$ or $\frac{5\pi}{6}$
 $\therefore t = 1$ or 5 hours after 5 am.
 \therefore Ship can enter harbour between 1 hour after 5 am and 5 hours after 5 am
 i.e. from 6 am up to, and including, 10 am.

Question 8

- (a) (i) $-1 < x < 3.$
 (ii) For x large, $f'(x) \rightarrow 0.$
 (iii)



- (b) (i) $P = 350\,000$
 $r = \frac{9}{12} \% = 0.0075$
 $A_1 = 350\,000(1.0075) - 2937$
 $= \$349\,688.$
- (ii) $A_{12} = 346\,095$
 $r = \frac{6}{12} \% = 0.005$
 $n = 24 \times 12 = 288$
 $A_{288} \equiv 0$
 Substituting into the give formula and rearranging,
 $A_{288} = 346\,095 \times 1.005^{288}$
 $= M(1 + 1.005 + \dots + 1.005^{287})$
 $\therefore 0 = 346\,095(1.005)^{288}$
 $\frac{M(1.005^{288} - 1)}{0.005}$
 $M(1.005^{288} - 1)$
 $\frac{0.005}{0.005}$
 $= 346\,095(1.005)^{288}$
 $\therefore M = \frac{346\,095(1.005)^{288}(0.005)}{1.005^{288} - 1}$
 $= \$2270.31$ (nearest cent).
- (iii) $M = 2937, A_n = 0$
 From (ii),
 $A_n = 346\,095(1.005)^n$
 $\frac{M(1.005^n - 1)}{0.005}$
 $\therefore 0 = 346\,095(1.005)^n$
 $\frac{2937(1.005^n - 1)}{0.005}$
 $\frac{2937(1.005^n - 1)}{0.005}$
 $= 346\,095(1.005)^n$

$$2937(1.005)^n - 2937 = 1730.475(1.005)^n$$

$$(1.005)^n (2937 - 1730.475) = 2937$$

$$(1.005)^n = 2.4342\dots$$

$$n \ln 1.005 = \ln 2.4342\dots$$

$$n = \frac{\ln 2.4342\dots}{\ln 1.005}$$

$$\approx 178.373\ 317\ 5$$

∴ It will take 179 months to repay the \$346 095.

- (iv) Total repayments for (ii) $\approx 288 \times \$2270.31 = \$653\ 849.28$
 Total repayments for (iii) $\approx 178.373\ 317\ 5 \times \$2937 = \$523\ 882.43$ (nearest cent)
 ∴ Saving over term of loan = $\$653\ 849.28 - \$523\ 882.43 = \$129\ 966.85$.

Question 9

- (a) For one draw each:

$$P(\text{Van wins}) = \frac{1}{9}, \quad P(\text{Van loses}) = \frac{8}{9},$$

$$P(\text{Marie wins}) = \frac{1}{16}, \quad P(\text{Marie loses}) = \frac{15}{16}$$

For 3 draws each:

$$P(\text{at least 1 win}) = 1 - P(\text{no wins})$$


$$= 1 - \left[\left(\frac{8}{9} \right)^3 \times \left(\frac{15}{16} \right)^3 \right]$$

$$= 1 - \frac{125}{216}$$

$$= \frac{91}{216}$$

- (b) (i) Cost of PRS = $5 \times \$1000 + 3 \times \$2600 = \$12\ 800$.

(ii)



Pythagoras' theorem in ΔPSR :

$$PS^2 = 5^2 + 3^2$$

$$PS = \sqrt{34} \text{ km}$$

$$\text{Cost of } PS = \sqrt{34} \times \$2600$$


$$= \$15\ 160.47$$

(nearest cent).

(iii)

$$PQ = PR - QR$$

$$\therefore PQ = (5 - x) \text{ km}$$



Pythagoras' theorem in ΔQSR :

$$QS^2 = x^2 + 3^2$$

$$QS = \sqrt{x^2 + 9}$$

Cost in \$ of PQS:

$$C = (5 - x) \times 1000 + \sqrt{x^2 + 9} \times 2600$$

$$= 1000 \left(5 - x + 2.6\sqrt{x^2 + 9} \right)$$

(iv)
$$C = 1000 \left(5 - x + 2.6(x^2 + 9)^{\frac{1}{2}} \right)$$

$$\frac{dC}{dx} = 1000 \left(-1 + 1.3(x^2 + 9)^{\frac{1}{2}} \times 2x \right)$$

$$= 1000 \left(\frac{2.6x}{\sqrt{x^2 + 9}} - 1 \right)$$

Max/min when $\frac{dC}{dx} = 0$

i.e. $1000 \left(\frac{2.6x}{\sqrt{x^2 + 9}} - 1 \right) = 0$

$$\frac{2.6x}{\sqrt{x^2 + 9}} - 1 = 0$$

$$\frac{2.6x}{\sqrt{x^2 + 9}} = 1$$

$$2.6x = \sqrt{x^2 + 9} \quad \text{①}$$

$$6.76x^2 = x^2 + 9$$

$$5.76x^2 = 9$$

$$x^2 = 1.5625$$

$$x = \pm 1.25$$

Since $x > 0$, then $x = 1.25$

Test $x = 1.25$:

x	1	1.25	1.5
$\frac{dC}{dx}$	-177.8...	0	162.7...

∴ Minimum cost when $x = 1.25$

$$\therefore C = 1000 \left(5 - 1.25 + 2.6\sqrt{1.25^2 + 9} \right)$$

$$= 12\ 200$$

∴ Minimum cost of PQS = \$12 200.

- (v) If cost underwater changes from \$2600/km to \$1100/km, then equation ① in (iv) becomes

$$1.1x = \sqrt{x^2 + 9}$$

$$1.21x^2 = x^2 + 9$$

$$0.21x^2 = 9$$

$$x^2 = 42.8571\dots$$

$$x = \pm 6.5465\dots$$

Since $0 \leq x \leq 5$, minimum cost occurs at either $x = 0$ or $x = 5$.

When $x = 0$,

$$\text{Cost} = \text{cost}(PR) + \text{cost}(RS)$$

$$= 5 \times \$1000 + 3 \times \$1100$$

$$= \$8300$$

When $x = 5$,

$$\text{Cost} = \text{cost}(PS)$$

$$= \sqrt{34} \times \$1100$$

$$= \$6414.05 \text{ (nearest cent)}$$

∴ Cable should be laid directly from P to S for minimum cost.

Question 10

(a)
$$f(x) = x - \frac{x^2}{2} + \frac{x^3}{3}$$

$$f'(x) = 1 - \frac{2x}{2} + \frac{3x^2}{3}$$

$$= 1 - x + x^2$$

Turning point when $f'(x) = 0$

i.e. $0 = 1 - x + x^2$

$$\Delta = (-1)^2 - 4 \times 1 \times 1$$

$$= -3$$

$$< 0$$

$f'(x) = 0$ has no solutions

∴ $y = f(x)$ has no turning points.

(b)
$$f''(x) = -1 + 2x$$

Point of inflexion when $f''(x) = 0$

i.e. $0 = -1 + 2x$

$$x = \frac{1}{2}$$

x	0	$\frac{1}{2}$	1
$f''(x)$	-1	0	1

Concavity changes, ∴ point of inflexion.

When $x = \frac{1}{2}$, $f(x) = \frac{5}{12}$

∴ $y = f(x)$ has point of inflexion

at $\left(\frac{1}{2}, \frac{5}{12} \right)$.

$$\begin{aligned}
 \text{(c) (i)} \quad & 1-x+x^2 - \frac{1}{1+x} \\
 &= \frac{(1-x+x^2)(1+x)-1}{1+x} \\
 &= \frac{1+x-x-x^2+x^2+x^3-1}{1+x} \\
 &= \frac{x^3}{1+x}, \text{ as required.}
 \end{aligned}$$

$$\text{(ii)} \quad f'(x) = x^2 - x + 1 \text{ (from (a))}$$

$$g(x) = \ln(1+x)$$

$$g'(x) = \frac{1}{1+x}$$


$$f'(x) - g'(x) = x^2 - x + 1 - \frac{1}{1+x}$$

$$= \frac{x^3}{1+x} \text{ (from (i))}$$

$$\text{Since } x \geq 0, \frac{x^3}{1+x} \geq 0$$

$$\therefore f'(x) - g'(x) \geq 0$$

$$\therefore f'(x) \geq g'(x).$$



$$\begin{aligned}
 \text{(e)} \quad & \frac{d}{dx} [(1+x)\ln(1+x) - (1+x)] \\
 &= (1+x) \times \frac{1}{1+x} + \ln(1+x) \times 1 - 1 \\
 &= 1 + \ln(1+x) - 1 \\
 &= \ln(1+x).
 \end{aligned}$$

$$\begin{aligned}
 \text{(f)} \quad & \int_0^1 [f(x) - g(x)] dx \\
 &= \int_0^1 \left(x - \frac{x^2}{2} + \frac{x^2}{3} - \ln(1+x) \right) dx \\
 &= \left[\frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - [(1+x)\ln(1+x) - (1+x)] \right]_0^1 \\
 &= \left(\frac{1}{2} - \frac{1}{6} + \frac{1}{12} - 2\ln 2 + 2 \right) - (-\ln 1 + 1) \\
 &= \left(2\frac{5}{12} - 2\ln 2 \right) - 1 \\
 &= \left(1\frac{5}{12} - 2\ln 2 \right) \text{ units}^2 \\
 &= 0.03 \text{ units}^2 \text{ (to 2 decimal places).}
 \end{aligned}$$

End of Mathematics solutions