

## KAMBALA

### MATHEMATICS

#### Assessment Task #3

Monday 31<sup>st</sup> May 2004

Time allowed: 70 minutes

#### INSTRUCTIONS

- There are 4 questions of equal value.
- Marks for each part of a question are indicated
- All questions should be attempted.
- All necessary working should be shown
- Start each question on a new page
- Approved scientific calculators and drawing templates may be used

#### Question 1 (START A NEW PAGE)

12 MARKS

(a) Evaluate  $e^{-3}$  (to 3 significant figures) [1]

(b) Evaluate  $\log_2 16 - \log_3 \frac{1}{9}$  [2]

(c) Simplify  $3\log_a(x-y) - \log_a(x+y)$  [2]

(d) Differentiate

(i)  $y = e^{2x-3}$  [1]

(ii)  $y = \frac{\ln x}{e^x}$  [2]

(iii)  $y = (e^{-x} + e^x)^3$  [2]

(e) Find  $\int 5e^{2x+1} dx$  [2]

#### Question 2 (START A NEW PAGE)

12 MARKS

- (a) Find the equation of the tangent to the curve  $f(x) = x \log_e x$  at the point where  $x = 1$ . Leave your answer in general form. [3]

(b) If  $f(x) = \log_e \left( \frac{x}{2-x} \right)$  find  $f'(x)$  [2]

- (c) What is the domain of the function  $y = \log_e(x-2)$ ? [1]

- (d) Find the volume of the solid of revolution formed when the curve  $y = e^{-\frac{1}{2}x^2}$  from  $x = 0$  to  $x = 1$  is rotated about the  $x$  axis.

Leave your answer in exact form. [3]

- (e) Given that  $\int \frac{1}{x} dx = \ln x + c$

Show that the exact area under the curve  $y = \frac{1}{2x}$  between  $x = 1$  and  $x = 2$  is  $\ln\sqrt{2}$  units<sup>2</sup> [3]

---

Question 3 (START A NEW PAGE)

12 MARKS

- (a) (i) Show that  $\int_1^k 2xe^{x^2-1} dx = e^{k^2-1} - 1$  [2]

- (ii) Hence find the values of  $k$  for which  $\int_1^k 2xe^{x^2-1} dx = e^7 - 1$  (in simplest exact form) [2]

*Question 3 continues over the page*

- (b) Verity, Constance and Prudence are asked to approximate the area under the curve  $f(x) = xe^x$  from  $x = 1$  to  $x = 3$ .

- (i) Verity decides she will use the Trapezoidal rule with three function values to approximate the area. Show her working and answer, leaving your answer in terms of  $e$ . [3]

- (ii) Constance uses Simpson's Rule once to approximate the answer. Show her working and answer, leaving your answer in terms of  $e$ . [3]

- (iii) Prudence decides to work out the difference between her friends' two values. What is the difference between the two values? (to 2 decimal places) [2]

---

Question 4 (START A NEW PAGE)

12 MARKS

The function  $f(x)$  is given by  $f(x) = xe^x$

- (a) Show that  $f'(x) = e^x(x+1)$  [2]

- (b) Find any stationary point(s) and determine their nature [2]

- (c) Show that  $f(x)$  has a point of inflection at  $x = -2$  [2]

- (d) What happens to the curve as  $x$  get large? (ie what happens as  $x \rightarrow +\infty$  and  $x \rightarrow -\infty$ ) [2]

- (e) Sketch  $f(x)$  for  $-3 \leq x \leq 1$  [3]

- (f) On your sketch, shade in the region where the inequalities  $y \geq xe^x$  and  $x \geq 0$  both apply [1]

END OF TEST

## 12 Z-unit Task #3

31.5.04

Qn	Solutions	Marks	Comments+Criteria
1(a)	$e^{-3} = \frac{1}{e^3} = 0.049787$ $\approx 0.0498$ (3sf)	✓ (1)	✓ off if not to 3sf
1(b)	$\log_2 16 - \log_3 \frac{1}{9}$ $= 4 - (-2)$ $= 6$	✓✓ (2)	1 mark each
1(c)	$3 \log_a (x-y) - \log_a (x+y)$ $= \log_a (x-y)^3 - \log_a (x+y)$ $= \log_a \frac{(x-y)^3}{(x+y)}$	✓ (2)	
1(d)	i, $y = e^{2x-3}$ $y' = 2e^{2x-3}$ ii, $y = \frac{\ln x}{e^{2x}} = \frac{u}{v}$ $y' = \frac{e^x \cdot \frac{1}{x} - \ln x \cdot e^x}{(e^x)^2}$ $= \frac{e^x}{x} - e^x \ln x \left[ \frac{1}{x} - \ln x \right]$ iii, $y = (e^{-x} + e^x)^3$ $y' = 3(e^{-x} + e^x)^2 \cdot (-e^{-x} + e^x)$ $= 3(e^{-x} + e^x)^2 (e^x - e^{-x})$	✓ ✓✓ ✓✓ ✓✓ (5)	1 for recognition of quotient and a start on the process $\rightarrow \frac{1}{x} - \ln x$ or $\frac{1-x \ln x}{x e^x}$
1(e)	$\int 5 e^{2x+1} dx$ $= 5 \int e^{2x+1} dx$ $= \frac{5}{2} e^{2x+1} + C$	✓✓ (2)	For 2 mks must be $\frac{5}{2}$ not $\frac{1}{2} 5$


12

Qn	Solutions	Marks	Comments+Criteria
2(a)	$f(x) = x \ln x$ $f'(x) = x \cdot \frac{1}{x} + \ln x \cdot 1$ $= 1 + \ln x$		
2(b)	$f'(1) = 1 + \ln 1 = 1 + \ln 1 = 1$ pt is $x=1$ $y = 1 \ln 1$ ie $(1, 0)$ $= 0$ $\therefore$ eqn is $y-0 = 1(x-1)$ ie $y = x-1$	✓ (3)	1/3 Incorrect $f'(1)$ followed through correctly
2(c)	$f(x) = \ln \frac{x}{2-x}$ $= \ln x - \ln(2-x)$ $f'(x) = \frac{1}{x} - \frac{-1}{2-x}$ $= \frac{1}{x} + \frac{1}{2-x}$	✓ (2)	$= \frac{2}{x+2-2x}$
2(d)	Domain: $x-2 > 0$ $x > 2$	✓ (1)	$x \neq 2$ (1) $x < 2$
2(e)	$V = \pi \int_a^b y^2 dx$ $= \pi \int_0^1 e^{-2x} dx$ $= \pi \left[ -e^{-x} \right]_0^1$ $= \pi (-e^{-1} + e^0)$ $= \pi \left( 1 - \frac{1}{e} \right)$ unit <sup>3</sup>	✓ ✓ ✓ ✓ (3)	1/3 (Incorrect) or 2/3 (but correctly completed)

Qn	Solutions	Marks	Comments+Criteria								
(e)	$A = \int_1^2 \frac{1}{2x} dx$ $= \frac{1}{2} \int_1^2 \frac{1}{x} dx$ $= \frac{1}{2} [\ln x]_1^2$ $= \frac{1}{2} (\ln 2 - \ln 1)$ $= \frac{1}{2} \ln 2 = \ln \sqrt{2}$	✓ ✓ ✓ ✓ ✓									
3(a)	(i) $\int_1^k 2x e^{x^2-1} dx$ $= [e^{x^2-1}]_1^k$ $= e^{k^2-1} - e^0$ $= e^{k^2-1} - 1$ (ii) $k^2 - 1 = 7$ $k^2 = 8$ $k = \pm\sqrt{8} = \pm 2\sqrt{2}$	✓ ✓ ✓	Let $u = x^2 - 1$ $du = 2x dx$ $x=k, u=k^2-1$ $x=1, u=0$ $I = \int_0^{k^2-1} e^u du$ $= [e^u]_0^{k^2-1}$ $= e^{k^2-1} - e^0$ $= e^{k^2-1} - 1$								
(b)	$f(x) = x e^x$ $a=1 \rightarrow x=3$ <table border="1"> <tr> <td><math>x</math></td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td><math>f(x)</math></td> <td><math>e^1</math></td> <td><math>2e^2</math></td> <td><math>3e^3</math></td> </tr> </table> $A = \frac{b-a}{4} \left[ f(a) + 2f\left(\frac{a+b}{2}\right) + f(b) \right]$ $= \frac{3-1}{4} \left[ e + 4e^2 + 3e^3 \right]$	$x$	1	2	3	$f(x)$	$e^1$	$2e^2$	$3e^3$	✓ ✓	$h = \frac{b-a}{n} = \frac{3-1}{2} = 1$ $A = \frac{1}{2} [e + 3e^3 + 2.2e^2]$
$x$	1	2	3								
$f(x)$	$e^1$	$2e^2$	$3e^3$								

Qn	Solutions	Marks	Comments+Criteria
	$A = \frac{1}{2} (e + 4e^2 + 3e^3)$	✓	
	(ii) $A = \frac{b-a}{6} (f(a) + 4f\left(\frac{a+b}{2}\right) + f(b))$ $= \frac{3-1}{6} (e + 8e^2 + 3e^3)$ $= \frac{1}{3} (e + 8e^2 + 3e^3)$	✓✓ ✓	$A = \frac{1}{3} (e + 3e^3 + 4 \times 2e^2)$
	" Difference is		
	$\frac{1}{2}(e + 4e^2 + 3e^3) - \frac{1}{3}(e + 8e^2 + 3e^3)$ $= \frac{1}{6}e + (2e^2 - \frac{8}{3}e^2) + \frac{1}{6} \cdot 3e^3$ $= \frac{1}{6}e - \frac{2e^2}{3} + \frac{e^3}{2}$ $= 0.453\dots - 4.926\dots + 10.04\dots$ $= 5.569\dots \div 5.57$	✓✓	1 for post calculation
4(a)	$f(x) = x e^x = u \cdot v$ $f'(x) = x e^x + e^x \cdot 1$ $= e^x(x+1)$	✓ ✓	
(b)	SP's when $f'(x) = 0$ i.e. $e^x(x+1) = 0$ $\therefore x = -1$ as $e^x \neq 0$ $f''(x) = e^x(x+2) + e^x$ $= e^x(x+2)$ $f''(-1) = e^{-1}(-1+2) > 0$ $\therefore \min P^+$ at $x = -1$ ( $y = \frac{-1}{e}$ )	✓ ✓ ✓	

Qn	Solutions	Marks	Comments+Criteria								
	$\therefore \text{min pt is } x = -1$ $y = -1 e^{-1} = -\frac{1}{e}$ ie $(-1, -\frac{1}{e})$		Ignore y coord.								
(c)	$f''(x) = e^x(x+2)$ $f''(x) = 0 \text{ for POI}$ ie $e^x(x+2) = 0$ $\therefore x = -2$ <table border="1"> <tr> <td><math>x</math></td> <td>-1</td> <td>-2</td> <td>-3</td> </tr> <tr> <td><math>f''(x)</math></td> <td><math>e^{-1}</math></td> <td>0</td> <td><math>-e^{-3}</math></td> </tr> </table> <p>+ <math>\overset{0}{-}</math>          change in concavity  <math>\therefore (-2, -2e^{-2})</math>          is a POI</p>	$x$	-1	-2	-3	$f''(x)$	$e^{-1}$	0	$-e^{-3}$	✓ if not found before	
$x$	-1	-2	-3								
$f''(x)$	$e^{-1}$	0	$-e^{-3}$								
(d)	$\text{as } x \rightarrow \infty \quad xe^x \rightarrow \infty$ $x \rightarrow -\infty \quad xe^x \rightarrow 0$ from beneath $\therefore f(x) = xe^x \quad \text{try } x = -100$ $= -100 e^{-100}$ $= \frac{-100}{e^{100}} \quad \text{v.v. small + (-)}$ etc	✓ ✓									
(e)	See over										

Qn	Solutions	Marks	Comments+Criteria
4(e) cont			
(f)	 $f(x) = xe^x$ $(-3, -\frac{3}{e^3}), (-2, -\frac{2}{e^2}), \text{POI}, (-1, -\frac{1}{e}), \text{min}$	✓ correct region	