

Student ID: _____



KAMBALA

Mathematics Extension 2

HSC Assessment Task 1

February 2008

*Time Allowed: 50 minutes working time***Outcomes Assessed**

- E2** chooses appropriate strategies to construct arguments and proofs in both concrete and abstract settings
- E3** uses the relationship between algebraic and geometric representations of complex numbers
- E6** combines the ideas of algebra and calculus to determine the important features of the graphs of a wide variety of functions
- E9** communicates abstract ideas and relationships using appropriate notation and logical argument

INSTRUCTIONS

- This task contains 3 questions of 10 marks each. Marks for each part question are shown.
- Answer all questions on the writing paper provided. **Start each question on a new page.**
- Calculators may be used.
- Show all necessary working.
- Marks may be deducted for careless or badly arranged work.
- More marks will be allocated to questions involving higher order thinking.

Question 1 (Start a new page.)**10 Marks**

- (a) If $z = 2 + 3i$ and $w = 4 - i$, find:
- (i) $z + w$ 1
- (ii) iw 1
- (b) If $(1 - i)z = 3 - 4i$, find:
- (i) z 2
- (ii) $z\bar{z}$ 2
- (c) Given that \vec{p} is the vector that represents the complex number $-2 - 3i$ and \vec{q} is the vector that represents the complex number $1 - i$:
- (i) Show these vectors on an Argand diagram. 2
- (ii) Show the position vector $\vec{p} + \vec{q}$. 1
- (iii) What complex number does $\vec{p} + \vec{q}$ represent? 1

Question 2 (Start a new page.)**10 Marks**

- (a) Sketch $|z + i| \leq 2$ and describe the locus geometrically. 2
- (b) (i) Write $\sqrt{3} - i$ in mod-arg form. 2
- (ii) Hence find the two square roots of $\sqrt{3} - i$. 3
- (c) Solve $z^2 + (1 + i)z + 2i = 0$ expressing the roots in the form $a + ib$ where a and b are real. Let $z = x + iy$ where x and y are real. 3

Question 3 (Start a new page.)

10 Marks

- (a) (i) Sketch the graph of $y = \frac{2x}{x^2+1}$, indicating the co-ordinates of its stationary points. **3**
- (ii) What geometrical feature of the graph of the curve $y = \frac{2x}{x^2+1}$ shows that the function is odd? **1**
- (b) Consider the function $f(x) = 6x - x^2$:
On separate diagrams, sketch the graphs of the following:
- (i) $y = f(x)$, clearly showing the co-ordinates of any turning points. **1**
- (ii) $y = \sqrt{f(x)}$ **1**
- (iii) $y = |f(x)|$ **1**
- (iv) $y = \frac{1}{f(x)}$ **2**
- (v) $|y| = f(x)$ **1**

End of Assessment Task

10/10

19465128

Q1:

(a) $z = 2+3i$ $w = 4-i$
 (i) $z+w = 2+3i+4-i = 6+2i$ ✓


(ii) $iw = i(4-i) = 4i+1 = 1+4i$ ✓

(b) $(1-i)z = 3-4i$
 (i) $z = \frac{3-4i}{1-i} = \frac{(3-4i)(1+i)}{1+1} = \frac{3+3i-4i+4}{2} = \frac{7-i}{2}$

(ii) $z\bar{z} = \left(\frac{7}{2}\right)^2 + \left(\frac{1}{2}\right)^2 = \frac{49}{4} + \frac{1}{4} = \frac{50}{4} = 12\frac{1}{2}$ ✓

(c) (i) $\vec{p} = -2-3i$
 $\vec{q} = 1-i$


(ii) See diagram
 (iii) $\vec{p} + \vec{q} = -1-4i$ ✓



10/10

or.

(a) $|z+1| \leq 2$
 the locus is a circle w radius = 2 units & centre is $(-1, 1)$



(b) (i) $\sqrt{3}-i = 2 \cos(-\frac{\pi}{6})$ ✓ working?

(ii) let $z^2 = 2 \cos(-\frac{\pi}{6})$
 $\therefore z = 2^{\frac{1}{2}} \cos(\frac{-\frac{\pi}{6} + 2k\pi}{2})$ where $k=0, 1$

$z_1 = \sqrt{2} \cos(-\frac{\pi}{12})$

$z_2 = \sqrt{2} \cos(\frac{-\frac{\pi}{6} + 2\pi}{2})$
 $= \sqrt{2} \cos(\frac{-\frac{\pi}{6} + 12\pi}{2})$
 $= \sqrt{2} \cos \frac{11\pi}{12}$

show clearly

\therefore two square roots are $\sqrt{2} \cos(\frac{\pi}{12})$ & $\sqrt{2} \cos \frac{11\pi}{12}$

(c) $z^2 + (1+i)z + 2i = 0$
 $z = \frac{-(1+i) \pm \sqrt{(1+i)^2 - 4 \times 2i}}{2}$
 $= \frac{-1-i \pm \sqrt{1+2i+i^2-8i}}{2}$

(c) $z^2 + (1+i)z + 2i = 0$
 $z = \frac{-(1+i) \pm \sqrt{(1+i)^2 - 4 \times 1 \times 2i}}{2}$
 $= \frac{-1-i \pm \sqrt{1+2i+i^2-8i}}{2}$
 $= \frac{-1-i \pm \sqrt{-6i}}{2}$
 $= \frac{-1-i \pm \sqrt{3-\sqrt{3}i}}{2}$

let $z^2 = -6i = 6 \cos(\frac{\pi}{2})$
 $w^2 = -6i$
 $(a+ib)^2 = -6i$
 $a^2 + 2abi - b^2 = -6i$
 by equating real & imag parts,

$a^2 - b^2 = 0$ $2ab = -6$
 $a^2 - (\frac{-3}{a})^2 = 0$ $ab = -3$
 $b = \frac{-3}{a}$

$a^2 - \frac{9}{a^2} = 0$
 $a^4 - 9 = 0$
 $(a^2-3)(a^2+3) = 0$
 $a^2 = 3$ or $a^2 = -3$
 $a = \pm\sqrt{3}$ \therefore no solns since $a \neq 0$

$\therefore b = \mp \frac{3}{\sqrt{3}}$
 $= \mp \frac{\sqrt{3}}{1}$
 $= \mp \sqrt{3}$

$\therefore w = \pm (\sqrt{3} - \sqrt{3}i)$

$\frac{4+6}{10} = \frac{10}{10}$ well done
 $x^2 \neq -1$ Taylor

Q3.

(a) (i) $y = \frac{2x}{x^2+1}$
 $\frac{dy}{dx} = \frac{u'v - v'u}{v^2}$
 $= \frac{2(x^2+1) - 2x(2x)}{(x^2+1)^2}$
 $= \frac{2x^2+2-4x^2}{(x^2+1)^2}$
 $= \frac{-2x^2+2}{(x^2+1)^2}$


check limits!!

∴ stat pts occur when:
 $-2x^2+2=0$
 $2(-x^2+1)=0$
 $-x^2+1=0$
 $x^2=1$
 $x = \pm 1$
 when $x=1, y = \frac{2}{1+1} = 1$
 when $x=-1, y = \frac{-2}{2} = -1$

$\frac{3}{3} = \frac{0}{1} = 0$


x	0	1	2
y'	ve	0	-ve

x	-2	-1	0
y'	-ve	0	+ve





(b) $f(x) = 6x - x^2 = x(6-x)$

(i) $f(x) = 6x - x^2$
 $f'(x) = 6 - 2x$
 ∴ stat pts when $6 - 2x = 0$
 $6 = 2x$
 $3 = x$
 when $x = 3, y = 9$
 $f''(x) = -2$
 ∴ when $x = 3, f''(x) = -2 < 0$
 ∴ max tip




(ii) $y = f(x) = \sqrt{f(x)}$



1465128


(iv) $y = \frac{1}{f(x)}$



$y \neq 0$
 $x \neq 0, 6$

$\frac{2}{2}$

(v) $|y| = f(x)$
 $y = f(x)$ for $y \geq 0$
 $y = -f(x)$ for $y < 0$



1