

J.M.J.Ch.

MARCELLIN COLLEGE RANDWICK



YEAR 11 ACCELERATED

MATHEMATICS

HSC ASSESSMENT TASK # 2

2006

Weighting: 20% of HSC Assessment Mark.

STUDENT NAME: _____ MARK: _____ / 30

RANK ON THIS TASK: _____ /

Time Allowed: 50 minutes.

Directions:

- Answer all questions on separate lined paper.
- Show all necessary working.
- Marks may not be awarded for careless or badly arranged work.
- Begin your answers to each new question on a new answer page.

Structure: 2 questions each worth 15 marks -- Total 30 marks.

OUTCOMES TO BE ASSESSED:

- H1 – seeks to apply mathematical techniques to problems in a wide range of practical contexts
- H4 – expresses practical problems in mathematical terms based on simple given models
- H5 – applies appropriate techniques from the study of calculus and trigonometry to solve problems
- H6 – uses the derivative to determine the features of the graph of a function
- H7 – uses the features of a graph to deduce information about the derivative
- H8 – uses techniques of integration to calculate areas and volumes

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Question 1

a. If $y = e^x \cos x$, find $\frac{dy}{dx}$.

2

b.




Figure not to scale

PS and QR are arcs of concentric circles with O as the centre. Find the perimeter of the shaded region. Leave your answer in exact form.

2

c. i. Differentiate $\cos^3 x$ with respect to x .

1

ii. Hence or otherwise, find $\int_0^1 \cos^2 x \sin x \, dx$.

2

d. Solve the equation $2 \sin 2x + 1 = 0$ in the domain $0 \leq x \leq \pi$.

2

e. If $\frac{dy}{dx} = 3 \sin x + \sec^2 x$, find y in terms of x , if $y = 4$ when $x = 0$.

2

f. i. Sketch the graph of $y = \sin x$ and $y = \cos x$ on the same diagram in the domain

1

$$0 \leq x \leq \frac{\pi}{2}$$

ii. Find the coordinates of their point/s of intersection in this interval.

1

iii. Determine the area bounded by the curves and the x -axis for $0 \leq x \leq \frac{\pi}{2}$.

2

Question 2

a. The rate of decrease of water in a leaking container after t minutes is given by $\frac{dV}{dt} = -40 + 0.3t$. Initially, there was 2000 cm^3 in the container. Find how much water remained in the container after 10 minutes.

2

b. An isotope of carbon, C_{14} decays at a rate proportional to the mass present. The rate of change is given by $\frac{dM}{dt} = -kM$ where k is a positive constant and M is the mass present.

i. Show $M = M_0 e^{-kt}$ is a solution to this equation.

1

ii. The half-life of this isotope of C_{14} is 5600 years. This means it takes 5600 years for 100 grams of C_{14} to decay to 50 grams. Find the value of k correct to 3 significant figures.

2

iii. Archaeologists use radiocarbon dating to establish the age of discoveries. Calculate the age of an item in which only one-eighth of the original carbon remains.

2

c. A particle is moving in a straight line. It starts 2 metres to the right of a fixed point O on the line and at time t seconds its velocity $v \text{ ms}^{-1}$ is given by $v = 2 - 4e^{-t}$.

i. Find when the particle comes to rest.

1

ii. Find the distance travelled by the particle before it comes to rest.

2

d. A defective rocket rises vertically upwards and then crashes back to the ground. The rocket's height above the ground at time t seconds after take-off, is h metres, where $h = 12t^2 - 2t^3$.

i. When does the rocket crash?

1

ii. What is its velocity at this time?

1

iii. When is the velocity of the rocket zero?

1

iv. What is its maximum height?

2

Question 1


(13)

A $u = e^x$
 $\frac{du}{dx} = e^x$
 $u = \cos x = e^x (\cos x - \sin x)$
 $\frac{du}{dx} = -\sin x$

E $y = 4$ $z \neq 0$
 $y' = -3 \cos x + \tan x + 1$
 $4 = -3 + 1$
 $\Rightarrow y' = -3 \cos x + \tan x + 7$

B $(=R\theta)$ $(=R\theta)$
 $= \frac{15\pi}{3}$ $= \frac{25\pi}{3}$
 $= 5\pi$
 $P = \frac{5\pi + 25\pi}{3} = \frac{30\pi}{3} = 10\pi$
 $= 40\pi + 20$

F.I. y



$\cos x = \sin x$
 $\tan x = 1$
 $x = \frac{\pi}{4}$

where $x = \frac{\pi}{4}$ $y = \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$ or $\frac{\sqrt{2}}{2}$

\therefore P.O.I = $(\frac{\pi}{4}, \frac{1}{\sqrt{2}})$

C I $(\cos x)^3$
 $3(\cos x)^2(-\sin x)$
 $= -3 \sin x (\cos x)^2$

II $\int \cos^3 x dx$
 $= \int (\cos x)^2 \sin x dx$
 $= \int (1 - \sin^2 x) \sin x dx$
 $= \int \sin x dx - \int \sin^3 x dx$
 $= -\cos x + \frac{1}{3} \cos^3 x + C$

D $\sin 2x = -\frac{1}{2}$
 $2x = \frac{7\pi}{6}, \frac{11\pi}{6}$
 $x = \frac{7\pi}{12}, \frac{11\pi}{12}$

Question 1

F $\int_0^{\pi/4} \sin x + \int_{\pi/4}^{\pi/2} \cos x$
 $= [-\cos x]_0^{\pi/4} + [\sin x]_{\pi/4}^{\pi/2}$
 $= \left[-\frac{1}{\sqrt{2}} + 1\right] + \left[1 - \frac{1}{\sqrt{2}}\right]$
 $= \frac{1 - \sqrt{2}}{2} + \frac{1 - \sqrt{2}}{2}$
 $= \frac{2 - 2\sqrt{2}}{2} = 2 - 2\sqrt{2}$

* (i) (c) (ii) Since $\frac{d}{dx} [\cos^3 x] = -3 \sin x \cos^2 x$ from (i)
 $\therefore \int \frac{d}{dx} [\cos^3 x] dx = -3 \int \sin x \cos^2 x dx$
 $\therefore -\frac{1}{3} \cos^3 x + C = \int \cos^2 x \cdot \sin x dx$

Question 2

A $\Gamma = -40 + 0.3t$

(13) C.I. $T=0$ $\mathcal{R}=2$
 $V = 2 - 4e^{-t}$

$V=0$

$T=0$ $V=2000$

$2 - 4e^{-t} = 0$

$V = \frac{-40t + 0.3t^2 + C}{2}$

$4e^{-t} = 2$

$e^{-t} = \frac{1}{2}$

$2000 = 0 + 0 + C$

$t = -\ln(1/2) \checkmark$


$C = 2000$

$f = 0.69$

$V = \frac{-40t + 0.3t^2 + 2000}{2}$

II $V = 2 - 4e^{-t}$

$V = \frac{-40(10) + 0.3(10)^2 + 2000}{2}$
 $= \frac{-400 + 30 + 2000}{2}$
 $= \frac{1630}{2} = 815$



B $\frac{dm}{dt} = -k m e^{-kt}$
 as $m = m_0 e^{-kt}$
 $\therefore = -k m$

$a = 4e^t$
 $\int_0^{-\ln(1/2)} 4e^t \cdot 2 + 4e^t$
 $= 4e^t (2t + 1)$

Distance = Area under curve

II $50 = 100e^{-k \cdot 5600}$

$\ln 1/2 = -k \cdot 5600$

$k = \frac{-\ln(1/2)}{5600}$

$k = \frac{0.693}{5600} = 0.000124$

III $m = m_0 e^{-kt}$

$x=2$ $t=0$

$c=2$

$\therefore x = 2 + 4e^t + 2$

area under V curve gives distance

$V = 2 - 4e^{-t}$

$\therefore \int_0^{-\ln(1/2)} (2 + 4e^t) dt$
 $= 2t + 4e^t \Big|_0^{-\ln(1/2)}$
 $= 2 \ln 2 + 4 - [0 + 4]$
 $= 2 \ln 2$

$\ln(1/2) = -kt$

$t = \frac{\ln(1/2)}{-k} = 16800$ years

Question 2


I $h = 12t^2 - 2t^3$

$h=0$

$12t^2 - 2t^3 = 0$

$t^2(12 - 2t) = 0$

$t=0$ $t=6$ ✓



II $V = 24t - 6t^2$

$= 24(6) - 6(6)^2$

$= 144 - 216$

$= -72 \text{ ms}^{-1}$ ✓

III $V=0$

$24t - 6t^2 = 0$

$t(24 - 6t) = 0$

$t=0$ $t=4$ ✓

IV max height - turning pt

$v=0$

$v=0$ at $t=4$

when $t=4$ $h=64$

\therefore max height is $h=64$ at $t=4$