

Name..... Teacher.....



MORIAH COLLEGE

Year 12

MATHEMATICS EXTENSION 1 TRIAL

Date: Monday 18th August 2008.

Time Allowed: 2 hours plus 5 minutes reading time.

Examiners: O. Golan, L. Bornstein, G. Wagner

Instructions:

- Attempt ALL questions.
- All necessary working should be shown in every question.
- Marks may be deducted for careless or badly arranged work.
- Start each question in a new booklet
- All questions are of equal value
- Board approved calculators may be used.

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$


Note: $\ln x = \log_e x, \quad x > 0$

Total marks 84
 Attempt Questions 1–7
 All questions are of equal value
 Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

	Marks
Question 1 (12 Marks) Use a SEPARATE writing booklet.	
(a) Express $(\sqrt{2} - 1)^4$ in the form of $a\sqrt{2} + b$, where a and b are integers.	2
(b) Evaluate $\lim_{x \rightarrow 0} \frac{\sin 3x}{2x}$.	2
(c) The point $(6, 4)$ divides the interval joining $(4, 2)$ to $(9, 7)$ in the ratio of $1:k$. Calculate the value of k .	2
(d) Determine the exact value of $\sin^{-1}\left(\sin \frac{5\pi}{4}\right)$.	2
(e) Sketch the graph of the function: $y = \frac{x}{ x }$	2
(f) Determine the exact value of $\int_0^{\frac{\pi}{24}} \sin^2 6x dx$.	2

Question 2 (12 Marks) Use a SEPARATE writing booklet. Marks

- (a) Use the substitution $x = \cos \theta$ to evaluate $\int_{\frac{1}{2}}^1 \frac{\sqrt{1-x^2}}{x^2} dx$. 3
- (b) The graphs $y = x^2$ and $y = x + 6$ intersect at $(3, 9)$. 3
 Determine the size of the acute angle between the line and the curve at $(3, 9)$.
 Give your answer in radians correct to two decimal places.
- (c) (i) Determine the domain and range of $y = 1 + 2\sin^{-1}3x$. 2
 (ii) Sketch the graph of $y = 1 + 2\sin^{-1}3x$. 1
- (d)



The diagram shows unit circle centre O . Points $P(\cos \theta, \sin \theta)$ and $K(\cos \alpha, \sin \alpha)$ are on the circumference of the circle.

- (i) Use the cosine rule in $\triangle PKO$ to find an expression for $(PK)^2$. 1
- (ii) By using Pythagoras' theorem, the distance formula, or otherwise, determine a different expression for $(PK)^2$ than the expression in part (i). 1
- (iii) Hence show that $\cos(\theta - \alpha) = \cos \theta \cos \alpha + \sin \theta \sin \alpha$. 1

Marks

Question 3 (12 Marks) Use a SEPARATE writing booklet.

(a) Solve the inequality $\frac{x}{x-1} \geq 2$.

2

(b) A couple takes a \$300,000 bank loan to buy a home. The bank charges interest on the loan at a rate of 8.4% p.a, compounded monthly. If the couple wishes to fully repay the loan at the end of 25 years, calculate the amount of their monthly repayment.


4

(c) Use mathematical induction to prove that $3^{2n-1} + 5$ is divisible by 8, for all integers $n, n \geq 1$.

3

(d)

3



NOT TO SCALE

In the diagram AT is a tangent at T to the circle centre O . Points B, C and D lie on the circumference of the circle. $\angle BCO = 50^\circ$ and $\angle TDC = 106^\circ$.

Copy or trace this diagram into your writing booklet.

Determine the size of the obtuse angle $\angle TOC$ and the size of angle $\angle ATB$

Marks

Question 4 (12 Marks) Use a SEPARATE writing booklet.

(a) (i) Prove that $\tan \frac{\alpha}{2} = \frac{\sin \alpha}{1 + \cos \alpha}$.

2

(ii) Hence find the value of $\tan \frac{\alpha}{2}$, $\frac{\pi}{2} < \alpha < 2\pi$, when $\sin \alpha = \frac{3}{5}$.

2


(b) (i) Show that there is a root to the equation $2 \tan x + 2x - \pi = 0$ between $x = 0.6$ and $x = 0.75$.

1

(ii) Start with $x = 0.6$ and use one application of Newton's method to approximate the root to $2 \tan x + 2x - \pi = 0$ in $0 < x < \frac{\pi}{2}$.

2

(c)



NOT TO SCALE

The diagram shows a rectangle inscribed under one arch of the curve $y = \sin x$ in $0 < x < \pi$

(i) The coordinates of point A are $(x, \sin x)$.

1

Explain why the coordinates of point B are $(\pi - x, \sin x)$.

(ii) Show that the area $A(x)$ of the rectangle is given by $A(x) = (\pi - 2x) \sin x$.

1

(iii) Using question 4(b) part (ii), or otherwise, determine the dimensions of the rectangle with the largest area that can be inscribed under one arch of the graph of $y = \sin x$.

3


Question 5 (12 Marks) Use a SEPARATE writing booklet.

Marks

(a) One of the factors of $P(x) = ax^3 - 7x^2 + kx + 4$ is $(x - 4)$ and the remainder when $P(x)$ is divided by $(x - 1)$ is -6 .

- (i) Determine the values of a and k . 2
- (ii) Calculate the sum of the roots of $P(x)$. 1

(b)



The diagram shows the graph of the parabola $x^2 = 4y$ and the tangents at $T(2t, t^2)$ and $P(2p, p^2)$. The tangents intersect at point K .

- (i) Prove that the equation of the tangent at T is $y = tx - t^2$. 2
- (ii) Show that the coordinates of point K , the point where the tangents at T and P intersect are $(p + t, pt)$. 2
- (iii) The angle TKP is a right angle. 1

Show that the locus of K is a straight line.

Question 5 continues on page 7

Question 5 (continued)


Marks

(c) Andrew hit a golf ball with a velocity of 15 m/s at an angle of 50° to the ground. The ball just cleared a tree 10 m horizontally away from Andrew, as shown in the diagram below.

Place the origin at the position where the ball was hit.

You may assume the equations of motion, i.e. $y = vt \sin \theta - \frac{1}{2}gt^2$ and $x = vt \cos \theta$.

Assume the acceleration due to gravity is 10 m/s^2 .




- (i) Calculate the height of the tree in metres. Give your answer correct to one decimal place. 2
- (ii) How far beyond the tree did the ball hit the ground? 2

Marks

Question 6 (12 Marks) Use a SEPARATE writing booklet.

(a) 4



The diagram shows a water wheel that is rotating once every four minutes in an anti-clockwise direction. As the wheel rotates the marker moves up and down. The distance y represents the height of the marker point above the centre O of the water wheel.

The marker point is moving in Simple Harmonic Motion given by the equation :

$$\ddot{y} = -\left(\frac{\pi^2}{4}\right)y \text{ m/s}^2$$

Initially the marker point is at position A going into the water.

- (i) Find A , n and α if $y = A \sin(nt + \alpha)$. 2
- (ii) How far is the marker below the surface of the water at $t = 1$ minute? 2


- (b) Consider the function $f(x) = \frac{x}{x^2 - 4}$.
- (i) Find the domain of $y = f(x)$. 1
- (ii) Show that $y = f(x)$ is monotonically decreasing throughout the domain. 2
- (iii) Given that $y = f(x)$ is an ODD function, sketch the graph of $y = f(x)$ clearly labelling all essential features. 2
- (iv) Find the largest domain containing the origin for which $f(x)$ has an inverse function $f^{-1}(x)$. 1

- (v) Calculate $\int_{-1}^1 f(x) dx$ 2

Marks

Question 7 (12 Marks) Use a SEPARATE writing booklet.

(a)



Cooking oil is being filtered. The oil is in a container that is in the shape of a cone and it is dripping into a cylindrical container at a rate of $288\pi \text{ cm}^3/\text{min}$.


The height and radius of the cone are equal. The radius of the cylinder is 10 cm.

- (i) At what rate is the depth of the oil in the cone decreasing when the depth is 12 cm? 3
- (ii) At what rate is the depth of oil in the cylinder increasing when the depth of oil in the cone is 12 cm? 1

- (b) The velocity of a particle, v m/s, when it is x m from the origin is given by $v = e^{-2x}$. Initially the particle is at the origin.
- (i) Prove that $\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = \ddot{x}$ 1
- (ii) Find the acceleration of the particle at the origin. 1
- (iii) Prove that the displacement, x , from the origin at t seconds is given by:
 $x = \ln \sqrt{2t + 1}$ 2

Question 7 (continued)

- (c) The following net is used to construct a square pyramid $SABCD$ with base $ABCD$ and apex S . All the edges are of equal length.



In the resulting pyramid:

- i. show that the angle between the edge SA and the base $ABCD$ is 45° 2
- ii. calculate the angle between the face SBC and the base $ABCD$ to the nearest degree. 2

End of paper

Question 1	Sample answer	Syllabus outcomes and marking guide
(a)	$(\sqrt{2})^4 - 4(\sqrt{2})^3 + 6(\sqrt{2})^2 - 4(\sqrt{2}) + 1$ $= 4 - 8\sqrt{2} + 12 - 4\sqrt{2} + 1$ $= 17 - 12\sqrt{2}$	HE3 • Gives correct answer 2 • Shows the first line in the worked solution: 1
(b)	$\lim_{x \rightarrow 0} \frac{\sin Ax}{Ax} = 1$ $\lim_{x \rightarrow 0} \frac{\sin 3x}{2x} = \frac{3}{2} \lim_{x \rightarrow 0} \frac{\sin 3x}{3x}$ $= \frac{3}{2} \times 1 = \frac{3}{2}$	HE4 • Gives correct answer 2 • Uses $\lim_{x \rightarrow 0} \frac{\sin(Ax)}{Ax} = 1$ 1
(c)	$\frac{4k+9}{k+1} = 6$ and $\frac{2k+7}{1+k} = 4$ $4k+9 = 6k+6$ $k = 1.5$	P4 • Gives correct answer 2 • Determines a correct equation in k 1
(d)	$\sin^{-1}\left(\sin \frac{5\pi}{4}\right) = \sin^{-1}\left(\frac{-1}{\sqrt{2}}\right)$ $= \frac{\pi}{4}$	HE4 • Gives correct answer 2 • Obtains $\frac{5\pi}{4}$ or $\frac{\pi}{4}$ 1
(e)	$y = \frac{x}{ x }$ 	HE3 • Gives correct answer 2
(f)	$\int_0^{\frac{\pi}{24}} \sin^2 6x dx = \frac{1}{2} \int_0^{\frac{\pi}{24}} (1 - \cos 12x) dx$ $= \frac{1}{2} \left[x - \frac{1}{12} \sin 12x \right]_0^{\frac{\pi}{24}}$ $= \frac{\pi - 2}{48}$	H5 • Gives correct answer 2 • Obtains an expression involving $\int (1 - \cos 12x) dx$ 1

Question 2	Sample answer	Syllabus outcomes and marking guide
(a)	$x = \cos \theta$ $\frac{dx}{d\theta} = -\sin \theta$ $dx = -\sin \theta d\theta$ When $x = 1$, $\theta = 0$. When $x = \frac{1}{2}$, $\theta = \frac{\pi}{3}$. $\int_{\frac{\pi}{3}}^0 \frac{\sqrt{1 - \cos^2 \theta}}{\cos^2 \theta} \times -\sin \theta d\theta$ $= \int_0^{\frac{\pi}{3}} \tan^2 \theta d\theta$ $= \int_0^{\frac{\pi}{3}} (\sec^2 \theta - 1) d\theta$ $= [\tan \theta - \theta]_0^{\frac{\pi}{3}}$ $= \sqrt{3} - \frac{\pi}{3}$	HE6 • Gives correct answer 3 • Makes significant progress 2 • Makes some progress, e.g. determines the value of the limits in terms of θ or shows that $\frac{dx}{d\theta} = -\sin \theta$ 1
(b)	Gradient of the line $y = x + 6$ is 1. For $y = x^2$ $y' = 2x$ $m_T = 3 \times 2 = 6$ $\tan \theta = \left \frac{6 - 1}{1 + 6 \times 1} \right = \frac{5}{7}$ $\theta = 0.62$	P3 • Gives correct answer, ignore rounding ... 3 • Gives $\theta = \tan^{-1}\left(\frac{5}{7}\right)$ 2 • Determines that $m_1 = 6$ and $m_2 = 1$... 1

Question 2	(Continued)	Sample answer	Syllabus outcomes and marking guide
(c)	(i)	Domain: $-\frac{1}{3} \leq x \leq \frac{1}{3}$ Range: $1 - \pi \leq y \leq 1 + \pi$	HE4 • Both domain and range correct 2 • Either domain or range correct 1
	(ii)		HE4 • Draws correct graph 1
(d)	(i)	$\angle POK = \theta - \alpha$ $(PK)^2 = 1^2 + 1^2 - 2 \times 1 \times 1 \times \cos(\theta - \alpha)$ $= 2 - 2\cos(\theta - \alpha)$	PE2 • Gives the correct answer 1
	(ii)	 $(PK)^2 = (\cos \alpha - \cos \theta)^2 + (\sin \theta - \sin \alpha)^2$ $= \cos^2 \alpha + \cos^2 \theta - 2 \cos \alpha \cos \theta$ $+ \sin^2 \theta + \sin^2 \alpha - 2 \sin \theta \sin \alpha$ $= 2 - 2 \cos \alpha \cos \theta - 2 \sin \theta \sin \alpha$	PE2 • Gives the correct answer 1
	(iii)	$2 - 2 \cos(\theta - \alpha) = 2 - 2 \cos \alpha \cos \theta - 2 \sin \theta \sin \alpha$ $2 \cos(\theta - \alpha) = 2 \cos \alpha \cos \theta + 2 \sin \theta \sin \alpha$ $\cos(\theta - \alpha) = \cos \theta \cos \alpha + \sin \theta \sin \alpha$	PE2 • Gives a correct proof 1

Question 3	Sample answer	Syllabus outcomes and marking guide
(a)	$x \neq 1$ Multiply both sides by $(x-1)^2$. $x(x-1) \geq 2(x-1)^2$ $x(x-1) - 2(x-1)^2 \geq 0$ $(x-1)(2-x) \geq 0$ $\therefore 1 < x \leq 2$	PE3 • Gives correct solution 2 • Indicates that $x \neq 1$ 1
(b)	(i) $0.8 \times 0.8 = 0.64$	HE3 • Gives correct answer 1
	(ii) ${}^5C_3(0.8)^2(0.2)^3 = 0.0512$	HE3 • Gives correct answer, ignore rounding. . . 1
	(iii) Need zero, one or two women. ${}^{12}C_0(0.8)^{12} + {}^{12}C_1(0.8)^{11}(0.2) + {}^{12}C_2(0.8)^{10}(0.2)^2$ $= 0.558$	HE3 • Gives correct answer, ignore rounding . . 2 • Uses the correct terms in the appropriate binomial expansion 1
(c)	Test for $n = 1$: $3^{(2 \times 1) - 1} + 5 = 8$ which is divisible by 8. Thus it is true for $n = 1$. Assume 8 divides $3^{2k-1} + 5$. $\therefore 3^{2k-1} + 5 = 8\lambda$ where λ is an integer i.e. $3^{2k-1} = 8\lambda - 5$ Test for $n = k + 1$: $3^{2(k+1)-1} + 5 = 3^2 \times 3^{2k-1} + 5$ $= 9 \times (8\lambda - 5) + 5$ $= 72\lambda - 40$ $= 8(9\lambda - 5)$ $8(9\lambda - 5)$ is divisible by 8 as $9\lambda - 5$ is an integer. Thus if $3^{2n-1} + 5$ is divisible by 8 for an integer value of n , then it is divisible by 8 for the following integer value of n . Since it is divisible by 8 for $n = 1$ and $n = 2$, it is divisible by 8 for all positive integers n .	H5, HE7 • Gives a complete proof 3 • Provides a proof that lacks a minor component 2 • Makes some progress towards a proof, e.g. shows that $3^{2n-1} + 5$ is divisible by 8 for $n = 1$, and begins to test divisibility for $n = k + 1$ on the assumption that the expression is divisible by 8 for $n = k$. . . 1

Question 3	(Continued)	Sample answer	Syllabus outcomes and marking guide
(d)	$\angle TBC = 74^\circ$ $\angle TOC = 148^\circ$ Join TC in $\triangle TOC$. $\triangle TOC$ is isosceles as $OT = OC$ (radii). $\therefore \angle OCT = 16^\circ$ (Base angles of an isosceles triangle are equal) $\therefore \angle TCB = 66^\circ$ $\angle ATB = \angle TCB$ (The angle between a tangent and a chord is equal to the angle in the alternate segment.) $\therefore \angle ATB = 66^\circ$	(Opposite angles in a cyclic quadrilateral add up to 180° .) (The angle at the centre is twice the angle at the circumference, standing on the same arc.) $\triangle TOC$ is isosceles as $OT = OC$ (radii). $\therefore \angle OCT = 16^\circ$ (Base angles of an isosceles triangle are equal) $\therefore \angle TCB = 66^\circ$ $\angle ATB = \angle TCB$ (The angle between a tangent and a chord is equal to the angle in the alternate segment.) $\therefore \angle ATB = 66^\circ$	PE2 • Gives both angles correctly with supporting reasons 3 • Gives one angle found with supporting reasons and the other angle without supporting reasons 2 • Both angles found without supporting reasons. OR • One angle found with supporting reasons 1

Question 3

(b)
Monthly rate: $0.7\% = 0.007$
Number of monthly repayments: 300
Let P be the monthly repayment
Let A_n be the amount owing after n months.

So, $A_1 = 300000(1.007) - P$
 $A_2 = 300000(1.007)^2 - P(1.007) - P$
 $A_3 = 300000(1.007)^3 - P(1.007)^2 - P(1.007) - P$
 \dots
 $A_{300} = 300000(1.007)^{300} - P(1.007)^{299} - P(1.007)^{298} - \dots - P$
 $A_{300} = 300000(1.007)^{300} - P[(1.007)^{299} + (1.007)^{298} + \dots + 1]$
GP: $r = 1.007$ $a = 1$ $n = 300$

$$A_{300} = 300000(1.007)^{300} - P \frac{[(1.007)^{300} - 1]}{0.007}$$


Solve: $A_{300} = 0$

$$0 = 300000(1.007)^{300} - P \frac{[(1.007)^{300} - 1]}{0.007}$$

$$300000(1.007)^{300} = P \frac{[(1.007)^{300} - 1]}{0.007}$$

$$\frac{300000(1.007)^{300} \times 0.007}{[(1.007)^{300} - 1]} = P$$

$\$2,395.50 = P$ (nearest cent)

Question 4	Sample answer	Syllabus outcomes and marking guide
(a) (i)	$\tan \frac{\alpha}{2} = \frac{\sin \alpha}{1 + \cos \alpha}$ $\text{RHS} = \frac{2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}}{1 + \cos^2 \frac{\alpha}{2} - \sin^2 \frac{\alpha}{2}}$ $= \frac{2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}}{2 \cos^2 \frac{\alpha}{2}}$ $= \frac{\sin \frac{\alpha}{2}}{\cos \frac{\alpha}{2}}$ $= \tan \frac{\alpha}{2}$ $= \text{LHS}$ <p><i>Note: an alternate approach is to use the t results.</i></p>	P3, P4 • Gives a complete proof 2 • Uses two different half angle results appropriately 1
(ii)	$\sin \alpha > 0$ and $\frac{\pi}{2} < \alpha < 2\pi$. $\therefore \alpha$ is in the second quadrant.  <p>Using the identity proven in part (a) (i):</p> $\tan \frac{\alpha}{2} = \frac{3}{1 - \frac{4}{5}}$ $= 3$	P3, P4 • Gives correct answer 2 • Implies that $\cos \alpha = -\frac{4}{5}$ 1


Question 4	(Continued)	Sample answer	Syllabus outcomes and marking guide
(b)	(i)	$f(x) = 2 \tan x + 2x - \pi$ $f(0.6) = -0.57$ $f(0.75) = 0.22$ As there is a change in sign on a continuous curve there is a root between $x = 0.6$ and $x = 0.75$.	PE3, P6 • Gives a correct demonstration 1
	(ii)	$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$ $f'(x) = 2 \sec^2 x + 2$ $f'(0.6) = 4.93609$ $x_2 = 0.6 - \frac{-0.573319}{4.93609}$ $= 0.716148$ $= 0.71$ or 0.72	PE3, P6 • Obtains either 0.71, 0.7161... or 0.72. 2 • Makes progress towards a solution, e.g. finds the value of $f'(0.6)$ 1
(c)	(i)	The value where $y = \sin x$ crosses the x -axis is π . By symmetry, the length of the line joining π to the right side of the rectangle is x . Thus the x -coordinate is $\pi - x$. The y -coordinate is $\sin(\pi - x)$ and $\sin(\pi - x) = \sin x$.	P4 • Gives a correct demonstration 1
	(ii)	The length of AB is $\pi - 2x$. The height of the rectangle is $\sin x$. area = length \times breadth $= (\pi - 2x) \times \sin x$	H5 • Gives a correct demonstration 1
	(iii)	$\frac{dA}{dx} = -2 \sin x + (\pi - 2x) \cos x$ $-2 \sin x + \pi \cos x - 2x \cos x = 0$ $\cos x = 0$ is not a solution, so divide through by $\cos x$, $-2 \tan x + \pi - 2x = 0$ $2 \tan x + 2x - \pi = 0$ (the solution from part (b) (ii)) $x = 0.71$ Test: $\frac{d^2 A}{dx^2} = -2 \cos x - \pi \sin x - 2 \cos x + 2x \sin x$ $= -4 \cos x + (2x - \pi) \sin x$ $= -4.16$ when $x = 0.71$ The stationary point is a maximum. The dimensions of the rectangle are 1.72 by 0.65.	H5, PE3 • Provides the correct solution OR • Provides a solution correct with respect to part (b) (ii) 3 • Provides solution with minor errors or omissions, e.g. omits a test for maximum 2 • Makes some progress toward a correct solution, e.g. uses the product rule to correctly differentiate the equation and equates the derivative to zero. 1

Question 5	Sample answer	Syllabus outcomes and marking guide	
(a)	(i)	$P(x) = ax^3 - 7x^2 + kx + 4$ $P(4) = 64a - 112 + 4k + 4 = 0$ $64a + 4k = 108$ $16a + k = 27$ $P(1) = a - 7 + k + 4 = -6$ $a + k = -3$ $16a + k = 27$ $a + k = -3$ $15a = 30$ $a = 2, k = -5$	PE3 • Obtains the correct values for a and b ... 2 • Determines one correct equation relating a and b 1
	(ii)	sum of the roots $= \frac{7}{2}$ (Formula for the sum of roots $= \frac{b}{a}$.)	PE3 • Gives a correct answer with respect to part (a) (i) 1
(b)	(i)	$y = \frac{1}{4}x^2$ $y' = \frac{1}{2}x$ at $(2t, t^2)$, $y' = \frac{1}{2} \times 2t$ $y' = t$ $y - t^2 = t(x - 2t)$ $y = tx - 2t^2 + t^2$ $y = tx - t^2$	PE4 • Gives a correct proof 2 • Proves $y' = t$. OR • Derives the equation without proving $y' = t$ 1
	(ii)	solving $y = tx - t^2$ and $y = px - p^2$ simultaneously $px - p^2 = tx - t^2$ $px - tx = p^2 - t^2$ $x(p - t) = (p - t)(p + t)$ $p \neq t$ $x = p + t$ $y = t(p + t) - t^2$ $= tp + t^2 - t^2$ $= tp$ $K = (p + t, tp)$	PE4 • Gives correct answer 2 • Correctly demonstrates either the x coordinate or the y coordinate. OR • Makes an error in determining one of the coordinates then uses the wrong value to correctly determine the other value. 1
	(iii)	as $\angle TKP = 90^\circ$ $p \times t = -1$ $\therefore K$ is $(p + t, -1)$, which is a point on the line $y = -1$	PE4 • Gives a correct demonstration 1


Question 5 (Continued)

Sample answer

Syllabus outcomes and marking guide

(c) (i)	 <p> $y = 15 \sin 50t - 5t^2$ $x = 15 \cos 50t$ need the value of y when $x = 10$ $\therefore 10 = 15 \cos 50t$ $t = \frac{2}{3 \cos 50}$ $y = 15 \sin 50 \times \frac{2}{3 \cos 50} - 5 \times \left(\frac{2}{3 \cos 50}\right)^2$ $= 10 \tan 50 - 5 \left(\frac{2}{3 \cos 50}\right)^2$ $= 6.5391 \text{ m}$ $= 6.5 \text{ m to 1 decimal place}$ </p>	<p>HE3</p> <ul style="list-style-type: none"> Obtains a height of 6.5 m, ignore rounding 2 Determines the equation $10 = 15t \cos 50^\circ$ or equivalent 1
(ii)	<p>Need the value of x when $y = 0$.</p> <p>When $y = 0$, $-5t(t - 3 \sin 50^\circ) = 0$.</p> <p>$t = 3 \sin 50^\circ$ is required.</p> <p>Calculating x gives</p> <p>$x = 15 \cos 50^\circ \times 3 \sin 50^\circ$ $= 22.16 \text{ m}$</p> <p>The ball landed 22.16 m past the tree.</p>	<p>HE3</p> <ul style="list-style-type: none"> Gives correct answer, ignore rounding . . 2 <p>OR</p> <ul style="list-style-type: none"> Determines $x = 22.16$, ignore rounding. Correctly determines the distance past the tree using the range calculated by a valid method but including a minor error 1

#6
(a)



y represents the height above 0. Initially the marker point has negative vertical velocity as it is heading downwards. Hence, our equation could be $t \sin$.

① $y = -6 \sin\left(\frac{\pi t}{2} + \alpha\right)$ or ② $y = 6 \sin\left(\frac{\pi t}{2} + \alpha\right)$
with an appropriate value of α

① If $t=0 \rightarrow \sin \alpha = \frac{1}{2}$
 $y = -3$
 $\alpha = \frac{\pi}{6} + 2k\pi$ (as curve must be shifted completely to match)

$\therefore y = -6 \sin\left(\frac{\pi t}{2} + \frac{\pi}{6}\right)$ is easiest option.

② If $t=0 \rightarrow \sin \alpha = -\frac{1}{2}$
 $y = -3$
 $\alpha = \frac{7\pi}{6} + 2k\pi$ (again as in ①)

$\therefore y = 6 \sin\left(\frac{\pi t}{2} + \frac{7\pi}{6}\right)$ is easiest option

(b) Using either equation and substituting $t = 1$ gives


$y = -3\sqrt{3}$ ✓

Depth below water is $-3 - (-3\sqrt{3}) = (3\sqrt{3} - 3) \text{ m}$

Alternative solution:

In 1 minute the wheel will have rotated 90° in an anticlockwise direction.

$\sin \theta = \frac{3}{6}$
 $\theta = 30^\circ$
 $\beta = 60^\circ$
 $\phi = 30^\circ$
 $\frac{d}{6} = \cos 30^\circ$
 $d = 5.196$



$d - 3 = 2.2 \text{ metres}$

Alternative solution approach:

- Gives correct solution 4
- Determines that point A is approximately 5.2 metres below the centre of the wheel at $t = 1$ minute 3
- Determines the values of θ , β and ϕ 2
- Determines that the wheel has rotated through 90° at $t = 1$ minute 1

Question 6

(b)


$$f(x) = \frac{x}{x^2 - 4}$$

$$f'(x) = -\frac{(x^2 + 4)}{(x^2 - 4)^2}$$

(i) Domain: $x \neq 2, -2$

(ii) Both the denominator and the numerator are positive for all real values of x . So the derivative is negative for all real values of x , and, hence, monotonically decreasing.

(iii)



(iv) $-2 \leq x \leq 2$

(v) Since the function is odd, there is symmetry, and the integral is 0.


#7 (a) (i) $\frac{dV}{dt} = 288\pi \text{ cm}^3/\text{min}$

$$r = h$$

$$V = \frac{1}{3}\pi r^2 h$$

$$= \frac{1}{3}\pi h^3$$

$$\frac{dV}{dh} = \pi h^2$$



The value of $\frac{dh}{dt}$ when $h = 12$ is required.

$$\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt}$$

$$= \frac{1}{\pi h^2} \times 288\pi$$

$$= \frac{288}{144}$$

$$= 2 \text{ cm/min}$$

(ii) $\frac{dh}{dt}$ is constant.

$$V = \pi \times 10^2 \times h$$

In 1 minute:

$$288\pi = \pi \times 10^2 \times h$$

$$h = 2.88$$

HES

• Gives correct answer 3

• Makes significant progress towards the solution 2

• Obtains $\frac{dV}{dt} = 288\pi$.

OR

• Obtains $\frac{dV}{dh} = \pi h^2$.

OR

• Obtains $\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt}$ 1

HE3

• Gives correct solution 1

Question 7 (b)

(i) L.H.S. = $\frac{d}{dv} \left(\frac{v^2}{2} \right) \times \frac{dv}{dx}$

$$= v \times \frac{dv}{dx}$$

$$= \frac{dx}{dt} \times \frac{dv}{dx}$$

$$= \frac{dv}{dt}$$

$$= \ddot{x}$$

(ii) $\frac{v^2}{2} = \frac{e^{-4x}}{2}$

$$\dot{x} = \frac{d}{dx} \left(\frac{e^{-4x}}{2} \right)$$

$$\dot{x} = -2e^{-4x}$$

when $x = 0$, $\ddot{x} = -2 \text{ m/s}^2$

(iii) $\frac{dx}{dt} = \frac{1}{e^{2x}} \rightarrow \frac{dt}{dx} = e^{2x}$

$$\int dt = \int e^{2x} dx$$

$$t = \frac{1}{2}(e^{2x} - 1)$$




$$2t + 1 = e^{2x}$$

$$2x = \ln(2t + 1)$$

$$x = \frac{1}{2} \ln(2t + 1)$$

$$x = \ln \sqrt{2t + 1}$$

Question 7



Let the edge length be: $2a$

Let the centre of the square be O and the midpoint of BC , P

i. Consider the square $ABCD$:

Using Pythagoras' Theorem $AC = \sqrt{8a^2} = 2a\sqrt{2}$

O is the midpoint of AC , so $AO = a\sqrt{2}$



Now, $AS = 2a$

Call the angle α

$$\text{So } \frac{AO}{AS} = \cos \alpha$$

$$\cos \alpha = \frac{a\sqrt{2}}{2a} = \frac{\sqrt{2}}{2}$$

$$\alpha = 45^\circ$$



ii. Look at triangle SBP :

Using Pythagoras' Theorem: $SP = \sqrt{(2a)^2 - a^2} \Rightarrow SP = a\sqrt{3}$

Now, $OP = a$

Call the angle θ

In triangle SOP

$$\frac{OP}{SP} = \cos \theta$$

$$\frac{a}{a\sqrt{3}} = \cos \theta$$

$$\frac{1}{\sqrt{3}} = \cos \theta$$

$$\theta = 55^\circ$$