

BOARD OF STUDIES
NEW SOUTH WALES

2012

HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 1

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
Black pen is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11–14, show relevant mathematical reasoning and/or calculations

Total marks – 70

Section I Pages 3–6

10 marks

- Attempt Questions 1–10
- Allow about 15 minutes for this section

Section II Pages 7–14

60 marks

- Attempt Questions 11–14
- Allow about 1 hour and 45 minutes for this section

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10.

1 Which expression is a correct factorisation of $x^3 - 27$?

- (A) $(x - 3)(x^2 - 3x + 9)$
- (B) $(x - 3)(x^2 - 6x + 9)$
- (C) $(x - 3)(x^2 + 3x + 9)$
- (D) $(x - 3)(x^2 + 6x + 9)$

2 The point P divides the interval from $A(-2, 2)$ to $B(8, -3)$ internally in the ratio 3:2.

What is the x -coordinate of P ?

- (A) 4
- (B) 2
- (C) 0
- (D) -1


3 A polynomial equation has roots α , β and γ where

$$\alpha + \beta + \gamma = -2, \quad \alpha\beta + \alpha\gamma + \beta\gamma = 3 \quad \text{and} \quad \alpha\beta\gamma = 1.$$

Which polynomial equation has the roots α , β and γ ?

- (A) $x^3 + 2x^2 + 3x + 1 = 0$
- (B) $x^3 + 2x^2 + 3x - 1 = 0$
- (C) $x^3 - 2x^2 + 3x + 1 = 0$
- (D) $x^3 - 2x^2 + 3x - 1 = 0$

4 Which function best describes the following graph?



- (A) $y = 3\sin^{-1} 2x$
- (B) $y = \frac{3}{2}\sin^{-1} 2x$
- (C) $y = 3\sin^{-1} \frac{x}{2}$
- (D) $y = \frac{3}{2}\sin^{-1} \frac{x}{2}$

5 How many arrangements of the letters of the word OLYMPIC are possible if the C and the L are to be together in any order?

- (A) 5!
- (B) 6!
- (C) $2 \times 5!$
- (D) $2 \times 6!$

- 6 A particle is moving in simple harmonic motion with displacement x . Its velocity v is given by

$$v^2 = 16(9 - x^2).$$

What is the amplitude, A , and the period, T , of the motion?

- (A) $A = 3$ and $T = \frac{\pi}{2}$
 (B) $A = 3$ and $T = \frac{\pi}{4}$
 (C) $A = 4$ and $T = \frac{\pi}{3}$
 (D) $A = 4$ and $T = \frac{2\pi}{3}$

- 7 Which expression is equal to $\int \sin^2 3x \, dx$?

- (A) $\frac{1}{2} \left(x - \frac{1}{3} \sin 3x \right) + C$
 (B) $\frac{1}{2} \left(x + \frac{1}{3} \sin 3x \right) + C$
 (C) $\frac{1}{2} \left(x - \frac{1}{6} \sin 6x \right) + C$
 (D) $\frac{1}{2} \left(x + \frac{1}{6} \sin 6x \right) + C$

- 8 When the polynomial $P(x)$ is divided by $(x+1)(x-3)$, the remainder is $2x+7$.


What is the remainder when $P(x)$ is divided by $x-3$?

- (A) 1
 (B) 7
 (C) 9
 (D) 13

- 9 What is the derivative of $\cos^{-1}(3x)$?

- (A) $\frac{1}{3\sqrt{1-9x^2}}$
 (B) $\frac{-1}{3\sqrt{1-9x^2}}$
 (C) $\frac{3}{\sqrt{1-9x^2}}$
 (D) $\frac{-3}{\sqrt{1-9x^2}}$

- 10 The points A , B and P lie on a circle centred at O . The tangents to the circle at A and B meet at the point T , and $\angle ATB = \theta$.



What is $\angle APB$ in terms of θ ?

- (A) $\frac{\theta}{2}$
 (B) $90^\circ - \frac{\theta}{2}$
 (C) θ
 (D) $180^\circ - \theta$

Section II

60 marks

Attempt Questions 11–14

Allow about 1 hour and 45 minutes for this section

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (15 marks) Use a SEPARATE writing booklet.

(a) Evaluate $\int_0^3 \frac{1}{9+x^2} dx$. 3

(b) Differentiate $x^2 \tan x$ with respect to x . 2

(c) Solve $\frac{x}{x-3} < 2$. 3

(d) Use the substitution $u = 2 - x$ to evaluate $\int_1^2 x(2-x)^5 dx$. 3

(e) In how many ways can a committee of 3 men and 4 women be selected from a group of 8 men and 10 women? 1

(f) (i) Use the binomial theorem to find an expression for the constant term 2
in the expansion of $\left(2x^3 - \frac{1}{x}\right)^{12}$.

(ii) For what values of n does $\left(2x^3 - \frac{1}{x}\right)^n$ have a non-zero constant term? 1

Question 12 (15 marks) Use a SEPARATE writing booklet.

(a) Use mathematical induction to prove that $2^{3n} - 3^n$ is divisible by 5 for $n \geq 1$. 3

(b) Let $f(x) = \sqrt{4x-3}$.


(i) Find the domain of $f(x)$. 1

(ii) Find an expression for the inverse function $f^{-1}(x)$. 2

(iii) Find the points where the graphs $y = f(x)$ and $y = x$ intersect. 1

(iv) On the same set of axes, sketch the graphs $y = f(x)$ and $y = f^{-1}(x)$ showing the information found in part (iii). 2

(c) Kim and Mel play a simple game using a spinner marked with the numbers 1, 2, 3, 4 and 5.



The game consists of each player spinning the spinner once. Each of the five numbers is equally likely to occur.

The player who obtains the higher number wins the game.

If both players obtain the same number, the result is a draw.


(i) Kim and Mel play one game. What is the probability that Kim wins the game? 1

(ii) Kim and Mel play six games. What is the probability that Kim wins exactly three games? 2

Question 12 continues on page 9

Question 12 (continued)

- (d) Let $A(0, -k)$ be a fixed point on the y -axis with $k > 0$. The point $C(t, 0)$ is on the x -axis. The point $B(0, y)$ is on the y -axis so that $\triangle ABC$ is right-angled with the right angle at C . The point P is chosen so that $OBPC$ is a rectangle as shown in the diagram.



- (i) Show that P lies on the parabola given parametrically by 2

$$x = t \quad \text{and} \quad y = \frac{t^2}{k}.$$

- (ii) Write down the coordinates of the focus of the parabola in terms of k . 1

End of Question 12

Question 13 (15 marks) Use a SEPARATE writing booklet.

- (a) Write $\sin\left(2\cos^{-1}\left(\frac{2}{3}\right)\right)$ in the form $a\sqrt{b}$, where a and b are rational. 2

- (b) (i) Find the horizontal asymptote of the graph $y = \frac{2x^2}{x^2 + 9}$. 1

- (ii) Without the use of calculus, sketch the graph $y = \frac{2x^2}{x^2 + 9}$, showing the asymptote found in part (i). 2

- (c) A particle is moving in a straight line according to the equation

$$x = 5 + 6 \cos 2t + 8 \sin 2t,$$

where x is the displacement in metres and t is the time in seconds.

- (i) Prove that the particle is moving in simple harmonic motion by showing that x satisfies an equation of the form $\ddot{x} = -n^2(x - c)$. 2
- (ii) When is the displacement of the particle zero for the first time? 3

Question 13 continues on page 11

Question 13 (continued)

- (d) The concentration of a drug in the blood of a patient t hours after it was administered is given by

$$C(t) = 1.4te^{-0.2t},$$

where $C(t)$ is measured in mg/L.

- (i) Initially the concentration of the drug in the blood of the patient increases until it reaches a maximum, and then it decreases. 3

Find the time when this maximum occurs.


- (ii) Taking $t = 20$ as a first approximation, use one application of Newton's method to find approximately when the concentration of the drug in the blood of the patient reaches 0.3 mg/L. 2

End of Question 13

Question 14 (15 marks) Use a SEPARATE writing booklet.

- (a) The diagram shows a large semicircle with diameter AB and two smaller semicircles with diameters AC and BC , respectively, where C is a point on the diameter AB . The point M is the centre of the semicircle with diameter AB .

The line perpendicular to AB through C meets the largest semicircle at the point D . The points S and T are the intersections of the lines AD and BD with the smaller semicircles. The point X is the intersection of the lines CD and ST .



Copy or trace the diagram into your writing booklet.

- (i) Explain why $CTDS$ is a rectangle. 1
- (ii) Show that $\triangle MXS$ and $\triangle MXC$ are congruent. 2
- (iii) Show that the line ST is a tangent to the semicircle with diameter AC . 1


Question 14 continues on page 13

Question 14 (continued)

- (b) A firework is fired from O , on level ground, with velocity 70 metres per second at an angle of inclination θ . The equations of motion of the firework are

$$x = 70t \cos \theta \quad \text{and} \quad y = 70t \sin \theta - 4.9t^2. \quad (\text{Do NOT prove this.})$$

The firework explodes when it reaches its maximum height.




- (i) Show that the firework explodes at a height of $250 \sin^2 \theta$ metres. 2
- (ii) Show that the firework explodes at a horizontal distance of $250 \sin 2\theta$ metres from O . 1
- (iii) For best viewing, the firework must explode at a horizontal distance between 125 m and 180 m from O , and at least 150 m above the ground. 3

For what values of θ will this occur?

Question 14 continues on page 14

Question 14 (continued)

- (c) A plane P takes off from a point B . It flies due north at a constant angle α to the horizontal. An observer is located at A , 1 km from B , at a bearing 060° from B . Let u km be the distance from B to the plane and let r km be the distance from the observer to the plane. The point G is on the ground directly below the plane.



- (i) Show that $r = \sqrt{1 + u^2 - u \cos \alpha}$. 3
- (ii) The plane is travelling at a constant speed of 360 km/h. 2

At what rate, in terms of α , is the distance of the plane from the observer changing 5 minutes after take-off?

End of paper

2012 Higher School Certificate Solutions Mathematics Extension 1

SECTION I

Summary

1	C	4	C	7	C	9	D
2	A	5	D	8	D	10	B
3	B	6	A				

- 1 (C) $x^3 - 27 = x^3 - 3^3$
 $= (x-3)(x^2 + 3x + 9)$.
- 2 (A) $x = \frac{m \times x_2 + n \times x_1}{m+n}$
 $= \frac{3 \times 8 + 2 \times -2}{3+2}$
 $= \frac{20}{5}$
 $= 4$.
- 3 (B) $x^3 - \Sigma \alpha_i x^2 + \Sigma \alpha_i \beta_j x - \alpha \beta \gamma = 0$
 $x^3 - (-2)x^2 + (3)x - (1) = 0$
 $x^3 + 2x^2 + 3x - 1 = 0$.
- 4 (C) If the domain is $-2 \leq x \leq 2$ then
 $y = A \sin^{-1} \frac{x}{2}$ and if the range is
 $-\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$ then $y = 3 \sin^{-1} \frac{x}{2}$
- 5 (D) If L and C are together then there are 6 items to be arranged, then L and C can be arranged in 2 ways. The number of arrangements is $2 \times 6!$.


6 (A) $n = \sqrt{16} = 4, T = \frac{2\pi}{n} = \frac{2\pi}{4} = \frac{\pi}{2}$
 $A^2 = 9, A = 3$
 \therefore answer (A).

7 (C) $\int \sin^2 3x dx = \int \frac{1}{2}(1 - \cos 6x) dx$
 $= \frac{1}{2} \int (1 - \cos 6x) dx$
 $= \frac{1}{2} \left(x - \frac{1}{6} \sin 6x \right) + C$.

8 (D) $P(x) = (x+1)(x-3)Q(x) + 2x+7$
 $P(3) = (3+1)(3-3)Q(3) + 2 \times 3 + 7$
 $= (4)(0)Q(3) + 13$
 $= 13$.

9 (D) $\frac{d}{dx} \cos^{-1}(3x) = \frac{-3}{\sqrt{1-(3x)^2}}$
 $= \frac{-3}{\sqrt{1-9x^2}}$.

10 (B)



$TA = TB$ (tangents from an external point are equal)
 $\therefore \triangle TAB$ is isosceles.
 $\angle TBA + \angle TAB = 180^\circ - \theta$ (\angle sum of \triangle)
 $\therefore \angle TBA = 90^\circ - \frac{\theta}{2}$
 $\angle APB = 90^\circ - \frac{\theta}{2}$ (Alt. Seg. Theorem)

SECTION II

Question 11

(a) $\int_0^3 \frac{1}{9+x^2} = \left[\frac{1}{3} \tan^{-1} \frac{x}{3} \right]_0^3$
 $= \frac{1}{3} \left[\frac{\pi}{4} - 0 \right]$
 $= \frac{\pi}{12}$.

(b) $\frac{d}{dx} (x^2 \tan x) = 2x \tan x + x^2 \sec^2 x$.

(c) *Method 1:*

$\frac{x}{x-3} < 2$
 $x(x-3) < 2(x-3)^2$
 $x(x-3) - 2(x-3)^2 < 0$
 $(x-3)[x-2x+6] < 0$
 $(x-3)(-x+6) < 0$
 $\cdot x < 3 \text{ or } x > 6$.

OR

Method 2:

Boundary or critical points occur at discontinuities or at equalities.

There is a discontinuity at:

$x - 3 \neq 0$

$x \neq 3$


There is an equality at:

$\frac{x}{x-3} = 2$

$x = 2x - 6$

$x = 6$

Test points in the following regions.



Test $x = 0$: $\frac{0}{-3} < 2$ ✓

$x = 4$: $\frac{4}{1} < 2$ ✗

$x = 7$: $\frac{7}{4} < 2$ ✓

Hence $x < 3$ or $x > 6$.

(d)

Let $u = 2 - x \rightarrow x = 2 - u$

$du = -dx \rightarrow dx = -du$

At $x = 1, u = 1$

$x = 2, u = 0$

$\int_1^2 x(2-x)^5 dx = \int_1^0 (2-u)u^5 \cdot -du$
 $= \int_0^1 (2-u)u^5 du$
 $= \int_0^1 2u^5 - u^6 du$
 $= \left[\frac{2u^6}{6} - \frac{u^7}{7} \right]_0^1$
 $= \left(\frac{1}{3} - \frac{1}{7} \right) - (0)$
 $= \frac{4}{21}$

(e)

${}^8C_3 \times {}^{10}C_4 = 11760$.

(f) (i) *Method 1:*
 Each term is of the form ${}^{12}C_r (2x^3)^r (-x^{-1})^{12-r}$
 and $x^{3r} \times x^{-r(12)} = x^{4r-12}$.

The constant term is the coefficient of x^0 .
 Thus $4r - 12 = 0$
 $4r = 12$
 $r = 3$

${}^{12}C_3 (2x^3)^3 (-x^{-1})^9 = {}^{12}C_3 (8x^9) (-x^{-9})$
 $= {}^{12}C_3 \times 8 \times -1$
 $= -1760$

OR

Method 2:
 $\left(2x^3 - \frac{1}{x}\right)^{12} = \left(\frac{2x^4 - 1}{x}\right)^{12} = \frac{(2x^4 - 1)^{12}}{x^{12}}$

By inspection, the constant term is:
 $\frac{{}^{12}C_3 (2x^4)^3 (-1)^9}{x^{12}} = {}^{12}C_3 (8)(-1)$
 $= -1760$.

(ii) *Method 1:*
 From (i) replace the 12 with an n
 $x^{3r} \times x^{-r(12)} = x^{4r-n}$
 But $4r - n = 0$
 $4r = n$
 $n = 4r$

Thus n must be a multiple of 4.

OR

Method 2:

$$\left(2x^3 - \frac{1}{x}\right)^{12} = \left(\frac{2x^4 - 1}{x}\right)^{12} = \frac{(2x^4 - 1)^{12}}{x^{12}}$$

$$T_{r+1} = \frac{{}^n C_r (2x^4)^r (-1)^{n-r}}{x^n}$$

$$= \frac{{}^n C_r 2^r x^{4r} (-1)^{n-r}}{x^n}$$

For the constant term, the powers of x must be the same $\therefore n = 4r$.
 Thus n must be a multiple of 4.

Question 12

(a) Prove true when $n = 1$:
 $2^3 - 3 = 8 - 3$
 $= 5$ (which is divisible by 5)

Assume the following is true:
 $2^{3n} - 3^n = 5Q$ where Q is an integer

Thus $2^{3n} = 5Q + 3^n$

Need to prove that:
 $2^{3(n+1)} - 3^{n+1} = 5R$ where R is an integer

$$2^{3(n+1)} - 3^{n+1} = 2^3 \times 2^{3n} - 3 \times 3^n$$

$$= 8(5Q + 3^n) - 3 \times 3^n$$

$$= 40Q + 8 \times 3^n - 3 \times 3^n$$

$$= 40Q + 5 \times 3^n$$

$$= 5(8Q + 3^n)$$

$$= 5R$$


where $R = 8Q + 3^n$ is an integer.
 \therefore By the principle of mathematical induction, $2^{3n} - 3^n$ is divisible by 5 for all $n \geq 1$.

(b) (i) $4x - 3 \geq 0$
 $4x \geq 3$
 $x \geq \frac{3}{4}$
 \therefore The domain is $x \geq \frac{3}{4}$.

(ii) $f: y = \sqrt{4x - 3}$
 $f^{-1}: x = \sqrt{4y - 3} \quad (\because x \geq 0)$
 $x^2 = 4y - 3$
 $4y = x^2 + 3$
 $y = \frac{x^2 + 3}{4}$

(iii) $\frac{x^2 + 3}{4} = x$
 $x^2 + 3 = 4x$
 $x^2 - 4x + 3 = 0$
 $(x-1)(x-3) = 0$
 $x = 1, 3$

Since $y = x$, the points of intersection are (1,1) and (3,3).



(c) (i) *Method 1:*
 $P(\text{win}) = P(\text{loss})$
 $P(\text{draw}) = \frac{1}{5}$
 $P(\text{win}) + P(\text{loss}) = 1 - \frac{1}{5}$
 $P(\text{win}) = \frac{2}{5}$

OR

Method 2:

M e l

	1	2	3	4	5
1	d	-	-	-	-
2	w	d	-	-	-
3	w	w	d	-	-
4	w	w	w	d	-
5	w	w	w	w	d

w: Kim wins d: draw -: Kim loses

$$P(\text{win}) = \frac{10}{25} = \frac{2}{5}$$

(ii) ${}^6 C_3 \left(\frac{2}{5}\right)^3 \left(\frac{3}{5}\right)^3 = \frac{864}{3125} = 0.27648$.

(d) (i) $m_{AC} = \frac{k}{t} \quad m_{BC} = -\frac{y}{t}$

Since $BC \perp AC$, $m_{BC} \times m_{AC} = -1$

i.e. $-\frac{y}{t} \times \frac{k}{t} = -1$
 $-yk = -t^2$
 $y = \frac{t^2}{k}$

We have $C(t, 0)$ and $B\left(0, \frac{t^2}{k}\right)$.

Since $x_P = x_C$, $x_P = t$.

Since $y_P = y_B$, $y_P = \frac{t^2}{k}$.

$\therefore P = \left(t, \frac{t^2}{k}\right)$ as required.

(ii) $x = t, y = \frac{t^2}{k}$

$y = \frac{x^2}{k}$

$ky = x^2$

$x^2 = ky$

This is of the form $x^2 = 4ay$

$\therefore 4a = k$


$a = \frac{k}{4}$

\therefore the focus is $\left(0, \frac{k}{4}\right)$.

Question 13

(a) Let $\alpha = \cos^{-1} \frac{2}{3}$

i.e. $\cos \alpha = \frac{2}{3}$




$$\begin{aligned} \sin\left(2\cos^{-1}\frac{2}{3}\right) &= \sin(2\alpha) \\ &= 2\sin\alpha\cos\alpha \\ &= 2 \times \frac{\sqrt{5}}{3} \times \frac{2}{3} \\ &= \frac{4\sqrt{5}}{9} \end{aligned}$$

(b) (i) $y = \frac{2x^2}{x^2+9}$

$$= \frac{2}{1+\frac{9}{x^2}}$$

As $x \rightarrow \infty, y \rightarrow 2$
 \therefore the horizontal asymptote is $y = 2$.

- (ii) By inspection:
- y is an even function, therefore the graph is symmetrical about the x -axis
 - $y \geq 0$ for all x
 - The graph passes through the origin.



(c) (i) $x = 5 + 6\cos 2t + 8\sin 2t$
 $\dot{x} = -12\sin 2t + 16\cos 2t$
 $\ddot{x} = -24\cos 2t - 32\sin 2t$
 $= -4(6\cos 2t + 8\sin 2t)$
 $= -4(x-5)$
 Thus $\ddot{x} = -n^2(x-c)$
 where $n = 2$ and $c = 5$.

(ii) Solve for $x=0$:
 i.e. $6\cos 2t + 8\sin 2t = -5$ ①
 Writing LHS as $R\cos(2t-\alpha)$
 where $R > 0$ and $0^\circ \leq \alpha < 90^\circ$:
 $6\cos 2t + 8\sin 2t = R\cos(2t-\alpha)$
 $= R\cos 2t \cos \alpha + R\sin 2t \sin \alpha$

Equating coefficients:

$$\begin{aligned} R\cos\alpha &= 6 & \text{②} \\ R\sin\alpha &= 8 & \text{③} \end{aligned}$$

②²+③²:

$$\begin{aligned} R^2\cos^2\alpha + R^2\sin^2\alpha &= 6^2 + 8^2 \\ R^2(\cos^2\alpha + \sin^2\alpha) &= 6^2 + 8^2 \\ R^2 &= 6^2 + 8^2 \\ R &= 10 \quad [R > 0] \end{aligned}$$

③ ÷ ②:

$$\begin{aligned} \frac{\sin\alpha}{\cos\alpha} &= \frac{4}{3} \\ \tan\alpha &= \frac{4}{3} \\ \alpha &= \tan^{-1}\frac{4}{3} \quad (0 \leq \alpha < 90) \end{aligned}$$

Thus
 $6\cos 2t + 8\sin 2t = 10\cos(2t - \tan^{-1}\frac{4}{3})$

Returning to equation ①:

$$\begin{aligned} 10\cos(2t - \tan^{-1}\frac{4}{3}) &= -5 \\ \cos(2t - \tan^{-1}\frac{4}{3}) &= -\frac{1}{2} \\ 2t - \tan^{-1}\frac{4}{3} &= \frac{2\pi}{3} \\ 2t &= \frac{2\pi}{3} + \tan^{-1}\frac{4}{3} \\ t &= \frac{\pi}{3} + \frac{1}{2}\tan^{-1}\frac{4}{3} \\ t &= 1.51084516... \\ t &\approx 1.5 \text{ seconds} \end{aligned}$$

(d) (i) $C'(t) = \frac{d}{dt} 1.4te^{-0.2t}$
 $= 1.4e^{-0.2t} + 1.4te^{-0.2t}(-0.2)$
 $= 1.4e^{-0.2t}(1-0.2t)$

For $C'(t)$ to be 0:
 $1.4e^{-0.2t}(1-0.2t) = 0$
 $1-0.2t = 0$
 $t = 5$

Table of gradients:

t	4	5	6
$C'(t)$	0.13	0	-0.08

\therefore maximum when $t = 5$ hours.

(ii) We need to solve $1.4te^{-0.2t} = 0.3$.

That is, $1.4te^{-0.2t} - 0.3 = 0$.

Let $f(t) = 1.4te^{-0.2t} - 0.3$

Then from (i):

$$f'(t) = 1.4e^{-0.2t}(1-0.2t)$$

Newton's method to approximate

$f(t) = 0$:

$$t_2 = t_1 - \frac{f(t_1)}{f'(t_1)}$$

Here, $t_1 = 20$


$$f(20) = 0.21283788...$$

$$f'(20) = -0.076925683...$$

$$\begin{aligned} \therefore t_2 &= 20 - \frac{0.212...}{-0.076...} \\ &\approx 22.8 \text{ hours.} \end{aligned}$$

Question 14

(a) (i)



$AD \perp DB$ (angle in semicircle)
 $AS \perp SC$ (angle in semicircle)
 $\therefore \angle DSC = 90^\circ$ (straight angle)
 $CT \perp TB$ (angle in semicircle)
 $\therefore \angle CTD = 90^\circ$ (straight angle)

$\therefore \angle SCT = 90^\circ$ (\angle sum of quadrilateral)
 $\therefore CTDS$ is a rectangle (4 right angles)

(ii) MX is common
 $MS = MC$ (radii of same circle)
 $SX = CX$ (diagonals of rectangle bisect one another)
 $\triangle MXS \equiv \triangle MXC$ (SSS)

(iii) $\angle MSX = \angle MCX$ (corresponding \angle s, congruent \triangle s)
 $\therefore \angle MSX = 90^\circ$ (since $\angle MCX = 90^\circ$)
 $\therefore ST$ is a tangent (tangent \perp radius)

(b) (i) $y = 70t \sin \theta - 4.9t^2$
 $\dot{y} = 70 \sin \theta - 9.8t$
 At maximum height, $\dot{y} = 0$
 $70 \sin \theta - 9.8t = 0$
 $t = \frac{70 \sin \theta}{9.8}$

$$\begin{aligned} y &= 70 \left(\frac{70 \sin \theta}{9.8} \right) \sin \theta - 4.9 \left(\frac{70 \sin \theta}{9.8} \right)^2 \\ &= \frac{4900 \sin^2 \theta}{9.8} - \frac{4.9 \times 4900 \sin^2 \theta}{9.8^2} \\ &= 500 \sin^2 \theta - 250 \sin^2 \theta \\ &= 250 \sin^2 \theta \end{aligned}$$


\therefore the maximum height is $250 \sin^2 \theta$ metres.

(ii) When $t = \frac{70 \sin \theta}{9.8}$,
 $x = 70t \cos \theta$
 $= 70 \left(\frac{70 \sin \theta}{9.8} \right) \cos \theta$
 $= 500 \sin \theta \cos \theta$
 $= 250 \times 2 \sin \theta \cos \theta$
 $= 250 \sin 2\theta$
 \therefore the horizontal distance from O is $250 \sin 2\theta$ metres.

- (iii) The firework explodes when $x = 250 \sin 2\theta$ and $y = 250 \sin^2 \theta$.

For $125 \leq x \leq 180$:
 $125 \leq 250 \sin 2\theta \leq 180$
 $0.5 \leq \sin 2\theta \leq 0.72$

The graph of $y = \sin 2\theta$ below shows two parts satisfying this inequality.



By calculation:
 $30^\circ \leq 2\theta \leq 46^\circ$ or $134^\circ \leq 2\theta \leq 150^\circ$

$15^\circ \leq \theta \leq 23^\circ$ or $67^\circ \leq \theta \leq 75^\circ$
 For $y \geq 150^\circ$ and $0^\circ \leq \theta \leq 90^\circ$:

$250 \sin^2 \theta \geq 150$
 $\sin^2 \theta \geq 0.6$
 $\sin \theta \geq 0.774\dots$ [$0^\circ \leq \theta \leq 90^\circ$]

As $\sin \theta$ is increasing in this domain,

$\theta \geq 50.76\dots^\circ$

Thus we have $\theta \geq 50.76\dots^\circ$ and
 $(15^\circ \leq \theta \leq 23^\circ$ or $67^\circ \leq \theta \leq 75^\circ)$

To satisfy both conditions:

$67^\circ \leq \theta \leq 75^\circ$.

- (c) (i) To find r , values for PG and AG are needed.

$\sin \alpha = \frac{PG}{u} \rightarrow PG = u \sin \alpha$

To find AG , BG is needed.

$\cos \alpha = \frac{BG}{u} \rightarrow BG = u \cos \alpha$

Using the cosine rule in $\triangle GBA$:

$AG^2 = BG^2 + 1^2 - 2 \times BG \times \cos 60^\circ$

$= BG^2 + 1^2 - 2 \times BG \times \frac{1}{2}$

$= BG^2 + 1^2 - BG$

$= u^2 \cos^2 \alpha + 1 - u \cos \alpha$

In $\triangle APG$:

$r^2 = PG^2 + AG^2$

$= (u \sin \alpha)^2 + (u^2 \cos^2 \alpha + 1 - u \cos \alpha)$

$= u^2 \sin^2 \alpha + u^2 \cos^2 \alpha + 1 - u \cos \alpha$

$= u^2 (\sin^2 \alpha + \cos^2 \alpha) + 1 - u \cos \alpha$

$= u^2 + 1 - u \cos \alpha$

$\therefore r = \sqrt{1 + u^2 - u \cos \alpha}$.

- (ii) Given $\frac{du}{dt} = 360 \text{ km/h}$ and $t = 5 \text{ min}$

($\therefore u = 30 \text{ km}$), we need to find $\frac{dr}{dt}$.

$r = \sqrt{1 + u^2 - u \cos \alpha}$

$= (1 + u^2 - u \cos \alpha)^{\frac{1}{2}}$

$\frac{dr}{du} = \frac{1}{2} (1 + u^2 - u \cos \alpha)^{-\frac{1}{2}} \cdot (2u - \cos \alpha)$

$= \frac{2u - \cos \alpha}{2\sqrt{1 + u^2 - u \cos \alpha}}$

As $u = 30$,

$\frac{dr}{du} = \frac{60 - \cos \alpha}{2\sqrt{901 - 30 \cos \alpha}}$

Now $\frac{dr}{dt} = \frac{dr}{du} \times \frac{du}{dt}$

$= \frac{60 - \cos \alpha}{2\sqrt{901 - 30 \cos \alpha}} \times 360$

$= 180 \left(\frac{60 - \cos \alpha}{\sqrt{901 - 30 \cos \alpha}} \right) \text{ km/h.}$