



Sydney Girls High School

12 MAY

2010
TRIAL HIGHER SCHOOL
CERTIFICATE EXAMINATION

Mathematics Extension 2

General Instructions

- Reading Time - 5 minutes
- Working time - 3 hours
- Attempt ALL questions
- ALL questions are of equal value
- All necessary working should be shown in every question. Marks may be deducted for careless or badly arranged work.
- Standard integrals are supplied
- Board-approved calculators may be used.
- Diagrams are not to scale
- Each question attempted should be started on a new page. Write on one side of the paper only.

This is a trial paper ONLY. It does not necessarily reflect the format or the content of the 2010 HSC Examination Paper in this subject.

2 0 0 5 9 9 1 5

Candidate Number

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Question One (15 marks)

- a) Find $\int \frac{\sin x}{\cos^3 x} dx$. 2
- b) Find $\int \frac{4x^3 - 2x^2 + 1}{2x - 1} dx$. 3
- c) i) Find A , B and C given that $\frac{4x - 6}{(x + 1)(2x^2 + 3)} = \frac{A}{x + 1} + \frac{Bx + C}{2x^2 + 3}$. 3
- ii) Hence, find $\int \frac{4x - 6}{(x + 1)(2x^2 + 3)} dx$. 1
- d) Find $\int \sin^{-1} x dx$. 3
- e) Find $\int \frac{1}{3 + 2 \cos \theta} d\theta$. 3

Question Two (15 marks)


- a) If $z = \sqrt{3} - i$ and $w = 1 + i$, find:
- i) $z w$ 1
- ii) $\arg z$ 1
- iii) $|w'|$ 1
- iv) $\operatorname{Im}\left(\frac{z}{w}\right)$ 2
- b) $OPQR$ is a rectangle on the Argand diagram labelled anti-clockwise where O represents the origin and point P represents the complex number $3 + 4i$. Find the complex number representing Q and R given that $PQ = 2QR$. 2
- c) i) Find the square roots of $21 + 20i$. 2
- ii) Hence, solve $(1 + i)z^2 + z - 5 = 0$. 3
- d) The complex number z is such that $|z - 1| = \operatorname{Re}(z)$.
- i) Find the cartesian equation of the locus of z . 2
- ii) Find the range of values of $|z|$. 1

Question Three (15 marks)


Marks

- a) The region bounded by $y = \log_e x$, $y = 1$ and $x = 3$ is rotated about the y axis.
- i) Sketch this region on the number plane. 1
- ii) Find the volume formed using the method of cylindrical shells. 3

- b) A solid is formed with the base and top both rectangles parallel to each other and 6 cm apart. The dimensions of the base are 11 cm and 15 cm and the dimensions of the top are 7 cm and 10 cm. If all other faces are trapeziums, find the volume of the solid. 5



- c) An object of mass m is lying on an inclined plane at an angle θ to the horizontal. As shown in the diagram below, the object is subject to a gravitational force mg , a normal reaction force N and a frictional force F .



The object is not moving.

Resolve the forces acting on the object, and hence find an expression for


$\frac{F}{N}$ in terms of θ . 3

- d) Find $\frac{dy}{dx}$ given $x^3 + x^2y^4 = 0$. 3

Question Four (15 marks)

Marks

- a) The diagram shows the graph of $y = f(x)$.



Draw separate one-third page sketches of the graphs of the following :

- i) $y = |f(x)|$ 1
- ii) $y = f(|x|)$ 1
- iii) $y = [f(x)]^2$ 2
- iv) $y = e^{f(x)}$ 2
- v) $y = \frac{1}{f(x)}$ 3

- b) The equation $x^3 + 3x^2 - 2x - 2 = 0$ has roots α , β and γ . Find the equation with roots $\frac{2\alpha}{\beta\gamma}$, $\frac{2\beta}{\alpha\gamma}$ and $\frac{2\gamma}{\alpha\beta}$. 3

- c) Determine the greatest and least values of $\arg(z)$ if $|z - 4i| = 2$. 3

Question Five (15 marks)


Marks

- a) Given $1 - i$ is a root of $x^3 - 3x^2 + 4x - 2 = 0$ find the other roots. 2
- b) For the equation $x^4 + x^3 - 16x^2 - 4x + 48 = 0$, the product of two of the roots is 6.
- i) Hence, express the equation in the form $(x^2 + ax + b)(x^2 + cx + d) = 0$. 3
- ii) Find the roots of the equation. 1
- c) i) Given $x = \alpha$ is a double root of the equation $ax^4 + 4bx + c = 0$, deduce that $\alpha^3 = -\frac{b}{a}$. 2
- ii) Also, deduce that $ac^3 = 27b^4$. 3
- iii) Hence or otherwise, solve the equation $27x^4 - 32x + 16 = 0$, given that it has a double root. 4

Question Six (15 marks)

Marks

- a) $P\left(p, \frac{1}{p}\right)$ and $Q\left(q, \frac{1}{q}\right)$ are two points on the rectangular hyperbola $xy = 1$.
- i) Derive the equation of the chord PQ and show that it can be expressed in general form as $x + pqy - (p + q) = 0$. 2
- ii) Hence, show that the area of $\triangle OPQ$ is $\frac{|p^2 - q^2|}{2|pq|}$ units². 4
- b) The point $P(x_1, y_1)$ lies on the hyperbola $\frac{x^2}{16} - \frac{y^2}{9} = 1$.
The tangent at P cuts the x -axis at T .




- i) Find the coordinates of the foci S and S' . 1
- ii) Show that the equation of the tangent at P is given by $\frac{xx_1}{16} - \frac{yy_1}{9} = 1$. 3
- iii) Show that $\frac{S'T}{ST} = \frac{S'P}{SP}$. 3
- iv) Hence, deduce that $\angle S'PT = \angle SPT$. 2

Question Seven (15 marks)

Marks

- a) A particle of mass m kg is attached to one end of a light string at B . The other end of the string is fixed at a point A . The particle rotates in a horizontal circle of radius r metres at g rad/s, the centre of the circle being directly below A .



The forces acting on the particle are the tension in the string T and the gravitational force mg .

Let $\angle BAO = \theta$.

- i) Show that $T \sin \theta = mg^2 r$. 1
- ii) Prove that $\theta = \tan^{-1}(gr)$. 1
- iii) Prove that $T = mg\sqrt{1+g^2 r^2}$. 2
- b) i) Use De Moivre's Theorem to express $\cos 4\theta$ and $\sin 4\theta$ as powers of $\cos \theta$ and $\sin \theta$. 2
- ii) Hence show that $\tan 4\theta = \frac{4t - 4t^3}{1 - 6t^2 + t^4}$ where $t = \tan \theta$. 1
- iii) By first solving the equation $\tan 4\theta = 1$ for $0 \leq \theta \leq 2\pi$, solve the equation $x^4 + 4x^3 - 6x^2 - 4x + 1 = 0$. 3
- iv) Hence find the value of $\tan \frac{\pi}{16} \tan \frac{3\pi}{16} \tan \frac{5\pi}{16} \tan \frac{7\pi}{16}$. 2
- c) Evaluate $\int_0^{\pi} x \cos 2x \, dx$. 3

Question Eight (15 marks)

Marks

- a) i) Use integration by parts to show that $I_n = \frac{n-1}{n} I_{n-2}$ for $n \geq 2$ given $I_n = \int_0^{\frac{\pi}{2}} (\sin x)^n \, dx$ where n is an integer ($n \geq 0$). 3
- ii) Deduce $I_{2n} = \frac{2n-1}{2n} \times \frac{2n-3}{2n-2} \times \dots \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2}$. 3
- b) Given the identity $\cos(A+B) + \cos(A-B) = 2 \cos A \cos B$, solve the equation $\cos 5x + \cos 3x + 2 \cos x = 0$ for $0 \leq x \leq \frac{\pi}{2}$. 3
- c) Two sides of a triangle are of length $2x$ cm and $3x$ cm. The angles opposite these sides differ by 45° . Show that the smaller of the two angles is given by $\tan^{-1}\left(\frac{2+3\sqrt{2}}{7}\right)$. 4
- d) The positive integers are bracketed as follows $(1), (2,3), (4,5,6), (7,8,9,10), \dots$. The n th bracket has n integers. Prove that the sum of the integers in the n th bracket is $\frac{n}{2}(n^2+1)$. 2

End of paper

BLANK PAGE

QUESTION 1

a. Let $u = \cos x$

$$\frac{du}{dx} = -\sin x$$

$$\begin{aligned} \int \frac{\sin x}{\cos^3 x} dx &= \int -u^{-3} du \\ &= -\frac{u^{-2}}{-2} + C \\ &= \frac{1}{2\cos^2 x} + C \end{aligned}$$

b.

$$\frac{2x^2}{2x-1} \sqrt{4x^3-2x^2+1}$$

$$\frac{4x^3-2x^2}{4x^3-2x^2} \cdot 0+1$$

$$\begin{aligned} \int \frac{4x^3-2x^2+1}{2x-1} dx &= \int \left(2x^2 + \frac{1}{2x-1} \right) dx \\ &= \frac{2x^3}{3} + \frac{1}{2} \ln(2x-1) + C \end{aligned}$$

c.

i. $4x - 6 = A(2x^2 + 3) + (Bx + C)(x+1)$

let $x = -1$
 $-10 = 5A$
 $\therefore A = -2$
 $0 = 2A + B$
 $\therefore B = 4$
 $-6 = 3A + C$
 $\therefore C = 0$

ii.

$$\begin{aligned} \int \frac{4x-6}{(x+1)(2x^2+3)} dx &= \int \frac{-2}{x+1} + \frac{4x}{2x^2+3} \\ &= -2\ln(x+1) + \ln(2x^2+3) + C \end{aligned}$$

d.

let

$$u = \sin^{-1} x \quad dv = dx$$

$$u' = \frac{1}{\sqrt{1-x^2}} \quad v = x$$

$$\int \sin^{-1} x dx = x \sin^{-1} x - \int \frac{x}{\sqrt{1-x^2}} dx$$

Let

$$u = 1 - x^2$$

$$du = -2x dx$$

$$-\frac{1}{2} du = x dx$$

$$\begin{aligned} x \sin^{-1} x - \int \frac{x}{\sqrt{1-x^2}} dx &= x \sin^{-1} x - \int \frac{-du}{2\sqrt{u}} \\ &= x \sin^{-1} x + \frac{1}{2} \int u^{-\frac{1}{2}} \\ &= x \sin^{-1} x + \frac{1}{2} \times 2u^{\frac{1}{2}} \\ &= x \sin^{-1} x + u^{\frac{1}{2}} \\ &= x \sin^{-1} x + \sqrt{1-x^2} \end{aligned}$$

e.

$$\int \frac{1}{3+2\cos\theta} d\theta = \int \frac{1}{3+2\left(\frac{1-t^2}{1+t^2}\right)} \cdot \frac{2dt}{1+t^2}$$

$$= \int \frac{1}{5+t^2} \cdot \frac{2dt}{1+t^2}$$

$$= \int \frac{2}{5+t^2} dt$$

$$= 2 \left(\frac{1}{\sqrt{5}} \tan^{-1} \frac{t}{\sqrt{5}} \right) + C$$

$$= 2 \left(\frac{1}{\sqrt{5}} \tan^{-1} \frac{\tan \frac{\theta}{2}}{\sqrt{5}} \right) + C$$

QUESTION 2

a.

i.

$$\begin{aligned} zw &= (\sqrt{3}-i)(1+i) \\ &= \sqrt{3}-i^2+i(\sqrt{3}-1) \\ &= \sqrt{3}+1+i(\sqrt{3}-1) \end{aligned}$$

ii.

$$\begin{aligned} \arg(z) &= -\tan^{-1} \left(\frac{1}{\sqrt{3}} \right) \\ &= -\frac{\pi}{6} \end{aligned}$$

iii.

$$\begin{aligned} |w| &= \sqrt{1^2+1^2} \\ &= \sqrt{2} \\ |w^7| &= |w|^7 \\ &= (\sqrt{2})^7 \\ &= 8\sqrt{2} \end{aligned}$$

iv.

$$\begin{aligned} \operatorname{Im} \left(\frac{z}{w} \right) &= \operatorname{Im} \left(\frac{\sqrt{3}-i}{1+i} \times \frac{1-i}{1-i} \right) \\ &= -\frac{\sqrt{3}+1}{2} \end{aligned}$$

c.

i.

$$\begin{aligned} \sqrt{21+20i} &= a+ib \\ 21+20i &= a^2+2abi-b^2 \\ a^2-b^2 &= 21 \\ 2ab &= 20 \\ a &= \pm 5 \\ b &= \pm 2 \\ \therefore & \pm(5+2i) \end{aligned}$$

ii.

$$\begin{aligned} z &= \frac{-1 \pm \sqrt{1^2+20(1+i)}}{2(1+i)} \\ &= \frac{-1 \pm (5+2i)}{2(1+i)} \\ &= \frac{4+2i}{2(1+i)} \quad \text{or} \quad \frac{-6-2i}{2(1+i)} \\ &= \frac{2+i}{1+i} \quad \text{or} \quad \frac{-3-i}{1+i} \end{aligned}$$

d.

i.

$$|z-1| = \operatorname{Re}(z)$$


If $z = x+iy$

$$\begin{aligned} |(x-1)+iy| &= x \\ \sqrt{(x-1)^2+y^2} &= x \\ (x-1)^2+y^2 &= x^2 \\ x^2-2x+1+y^2 &= x^2 \end{aligned}$$

$$y^2 = 2x-1 \quad \text{or} \quad x = \frac{1}{2}(y^2+1)$$

ii.

$$|z| \geq \frac{1}{2}$$




ii) $V_{\text{rot}} = \pi(R^2 - r^2)$
 $= \pi((x+5)^2 - x^2)(y-1)$
 $= 2\pi x(y-1)dx$
 $V_{\text{rot}} = \int_{x=0}^{x=e} 2\pi x(y-1)dx$

Let $u = \log x$, $v' = x$
 $u' = \frac{1}{x}$, $v = \frac{x^2}{2}$
 $V_{\text{rot}} = 2\pi \int_0^1 [L_{\log x} \cdot \frac{x^2}{2}]' - \int_0^1 x^2 \cdot [-\frac{x^2}{2}]'$
 $= 2\pi \int_0^1 (x \log x - 2\pi \frac{x^2}{2})' dx$
 $= 2\pi \int_0^1 x(\log x - x) dx$

$V_{\text{rot}} = 2\pi \left([L_{\log x} \cdot \frac{x^2}{2}]' - \int_0^1 x^2 \cdot [-\frac{x^2}{2}]' \right)$
 $= 2\pi \left(\frac{1}{2} \log x \cdot x^2 - \frac{1}{2} \int_0^1 x^2 \cdot x^2 dx \right)$
 $= 2\pi \left(\frac{1}{2} \log 3 - \frac{1}{2} \int_0^1 x^4 dx \right)$
 $= 2\pi \left(\frac{1}{2} \log 3 - \frac{1}{2} \cdot \frac{1}{5} \right)$
 $= \pi \left(\log 3 - \frac{1}{5} \right)$

$V_{\text{rot}} = \int_a^b \pi(y^2 - x^2) dx$
 $= \int_0^e \pi \left(\frac{1}{x^2} - x^2 \right) dx$
 $= \pi \left(\int_0^e \frac{1}{x^2} dx - \int_0^e x^2 dx \right)$
 $= \pi \left(\left[-\frac{1}{x} \right]_0^e - \left[\frac{x^3}{3} \right]_0^e \right)$
 $= \pi \left(-\frac{1}{e} - \frac{e^3}{3} \right)$
 $= -\frac{\pi}{3} \left(\frac{e^3}{3} + \frac{1}{e} \right)$
 $= -\frac{\pi}{9} \left(e^3 + \frac{1}{e} \right)$
 $= -\frac{\pi}{9} \left(\frac{e^4 + 1}{e} \right)$
 $= -\frac{\pi}{9} \frac{e^4 + 1}{e}$
 $= -\frac{\pi}{9} \frac{685}{e}$

$\int_0^e \frac{1}{x^2} dx = \left[-\frac{1}{x} \right]_0^e = -\frac{1}{e}$
 $\int_0^e x^2 dx = \left[\frac{x^3}{3} \right]_0^e = \frac{e^3}{3}$
 $\int_0^e \frac{1}{x^2} dx - \int_0^e x^2 dx = -\frac{1}{e} - \frac{e^3}{3}$
 $= -\frac{1}{e} - \frac{e^3}{3}$
 $= -\frac{e^3 + 1}{3e}$
 $= -\frac{685}{3e}$








$F \sin \theta + N \cos \theta - mg = 0$
 $N \sin \theta - F \cos \theta = 0$

$F \sin^2 \theta + N \sin \theta \cos \theta = mg \sin \theta$
 $N \sin \theta \cos \theta - F \cos^2 \theta = 0$

$F \sin \theta \cos \theta + N \cos^2 \theta = mg \cos \theta$
 $N \sin^2 \theta - F \sin \theta \cos \theta = 0$


$F = \frac{mg \sin \theta}{\cos \theta}$
 $N = mg \cos \theta$

d) $3x^2 + x^2 \sqrt{4x^3 dy + 2xy^2} = 0$
 $dy = \frac{-2xy^2}{4x^2 y^2} dx$
 $= -\frac{2y^2 + 3x}{4x y^3}$



(a) $2x^2 = 2x$
 $x^2 = x$
 $x^2 - x = 0$
 $x(x-1) = 0$
 $x = 0$ or $x = 1$

$(x^2 + 3)(x^2 - 2\sqrt{x} - 2) \geq 0$
 $x^2 + 3 > 0$
 $x^2 - 2\sqrt{x} - 2 \geq 0$
 $x^2 - 4\sqrt{x} + 4\sqrt{x} - 12\sqrt{x} + 9\sqrt{x} - 13\sqrt{x} + 13\sqrt{x} - 4 \geq 0$



Gradient at $x = \frac{2}{\sqrt{3}} + \frac{2}{\sqrt{3}}$
 $= \frac{4}{\sqrt{3}}$

Local at $x = \frac{1}{\sqrt{3}} - \frac{1}{\sqrt{3}}$
 $= 0$

Question Five

(a) If $1-i$, then $1+i$ must also be a root.

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

$$\alpha + 1 - i + 1 + i = -\frac{(-3)}{1}$$

$$\alpha + 3 - 2 = 1$$

Hence, roots are $1-i, 1+i, 1$.

(b)(i) $\alpha\beta = 6$

$$\alpha\beta\gamma\delta = 48 \Rightarrow \gamma\delta = 8$$

$$(x^2 + ax + 6)(x^2 + cx + 8) = 0$$

coefficient of x^3 $a + c = 1$

coefficient of x $8a + 6c = -4$

$$8a + 8c = 8$$

$$2a = -10 \Rightarrow a = -5, c = 6$$

$$(x^2 - 5x + 6)(x^2 + 6x + 8) = 0$$

(b)(ii) $(x-3)(x-2)(x+4)(x+2) = 0$

$$\therefore x = 3, 2, -4 \text{ or } -2$$

(c)(i) $P(x) = ax^4 + 4bx + c$

$$P'(x) = 4ax^3 + 4b$$

double root at $x = \alpha$

$$\Rightarrow P(\alpha) = P'(\alpha) = 0$$

$$4a\alpha^3 + 4b = 0$$

$$4a\alpha^3 = -4b$$

$$\alpha^3 = \frac{-4b}{4a} = -\frac{b}{a}$$

(c)(ii) $P(\alpha) = 0$

$$a\alpha^4 + 4b\alpha + c = 0$$

$$\alpha(a\alpha^3 + 4b) = -c$$

$$\alpha\left(a \times \frac{b}{a} + 4b\right) = -c$$

$$\alpha(-b + 4b) = -c$$

$$\alpha^3(3b) = -c^3$$

$$-\frac{b}{a} \times 27b^3 = -c^3$$

$$\therefore ac^3 = 27b^4$$

(c)(iii) $a = 27, b = -8, c = 16$

$$\alpha^3 = \frac{8}{27} \therefore \alpha = \frac{2}{3}$$

$$27x^4 - 32x + 16 = (3x-2)^2(3x^2 + 4x + 4)$$

$$x = \frac{-4 \pm \sqrt{16 - 4(12)}}{6} = \frac{-4 \pm \sqrt{-32}}{6}$$

$$\therefore x = \frac{2}{3}, \frac{2}{3}, \frac{-2 \pm 2\sqrt{2}i}{3}$$

Question Six

(a)(i) $m_{PQ} = \frac{\frac{1}{p} - \frac{1}{q}}{\frac{p}{p-q} - \frac{q}{pq(p-q)}} = -\frac{1}{pq}$

$$y - \frac{1}{p} = -\frac{1}{pq}(x - p)$$

$$pqy - q = -x + p$$

$$x + pqy - (p + q) = 0$$

(a)(ii) height (distance from O to PQ)

$$h = \frac{|0 + (pq)0 - (p + q)|}{\sqrt{1^2 + (pq)^2}}$$

$$= \frac{|p + q|}{\sqrt{1 + p^2q^2}}$$

$$PQ = \sqrt{(p-q)^2 + \left(\frac{1}{p} - \frac{1}{q}\right)^2}$$

$$= \sqrt{(p-q)^2 + \frac{1}{(pq)^2}(q-p)^2}$$

$$= \sqrt{(p-q)^2 \left(1 + \frac{1}{p^2q^2}\right)} \text{ as } (q-p)^2 = (p-q)^2$$

$$= \sqrt{\frac{(p-q)^2}{p^2q^2}} \times \sqrt{p^2q^2 + 1}$$

$$= \left|\frac{p-q}{pq}\right| \times \sqrt{1 + p^2q^2}$$

$$\text{Area} = \frac{1}{2} \times \left|\frac{p+q}{\sqrt{1+p^2q^2}}\right| \times \left|\frac{p-q}{pq}\right| \times \sqrt{1+p^2q^2}$$

$$= \frac{|p^2 - q^2|}{2|pq|} \text{ units}^2$$

(b)(i) $e^2 = 1 + \frac{9}{16} \therefore e = \frac{5}{4}$

$$\text{Focus} = (\pm ae, 0) = (\pm 5, 0)$$

(b)(ii) differentiate wrt x

$$\frac{2x}{16} - \frac{2y}{9} \times \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{x}{8} \times \frac{9}{2y} = \frac{9x}{16y}$$

(b)(ii) continued at P $m_T = \frac{9x_1}{16y_1}$

equation of tangent :

$$y - y_1 = \frac{9x_1}{16y_1}(x - x_1)$$

$$16y_1(y - y_1) = 9x_1(x - x_1)$$

$$16yy_1 - 16(y_1)^2 = 9xx_1 - 9(x_1)^2$$

$$\frac{xx_1}{16} - \frac{yy_1}{9} = \frac{(x_1)^2}{16} - \frac{(y_1)^2}{9}$$

i.e. $\frac{xx_1}{16} - \frac{yy_1}{9} = 1$ as (x_1, y_1) lies on $\frac{x^2}{16} - \frac{y^2}{9} = 1$

(b)(iii) T is x -int. where $y = 0$

$$T = \left(\frac{16}{x_1}, 0\right)$$

$$\frac{S'T}{ST} = \frac{5 + \frac{16}{x_1}}{5 - \frac{16}{x_1}}$$

i.e. $\frac{S'T}{ST} = \frac{5x_1 + 16}{5x_1 - 16}$

$$\frac{S'P}{SP} = \frac{ePM'}{ePM} \text{ where } M \text{ is corr. directrix}$$

$$= \frac{PM'}{PM} = \frac{x_1 + \frac{16}{5}}{x_1 - \frac{16}{5}}$$

i.e. $\frac{S'P}{SP} = \frac{5x_1 + 16}{5x_1 - 16} = \frac{S'T}{ST}$

(b)(iv) Let $\angle S'PT = \beta, \angle SPT = \gamma$ and $\angle PTS = \alpha$

$$\angle PTS' = 180 - \alpha \text{ (st. } \angle)$$

$$\frac{\sin(180 - \alpha)}{S'P} = \frac{\sin \beta}{S'T}$$


i.e. $\sin \beta = \frac{S'T \sin(180 - \alpha)}{S'P}$

$$= \frac{S'T \sin \alpha}{S'P}$$

Similarly, $\sin \gamma = \frac{ST \sin \alpha}{SP}$

Since $\frac{S'T}{ST} = \frac{S'P}{SP}$ then $\frac{S'T}{S'P} = \frac{ST}{SP}$

Hence $\sin \beta = \sin \gamma$ i.e. $\beta = \gamma$



1) $w = g$
 Resolving Horizontally
 $T \sin \theta = m w$
 $T \sin \theta = m r g$ (1)

ii) Balancing Vertically
 $T \cos \theta = m g$ (2)

$\frac{T \sin \theta}{T \cos \theta} = \frac{m r g}{m g}$
 $\tan \theta = r$
 $\therefore \theta = \tan^{-1}(g r)$ #

iii) From (1) $T^2 \sin^2 \theta = m^2 r^2 g$ (3)
 from (2) $T^2 \cos^2 \theta = m^2 g$ (4)

$T^2 (\sin^2 \theta + \cos^2 \theta) = m^2 r^2 g + m^2 g$ (5)

$T^2 = m^2 g^2 (1 + r^2)$ (2)

$T = m g \sqrt{1 + r^2}$

b) i) $\cos \theta + x \sin \theta = (\cos \theta + x \sin \theta)^2 + [D_x \text{ mean TR}]$
 $RHS = \cos^2 \theta + 4x \cos^2 \theta \sin \theta - 6x^2 \theta \sin^2 \theta - 4x \cos \theta \sin^2 \theta + \sin^4 \theta$
 Equate $Rx \cos \theta = \cos^2 \theta - 6x^2 \theta \sin^2 \theta - 4x \cos \theta \sin^2 \theta$ (1)
 Equate $\sin \theta = 4 \cos^2 \theta \sin \theta - 4x \cos \theta \sin^2 \theta$ (2)

ii) $\text{tan } \theta = \frac{4 \cos^2 \theta \sin \theta - 4x \cos \theta \sin^2 \theta}{\cos^2 \theta - 6x^2 \theta \sin^2 \theta - 4x \cos \theta \sin^2 \theta + \sin^4 \theta}$
 Calculating top & bottom
 (1) $\frac{1 - 6x^2 + x^4}{4x^2 - 4x^2}$

76) iii)

$\tan \theta = 1$
 $\theta = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4}$

roots are $\tan \frac{\pi}{4}, \tan \frac{5\pi}{4}, \tan \frac{9\pi}{4}, \tan \frac{13\pi}{4}$

$\tan \frac{5\pi}{4} = -\tan \frac{\pi}{4}$
 $\tan \frac{9\pi}{4} = -\tan \frac{\pi}{4}$
 $\tan \frac{13\pi}{4} = -\tan \frac{\pi}{4}$

is $\tan \frac{\pi}{4}, \tan \frac{5\pi}{4}, \tan \frac{9\pi}{4}, \tan \frac{13\pi}{4}$, $-\tan \frac{\pi}{4}$ or equivalent

iv) Product of roots = 1
 $\therefore \tan \frac{\pi}{6} \times \tan \frac{5\pi}{6} \times (-\tan \frac{7\pi}{6}) \times (-\tan \frac{11\pi}{6}) = 1$

or $\tan \frac{\pi}{6} \tan \frac{5\pi}{6} \tan \frac{7\pi}{6} \tan \frac{11\pi}{6} = 1$
 (Note: Hence true, a must follow from iii)

v) $I = \int_{\pi}^{\pi} x \cos 2x \, dx$ put $\pi - x$ in place of x

$I = \int_{\pi}^{\pi} (\pi - x) \cos (2\pi - 2x) \, dx$

$I = \int_{\pi}^{\pi} \pi \cos (2\pi - 2x) \, dx - \int_{\pi}^{\pi} x \cos (2\pi - 2x) \, dx$

$I = \int_{\pi}^{\pi} \pi \cos (2x) \, dx - \int_{\pi}^{\pi} x \cos (2x) \, dx$

$= \int_{\pi}^{\pi} \cos 2x \, dx - I$

$2I = -\pi [\sin 2x]_{\pi}^{\pi}$

$\therefore I = 0$

Let $u = x$ $v = \cos 2x$
 $u = 1$ $v = \cos 2x$
 $I = [\frac{x}{2} \sin 2x]_{\pi}^{\pi} - \int_{\pi}^{\pi} \frac{1}{2} \sin 2x$
 $= 0 - [-\frac{1}{4} \cos 2x]_{\pi}^{\pi}$
 $= 0 + \frac{1}{4} [1 - 1]$
 $= 0$



Question Eight

$$a) 1) I_n = \int_0^{\frac{\pi}{2}} (\sin x)^n dx$$

$$= \int_0^{\frac{\pi}{2}} (n-1) (\sin x)^{n-2} \cos x dx$$

let $u = \sin^{n-1} x$ $u' = \sin x$
 $du = (n-1) (\sin x)^{n-2} \cos x dx$ $v = -\cos x$ ✓

$$I_n = - [\cos x \sin^{n-1} x]_0^{\frac{\pi}{2}} + (n-1) \int_0^{\frac{\pi}{2}} \sin^{n-2} x \cos^2 x dx$$

$$= 0 + (n-1) \int_0^{\frac{\pi}{2}} (\sin^{n-2} x) (1 - \sin^2 x) dx$$

$$= (n-1) \left[\int_0^{\frac{\pi}{2}} \sin^{n-2} x dx - \int_0^{\frac{\pi}{2}} \sin^n x dx \right]$$

$$= (n-1) [I_{n-2} - I_n]$$
 ✓

$$I_n = (n-1) I_{n-2} - (n-1) I_n \quad (3)$$

$$n I_n = (n-1) I_{n-2}$$

$$I_n = \frac{n-1}{n} I_{n-2}$$

ii) put $2n$ in place of n

$$I_{2n} = \frac{2n-1}{2n} I_{2n-2}$$

$$= \frac{2n-1}{2n} \times \frac{2n-3}{2n-2} \times I_{2n-4} \dots$$

$$= \frac{2n-1}{2n} \times \frac{2n-3}{2n-2} \times \frac{2n-5}{2n-4} \times \dots \times I_2 \times I_0 \quad (3)$$

and $I_0 = \int_0^{\frac{\pi}{2}} 1 dx = \frac{\pi}{2}$ ✓

$$I_2 = \int_0^{\frac{\pi}{2}} \sin^2 x dx$$

$$= \frac{1}{2} [1 - \cos 2x]_0^{\frac{\pi}{2}}$$

$$= \frac{1}{2} [(1-1) - (1-0)]$$

$$= \frac{1}{2}$$
 ✓

∴ $I_{2n} = \frac{2n-1}{2n} \times \frac{2n-3}{2n-2} \times \frac{2n-5}{2n-4} \times \dots \times \frac{1}{2} \times \frac{\pi}{2}$
 Now numerator & denominator increase by 2 from right to left

$$\therefore I_{2n} = \frac{2n-1}{2n} \times \frac{2n-3}{2n-2} \times \frac{2n-5}{2n-4} \times \dots \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2}$$

Q 8 b) $\cos(A+B) + \cos(A-B) = 2 \cos A \cos B$
 $\cos(4x+2x) + \cos(4x-2x) = 2 \cos 4x \cos 2x$

∴ eqn becomes

$$2 \cos 4x \cos 2x + 2 \cos 2x = 0 \quad \checkmark$$

$$2 \cos 2x (\cos 4x + 1) = 0$$

$$2 \cos 2x = 0, \quad \cos 4x + 1 = 0$$

$$\cos 2x = 0, \quad \cos 4x = -1$$

$$2x = \frac{\pi}{2}, \quad 4x = \pi$$


$$x = \frac{\pi}{4}, \quad x = \frac{\pi}{4} \quad \checkmark$$

$$0 \leq x \leq \frac{\pi}{2}$$

$$0 \leq 4x \leq 2\pi$$



(3)



$$\frac{2x}{\sin \alpha} = \frac{3x}{\sin(\alpha + 45^\circ)} \quad \checkmark$$

$$\frac{\sin(\alpha + 45^\circ)}{\sin \alpha} = \frac{3}{2}$$

$$3 \sin \alpha = 2 \sin(\alpha + 45^\circ)$$

$$3 \sin \alpha = 2 (\sin \alpha \cos 45^\circ + \cos \alpha \sin 45^\circ)$$

$$3 \sin \alpha = 2 \left(\frac{\sin \alpha}{\sqrt{2}} + \frac{\cos \alpha}{\sqrt{2}} \right)$$

$$3 \sin \alpha = \sqrt{2} \sin \alpha + \sqrt{2} \cos \alpha$$

$$\sin \alpha (3 - \sqrt{2}) = \sqrt{2} \cos \alpha \quad \checkmark$$

$$\frac{\sin \alpha}{\cos \alpha} = \frac{\sqrt{2}}{3 - \sqrt{2}} \times \frac{3 + \sqrt{2}}{3 + \sqrt{2}}$$

(4)

$$\tan \alpha = \frac{3\sqrt{2} + 2}{7}$$

$$\alpha = \tan^{-1} \left(\frac{2 + 3\sqrt{2}}{7} \right) \quad \checkmark$$

↑
 Mark for
 same work
 or diagram

8d) (1), (2, 3), (4, 5, 6), (7, 8, 9, 10)

$$T_1 \text{ in first baskets} = 1$$

$$T_1 \text{ in 2nd " } = 1+1$$

$$T_1 \text{ in 3rd " } = 1+1+2$$

$$T_1 \text{ in 4th " } = 1+1+2+3$$

$$T_1 \text{ in 5th " } = 1+1+2+3+4$$

$$\text{or } T_1 \text{ in } n\text{th " } = 1 + \underbrace{(1+2+3+\dots+n-1)}_{S_n = \frac{n}{2}(a+d)}$$

$$= 1 + \frac{n-1}{2} (1+n-1)$$

$$= 1 + \frac{n^2-n}{2}$$

$$\text{or } a = \frac{n^2-n+2}{2} \quad \checkmark$$

(2)

$$S_n = \frac{n}{2} (2a + (n-1)d)$$

$$\text{Sum of } n \text{ terms} = \frac{n}{2} \left(2 \left(\frac{n^2-n+2}{2} \right) + (n-1)1 \right) \checkmark$$

$$= \frac{n}{2} (n^2-n+2+n-1)$$

$$= \frac{n}{2} (n^2+1)$$