

## MATHEMATICS (EXTENSION 1)

2012 HSC Course Assessment Task 3 (Trial Examination)

June 27, 2012

### General instructions

- Working time – 2 hours.  
(plus 5 minutes reading time)
- Write using blue or black pen. Where diagrams are to be sketched, these may be done in pencil.
- Board approved calculators may be used.
- Attempt all questions.
- At the conclusion of the examination, bundle the booklets + answer sheet used in the correct order within this paper and hand to examination supervisors.

### SECTION I

- Mark your answers on the answer sheet provided (numbered as page 9)

### SECTION II

- Commence each new question on a new page. Write on both sides of the paper.
- All necessary working should be shown in every question. Marks may be deducted for illegible or incomplete working.

STUDENT NUMBER: ..... # BOOKLETS USED: .....

Class (please ✓)

- |   |   |
|---|---|
| <input type="radio"/> 12M4A – Mr Weiss<br><input type="radio"/> 12M4B – Mr Ireland<br><input type="radio"/> 12M4C – Mr Fletcher | <input type="radio"/> 12M3C – Ms Ziaziaris<br><input type="radio"/> 12M3D – Mr Lowe<br><input type="radio"/> 12M3E – Mr Lam |
|---|---|

Marker's use only.

QUESTION	1-10	11	12	13	14	Total	%
MARKS	10	15	15	15	15	70	

### STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE:  $\ln x = \log_e x, \quad x > 0$





## Section I: Objective response

Mark your answers on the multiple choice sheet provided.

Marks

1. Which is the correct value of  $\lim_{x \rightarrow 0} \frac{3x}{\sin 2x}$ ? 1  
 (A) 0 (B)  $\frac{2}{3}$  (C)  $\frac{3}{2}$  (D) 3
2. Which of the following is the acute angle (correct to the nearest degree) between the two lines  $2x - y + 1 = 0$  and  $3x + y - 4 = 0$ ? 1  
 (A)  $11^\circ$  (B)  $45^\circ$  (C)  $79^\circ$  (D)  $135^\circ$
3. Which of the following expressions will result in the coordinates of the point  $P$  which divides the interval  $AB$  externally in the ratio  $3 : 2$ , given  $A$  is  $(-5, 2)$  and  $B$  is  $(4, 5)$ ? 1  
 (A)  $\left( \frac{(2)(-5) + (-3)(4)}{-3+2}, \frac{(2)(2) + (-3)(5)}{-3+2} \right)$   
 (B)  $\left( \frac{(-3)(-5) + (2)(4)}{-3+2}, \frac{(-3)(2) + (2)(5)}{-3+2} \right)$   
 (C)  $\left( \frac{(-2)(-5) + (-3)(4)}{-3+2}, \frac{(-2)(2) + (-3)(5)}{-3+2} \right)$   
 (D)  $\left( \frac{(2)(2) + (-3)(5)}{-3+2}, \frac{(2)(-5) + (-3)(4)}{-3+2} \right)$
4. Which of the following is the derivative of  $xe^{2x}$ ? 1  
 (A)  $e^{2x}(1+x)$  (B)  $e^{2x}(1+2x)$  (C)  $2x^2e^{2x}$  (D)  $\frac{1}{2}xe^{2x}$
5. Which of the following represents the complete solutions for  $-180^\circ < \theta \leq 180^\circ$  to the equation  $\cos^2 \frac{\theta}{2} = \frac{1}{4}$ ? 1  
 (A)  $60^\circ, 120^\circ$  (B)  $120^\circ, 240^\circ$  (C)  $\pm 60^\circ, \pm 120^\circ$  (D)  $\pm 120^\circ$
6. What should  $\int \cos^2 \frac{1}{2}x dx$  be transformed into, in order to find its primitive? 1  
 (A)  $\int \frac{1}{2} - \frac{\cos x}{2} dx$  (C)  $\int \frac{1}{2} - \frac{\cos 2x}{2} dx$   
 (B)  $\int \frac{1}{2} + \frac{\cos 2x}{2} dx$  (D)  $\int \frac{1}{2} + \frac{\cos x}{2} dx$

7. It is known that  $\log_e x + \sin x = 0$  has a root close to  $x = 0.5$ . Using one application of Newton's method, which of the following gives a better approximation to 2 decimal places? 1  
 (A) 0.43 (B) 0.73 (C) 0.57 (D) 0.27
8. Which of the following graphs represents  $y = \cos^{-1} 2x$ ? 1



9. If  $\sqrt{3} \cos x - \sin x \equiv R \cos(x + \alpha)$ , which of the following gives the correct value of  $\alpha$ ? 1  
 (A)  $\frac{\pi}{6}$  (B)  $\frac{5\pi}{6}$  (C)  $\frac{7\pi}{6}$  (D)  $\frac{11\pi}{6}$
10. Zac and Mitchell play a series of games. The series ends when one player has won two games. In any game the probability that Zac wins is  $\frac{3}{5}$  and the probability that Mitchell wins is  $\frac{2}{5}$ . 1

What is the probability that three games are played?

- (A)  $\frac{6}{25}$  (B)  $\frac{19}{25}$  (C)  $\frac{12}{25}$  (D)  $\frac{18}{25}$

End of Section I.  
 Examination continues overleaf.

## Section II: Short answer

Question 11 (15 Marks) Commence a NEW page. Marks

- (a) Solve for  $x$ :  $\frac{1}{x} > x$  3
- (b) Evaluate  $\int_0^2 \frac{dx}{\sqrt{16-x^2}}$  2
- (c) Find the exact value of  $\sin\left(2 \tan^{-1} \frac{3}{7}\right)$ , showing full working. 3
- (d) Solve  $\sin 2\theta = \sin \theta$ ,  $0 \leq \theta \leq 2\pi$ . 3
- (e) Using the substitution  $u = \tan x$ , find the exact value of 4

$$\int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{3 + \tan^2 x} dx$$

Question 12 (15 Marks) Commence a NEW page. Marks


- (a) If  $P(x) = x^3 - 6x^2 + ax - 4$ ,  $a > 0$ , 3
- Given all the roots of  $P(x) = 0$  are real and positive, and that one of the roots is the product of the other two roots, show that  $a = 10$ .
  - Show that  $x - 2$  is a factor of  $P(x) = x^3 - 6x^2 + 10x - 4$ . 2
- (b) Air is being pumped into a spherical balloon at a rate of  $20 \text{ cm}^3 \text{ s}^{-1}$ . Find the rate of increase of the surface area of the balloon when the radius is 5 cm. 3
- (c) A particle moves along the  $x$  axis such that its velocity  $v \text{ ms}^{-1}$  is given by 3

$$v^2 = -4x^2 + 8x + 32$$

- By expressing the acceleration as a function in terms of  $x$ , prove that the particle is undergoing simple harmonic motion. 3
- Find the amplitude. 2
- Find the maximum acceleration. 2


Question 13 (15 Marks) Commence a NEW page. Marks

- (a) Prove by mathematical induction that  $5^n + 2 \times 11^n$  is divisible by 3, where  $n$  is a positive integer. 3
- (b) Show that the shaded area is  $A = \frac{1}{2} \ln 2$  units<sup>2</sup>. 3



- (c) Two circles intersect at  $A$  and  $B$ .  $CAE$  is a straight line where  $C$  is a point on the first circle and  $E$  is a point on the second circle. The tangents to the circles at  $C$  and  $E$  meet at  $D$ . 4

Copy the diagram into your writing booklet.

Prove that  $BCDE$  is a cyclic quadrilateral, without adding any construction lines.

- (d) The acceleration of a raindrop which at time  $t$  seconds is falling with speed  $v$  metres per second is given by the equation

$$\frac{dv}{dt} = -\frac{1}{3}(v - 3g)$$

where  $g$  is a constant.


- Show that  $v = 3g + Ae^{-\frac{1}{3}t}$ , where  $A$  is a constant, satisfies the above equation. 1
- Given that the initial velocity has a value of  $g$ , find the value of  $A$ . 1
- After how many seconds is the raindrop falling with a speed of  $2g$  metres per second? Give your answer correct to 1 decimal place. 2
- What value does  $v$  approach as  $t \rightarrow \infty$ ? 1

## Question 14 (15 Marks)

Commence a NEW page.

Marks

- (a)  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  are two points on the parabola  $x^2 = 4ay$ . The tangent at  $P$  and the line through  $Q$  parallel to the axis of the parabola meet at the point  $R$ .




The tangent at  $Q$  and the line through  $P$  parallel to the axis of the parabola meet at the point  $S$ .

- Show that the equations of the tangents at  $P$  and  $Q$  are  $y = px - ap^2$  and  $y = qx - aq^2$  respectively. 2
- Show that the coordinates of  $S$  and  $R$  are 2  
 $S(2ap, 2apq - aq^2)$       $R(2aq, 2apq - ap^2)$
- Show that  $PQRS$  is a parallelogram. 2
- Show that the area of this parallelogram is  $2a^2 |p - q|^3$ . 2

Question 14 continues overleaf...

Question 14 continued from the previous page...

- (b) A projectile is thrown horizontally from the top of a 125 m tower with velocity  $V$  metres per second. It clears a second tower of height 100 m by a distance of  $c$  metres, as shown. The two towers are  $20\sqrt{5}$  metres apart.



- i. The equations of motion for this system are

$$\begin{cases} x = Vt \\ y = -5t^2 + 125 \end{cases}$$

(Do not prove this)

Where is the origin of the system being taken from?

- Show that  $V = \frac{100}{\sqrt{25 - c}}$ .
- Prove that the minimum initial speed of the projectile to just clear the 100 m tower is  $20 \text{ ms}^{-1}$ .
- Hence, find how far past the 100 m tower will the projectile strike the ground.
- Determine the vertical component of the velocity of the projectile when it strikes the ground.

End of paper.

## Suggested Solutions

## Section I

(Lowe)


1. (C) 2. (B) 3. (A) 4. (B) 5. (D)  
6. (D) 7. (C) 8. (D) 9. (A) 10. (C)

## Question 11 (Lam)

(a) (3 marks)

- ✓ [1] for multiplying by square of denominator.
- ✓ [0] for entire part if only multiplying by denominator.
- ✓ [1] for each correct inequality.

$$\begin{aligned} \frac{1}{x} &> \frac{x}{x^2} \\ x &> x^3 \\ x^3 - x &< 0 \\ x(x^2 - 1) &< 0 \\ x(x-1)(x+1) &< 0 \end{aligned}$$



From the sketch,

$$x < -1 \text{ or } 0 < x < 1$$


(b) (2 marks)

- ✓ [1] for correct primitive.
- ✓ [1] for correct evaluation of limits.

$$\begin{aligned} \int_0^2 \frac{dx}{\sqrt{16-x^2}} &= \left[ \sin^{-1} \frac{x}{4} \right]_0^2 \\ &= \sin^{-1} \frac{1}{2} - \sin^{-1} 0 \\ &= \frac{\pi}{6} \end{aligned}$$

(c) (3 marks)

- ✓ [1] for drawing relevant right-angled triangle.
- ✓ [1] for expanding  $\sin 2\alpha$ .
- ✓ [1] for final answer.

Let  $\alpha = \tan^{-1} \frac{3}{7}$ . Then  $\tan \alpha = \frac{3}{7}$ :

$$\begin{aligned} \sin \left( 2 \tan^{-1} \frac{3}{7} \right) &\equiv \sin 2\alpha \\ &= 2 \sin \alpha \cos \alpha \\ &= 2 \times \frac{3}{\sqrt{58}} \times \frac{7}{\sqrt{58}} \\ &= \frac{42}{58} = \frac{21}{29} \end{aligned}$$

(d) (3 marks)

- ✓ [1] for factorising expression into  $\sin \theta(2 \cos \theta - 1) = 0$ .
- ✓ [1] for solutions in positive integral multiples of  $\pi$ .
- ✓ [1] for solutions in multiples of  $\frac{\pi}{3}$ .

$$\begin{aligned} \sin 2\theta &= \sin \theta \\ 2 \sin \theta \cos \theta - \sin \theta &= 0 \\ \sin \theta(2 \cos \theta - 1) &= 0 \\ \sin \theta = 0 &\quad \cos \theta = \frac{1}{2} \\ \theta = 0, \pi, 2\pi &\quad \theta = \frac{\pi}{3}, \frac{5\pi}{3} \\ \therefore \theta &= 0, \pi, 2\pi, \frac{\pi}{3}, \frac{5\pi}{3} \end{aligned}$$

(e) (4 marks)

- ✓ [1] for changing limits.
- ✓ [1] for making algebraic substitution.
- ✓ [1] for correct primitive.
- ✓ [1] for final answer.

$$\int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{3 + \tan^2 x} dx$$

Letting  $u = \tan x$ ,

$$\begin{aligned} \frac{du}{dx} &= \sec^2 x \\ \therefore du &= \sec^2 x dx \\ x = 0 &\Rightarrow u = 0 \\ x = \frac{\pi}{4} &\Rightarrow u = 1 \end{aligned}$$

$$\begin{aligned} \int_0^1 \frac{\sec^2 x}{3 + \tan^2 x} dx &= \int_{u=0}^1 \frac{\sec^2 x dx}{3 + u^2} \\ &= \int_0^1 \frac{du}{3 + u^2} \\ &= \frac{1}{\sqrt{3}} \left[ \tan^{-1} \frac{u}{\sqrt{3}} \right]_0^1 \\ &= \frac{1}{\sqrt{3}} \left( \tan^{-1} \frac{1}{\sqrt{3}} - \tan^{-1} 0 \right) \\ &= \frac{1}{\sqrt{3}} \times \frac{\pi}{6} \\ &= \frac{\pi}{6\sqrt{3}} \left( = \frac{\pi\sqrt{3}}{18} \right) \end{aligned}$$

## Question 12 (Lowe)

(a) i. (3 marks)

- ✓ [1] for  $\alpha\beta = 2$ .
- ✓ [1] for  $\alpha + \beta = 4$ .
- ✓ [1] for final answer.

$P(x) = x^3 - 6x^2 + ax - 4$ . Let the roots be  $\alpha$ ,  $\beta$  and  $\alpha\beta$ .

- Sum of roots:

$$\alpha + \beta + \alpha\beta = -\frac{b}{a} = 6 \quad (12.1)$$

- Pairs of roots:

$$\alpha\beta + \alpha^2\beta + \beta^2\alpha = \frac{c}{a} = a \quad (12.2)$$

- Product of roots:

$$\begin{aligned} \alpha\beta(\alpha\beta) &= -\frac{d}{a} = 4 \\ \alpha^2\beta^2 &= 4 \\ \therefore \alpha\beta &= 2 \quad (12.3) \end{aligned}$$

as roots are positive.

Substitute (12.3) into (12.1):

$$\begin{aligned} \alpha + \beta + 2 &= 6 \\ \therefore \alpha + \beta &= 4 \quad (12.4) \end{aligned}$$

Substitute (12.4) into (12.2) to find  $a$ :

$$\begin{aligned} \alpha\beta^2 + \alpha\beta(\alpha + \beta) &= a \quad (12.5) \\ 2 + 2(4) &= a \\ \therefore a &= 10 \end{aligned}$$

ii. (2 marks)

- ✓ [2] for correct application of factor theorem.

If  $x - 2$  is a factor then  $P(2) = 0$ .

$$\begin{aligned} P(2) &= 2^3 - 6(2^2) + 10(2) - 4 \\ &= 8 - 24 + 20 - 4 = 0 \end{aligned}$$

(b) (3 marks)

- ✓ [1] for  $\frac{dV}{dt}$ .
- ✓ [1] for  $\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$ .
- ✓ [1] for final answer.

$$\frac{dV}{dt} = 20 = \frac{dV}{dr} \times \frac{dr}{dt}$$

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dV}{dr} = \frac{4}{3} \times \pi \times 3r^2 = 4\pi r^2$$

$$\therefore \frac{dV}{dt} = 20 = 4\pi r^2 \Big|_{r=5} \times \frac{dr}{dt}$$

$$= 4\pi \times 25 \times \frac{dr}{dt}$$

$$\therefore \frac{dr}{dt} = \frac{1}{5\pi}$$

$$\text{Now } \frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$$

$$SA = 4\pi r^2$$

$$\therefore \frac{dA}{dr} = 8\pi r$$

$$\frac{dA}{dt} = 8\pi r \Big|_{r=5} \times \frac{1}{5\pi}$$

$$= 8 \text{ cm}^2 \text{ s}^{-1}$$

(c) i. (3 marks)

- ✓ [1] for using  $\frac{d}{dx}(\frac{1}{2}v^2)$  to find acceleration.
- ✓ [1] obtaining  $\ddot{x} = -4x + 4$ .
- ✓ [1] factorising and noting form  $\frac{dv}{dt} = -n^2(x - x_0)$  for SHM.

$$v^2 = -4x^2 + 8x + 32$$

$$\frac{1}{2}v^2 = -2x^2 + 4x + 16$$

$$\ddot{x} = \frac{dv}{dt} = \frac{d}{dx} \left( \frac{1}{2}v^2 \right)$$

$$= \frac{d}{dx} (-2x^2 + 4x + 16)$$

$$= -4x + 4 = -4(x - 1)$$

As acceleration is proportion to the opposite direction of displacement, hence the particle is moving in simple harmonic motion with centre at  $x = 1$ .

ii. (2 marks)

- ✓ [1] for  $x = 4, x = -2$ .
- ✓ [1] for finding amplitude.

The amplitude occurs when  $\dot{x} = 0$ .

$$-4x^2 + 8x + 32 = 0$$

$$x^2 - 2x - 8 = 0$$

$$(x - 4)(x + 2) = 0$$

$$\therefore x = 4, -2$$

$$a = \frac{4 - (-2)}{2} = 3$$

As the centre of motion is  $x = 1$  and particle's maximum displacement is 4 or  $-2$ , therefore the amplitude is  $a = 3$ .

iii. (1 mark)

Maximum acceleration occurs at the amplitude, i.e.  $x = 4$  or  $x = -2$ .

$$\ddot{x} = -4(x - 1) \Big|_{x=-2}$$

$$= -4(-2 - 1) = 12$$

Question 13 (Ziaziaris)

(a) (3 marks)

- ✓ [1] for proving base case.
- ✓ [1] for inductive step.
- ✓ [1] for required proof.

Let  $P(n)$  be the statement  $5^n + 2 \times 11^n$  is divisible by 3, i.e.

$$5^n + 2 \times 11^n = 3J$$

where  $J \in \mathbb{N}$ .

- Base case:  $P(1)$ :

$$5^1 + 2 \times 11 = 5 + 22 = 27$$

which is divisible by 3. Hence  $P(1)$  is true.

- Inductive step:

– Assume  $P(k)$  is true for some  $k \in \mathbb{N}, k < n$ , i.e.

$$5^k + 2 \times 11^k = 3M$$

where  $M \in \mathbb{N}$ . Alternatively,

$$5^k = 3M - 2 \times 11^k$$

- Examine  $P(k+1)$ :

$$5^{k+1} + 2 \times 11^{k+1}$$

$$= 5^k 5^1 + 2 \times 11^k 11^1$$

$$= 5(3M - 2 \times 11^k) + 2 \times 11^k 11^1$$

$$= 3 \times 5M - 10 \times 11^k + 2 \times 11 \times 11^k$$

$$= 3 \times 5M - 10 \times 11^k + 22 \times 11^k$$

$$= 3 \times 5M + 12 \times 11^k$$

$$= 3 \underbrace{(5M + 4 \times 11^k)}_{\in \mathbb{N}} \equiv 3P$$

where  $P \in \mathbb{N}$ . Hence  $P(k+1)$  is true.

Since  $k \in \mathbb{N}$  and truth in  $P(k)$  also leads to truth in  $P(k+1)$ , therefore  $P(n)$  is true by induction.

(b) (3 marks)

- ✓ [1] for converting integrand to  $\frac{\sin y}{\cos y}$
- ✓ [1] for correct primitive
- ✓ [1] for final answer.

$$A = \int_0^{\frac{\pi}{4}} \tan y \, dy$$


$$= \int_0^{\frac{\pi}{4}} \frac{\sin y}{\cos y} \, dy$$

$$= \left[ -\log_e (\cos y) \right]_0^{\frac{\pi}{4}}$$

$$= -\log_e \cos \frac{\pi}{4} + \log_e \cos 0$$

$$= -\log_e \frac{1}{\sqrt{2}} + \log_e 1$$

$$= -\log_e 2^{-\frac{1}{2}} = \frac{1}{2} \log_e 2$$

(c) (4 marks) – marking scheme embedded inline. Presence of  $\checkmark$  indicates 1 mark.

- ✓ Let  $\angle DCE = \alpha$  and  $\angle DEC = \beta$ .
- ✓  $\therefore \angle CBA = \alpha$  ( $\angle$  in alternate segment)
- ✓ Similarly,  $\angle ABE = \beta$  ( $\angle$  in alternate segment)
- ✓ Also,  $\angle CDE = 180^\circ - (\alpha + \beta)$ . (Angle sum of  $\triangle CDE$ )
- Hence  $\angle CDE + \angle CBE = 180^\circ$
- ✓ Opposite  $\angle$  in  $BCDE$  are supplementary. Hence  $BCDE$  is a cyclic quadrilateral.

(d) i. (1 mark)

$$v = 3g + Ae^{-\frac{1}{3}t}$$

$$-3g = 3g + Ae^{-\frac{1}{3}t}$$

$$v - 3g = Ae^{-\frac{1}{3}t}$$

$$\frac{dv}{dt} = -\frac{1}{3} \underbrace{Ae^{-\frac{1}{3}t}}_{=(v-3g)}$$

$$= -\frac{1}{3}(v - 3g)$$

ii. (1 mark)

$$t = 0, v = g$$

$$\therefore g = 3g + Ae^0$$

$$\therefore A = -2g$$

iii. (2 marks)

$$v = 2g, t = ?$$

$$2g = 3g - 2ge^{-\frac{1}{3}t}$$

$$-g = -2ge^{-\frac{1}{3}t}$$

$$\frac{1}{2} = e^{-\frac{1}{3}t}$$

$$-\frac{1}{3}t = \log_e \frac{1}{2} = -\log_e 2$$

$$\therefore t = 3 \log_e 2 \approx 2.1 \text{ seconds}$$

iv. (1 mark)

As  $t \rightarrow \infty, v \rightarrow 3g$ .

## Question 14 (Ireland/Fletcher)

(a) i. (2 marks)

- ✓ [1] for proving  $\frac{dy}{dx} = p$  at  $P$ .  
 ✓ [1] for equation of tangent at  $P$ .

$$x^2 = 4ay \Rightarrow y = \frac{x^2}{4a}$$

$$\frac{dy}{dx} = \frac{2x}{4a} = \frac{x}{2a}$$

At  $x = 2ap$ ,

$$\frac{dy}{dx} = \frac{2ap}{2a} = p$$

Equation of the tangent at  $P$ :

$$y - ap^2 = p(x - 2ap) = px - 2ap^2$$

$$y = px - ap^2$$

Similarly, the tangent at  $Q$  is

$$y = qx - aq^2$$

ii. (2 marks)

- ✓ [1] for each  $y$  coordinate.  
 Coordinates of  $S$  arise from the intersection of  $x = 2ap$  and  $y = qx - aq^2$ :

$$y = q(2ap) - aq^2 = 2apq - aq^2$$

$$\therefore S(2ap, 2apq - aq^2)$$

Coordinates of  $R$  arise from the intersection of  $x = 2aq$  and  $y = px - ap^2$ :

$$y = p(2aq) - ap^2 = 2apq - ap^2$$

$$\therefore R(2aq, 2apq - ap^2)$$

iii. (2 marks)

- ✓ [1] for showing  $PS \parallel QR$ .  
 ✓ [1] for showing  $PS = QR$ .

$$d_{PS} = \sqrt{(2ap - 2ap)^2 + (ap^2 - (2apq - aq^2))^2}$$

$$= \sqrt{(a(p - q)^2)^2}$$

$$= a(p - q)$$

$$d_{QR} = \sqrt{(2aq - 2aq)^2 + (aq^2 - (2apq - ap^2))^2}$$

$$= \sqrt{a(q - p)^2} = \sqrt{a(p - q)^2}$$

$$= a(p - q)$$

As  $PS = QR$ , hence one pair of opposite sides equal and parallel. Hence  $PQRS$  is a parallelogram.

Alternatively, if  $PQRS$  is a parallelogram, then the diagonals bisect each other; i.e.  $QS$  and  $PR$  share the same midpoint. Show via midpoint formula results in

$$MP_{QS} = \left( \frac{a(p+q)}{2}, apq \right)$$

$$MP_{PR} = \left( \frac{a(p+q)}{2}, apq \right)$$

iv. (2 marks)

- ✓ [1] for  $h$  (fully)  
 ✓ [1] for area.  
 • Use  $A = bh$ .  
 •  $h$  is perpendicular distance from  $Q$  to  $PS$ .  
 • Use  $b = d_{PS}$ .

Using the perpendicular dist formula with  $x = 2ap$  &  $Q(2aq, aq^2)$ :

$$h = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

$$= \frac{|2aq(1) + 0 - 2ap|}{\sqrt{1^2 + 0}}$$

$$= \frac{|2aq - 2ap|}{1} = 2a|q - p|$$

$$= 2a|p - q|$$

As  $|p - q| = |q - p|$ .

$$A = bh$$

$$= 2a|p - q| \times a(p - q)^2$$

$$= 2a^2|p - q|^3$$

(b) i. (1 mark)

Origin is at the base of tower.

ii. (2 marks)

- ✓ [1] for  $100 + c = -5 \left( \frac{400 \times 5}{V^2} \right) + 125$ .  
 ✓ [1] for final result shown.

When  $x = 20\sqrt{5}$ ,  $y = 100 + c$ . Using  $x = Vt$ ,

$$20\sqrt{5} = Vt$$

$$\therefore t = \frac{20\sqrt{5}}{V}$$

Substitute into  $y = -5t^2 + 125$ ,

$$100 + c = -5 \left( \frac{20\sqrt{5}}{V} \right)^2 + 125$$

$$= -5 \left( \frac{400 \times 5}{V^2} \right) + 125$$

$$-25 + c = -5 \times \frac{400 \times 5}{V^2}$$

$$25 - c = \frac{25 \times 400}{V^2}$$

$$V^2 = \frac{10000}{25 - c}$$

$$\therefore V = \frac{100}{\sqrt{25 - c}} \quad (V > 0)$$

iii. (1 mark) Projectile just clears tower when  $c = 0$ .

$$V = \frac{100}{\sqrt{25 - c}} \Big|_{c=0} = 20 \text{ ms}^{-1}$$

iv. (2 marks)

- ✓ [1] for  $x = 100$ .  
 ✓ [1] for final answer.

Projectile strikes ground when  $y = 0$ .

$$-5t^2 + 125 = 0$$

$$5t^2 = 125$$

$$\therefore t^2 = 25 \Rightarrow t = 5$$

When  $t = 5$ ,

$$x = Vt = 20 \times 5 = 100$$

Hence projectile will strike ground  $100 - 20\sqrt{5}$  metres past second tower.

v. (1 mark)

$$y = -5t^2 + 125$$

$$\dot{y} = -10t \Big|_{t=5} = -50 \text{ ms}^{-1}$$