



**2012**  
TRIAL HSC EXAMINATION

# Mathematics Extension 2

**General Instructions**

- Reading Time – 5 minutes
- Working Time – 3 hours
- Write using black or blue pen
- Board approved calculators may be used
- A table of standard integrals is provided at the back of this page
- Show all necessary working in questions 11 – 16

**Total Marks – 100**

**Section I 10 Marks**

- Attempt Questions 1 – 10
- Allow about 15 minutes for this section.

**Section II 90 marks**

- Attempt Questions 11 – 16

NAME: \_\_\_\_\_

TEACHER: \_\_\_\_\_

NUMBER: \_\_\_\_\_

QUESTION	MARK
1 – 10	/10
11	/15
12	/15
13	/15
14	/15
15	/15
16	/15
TOTAL	/100

**STANDARD INTEGRALS**

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE:  $\ln x = \log_e x, \quad x > 0$

Section I  
Total marks – 10  
Attempt Questions 1 – 10

Objective Response Questions

Answer each question on the multiple choice answer sheet provided.

1.  $\frac{2-i}{-2-i} = ?$

(A)  $-\frac{3}{5} + \frac{4}{5}i$


(B)  $-1$

(C)  $-1 + \frac{4}{3}i$


(D)  $-\frac{5}{3}$

2. Which of the following is the graph of  $9x^2 - 16y^2 = 144$ ?


(A)




(B)




(C)



(D)



3.



Which of the following is NOT a valid algebraic description of this locus?


(A)  $\text{Re } z = 2$

(B)  $|z| = |z-4|$

(C)  $\arg(z-4) + \arg z = \pi$


(D)  $z + \bar{z} = 4$

4. The solid shown in the diagram has a pair of parallel faces, one a regular octagon, and one a square, with vertices of each end joined by straight lines.




Which of the following diagrams shows a typical cross-section taken parallel to the two end faces?


(A)




(B)



(C)



(D)



5. The polynomial equation  $P(x)=0$  has roots  $\alpha$ ,  $\beta$  and  $\gamma$ .  
What are the roots of the polynomial equation  $P(3x+2)=0$ ?

- (A)  $\frac{\alpha}{3}-2, \frac{\beta}{3}-2, \frac{\gamma}{3}-2$       (B)  $\frac{\alpha-2}{3}, \frac{\beta-2}{3}, \frac{\gamma-2}{3}$ ,  
(C)  $3\alpha+2, 3\beta+2, 3\gamma+2$       (D)  $\alpha+\frac{2}{3}, \beta+\frac{2}{3}, \gamma+\frac{2}{3}$

6. Consider a polynomial  $P(x)$  of degree 3.

You are given 2 numbers  $a$  and  $b$  such that

- $a < b$
- $P(a) > P(b) > 0$
- $P'(a) = P'(b) = 0$

The polynomial has

- (A) 3 real zeros      (B) 1 real zero  $\gamma$  such that  $\gamma < a$   
(C) 1 real zero  $\gamma$  such that  $a < \gamma < b$       (D) 1 real zero  $\alpha$  such that  $\gamma > b$

7. Consider the following two statements:

I:  $\int_0^1 \frac{dx}{1+x^n} < \int_0^1 \frac{dx}{1+x^{n+1}}$

II:  $\int_0^{\frac{\pi}{2}} \sqrt{\sin x} dx = \int_0^{\frac{\pi}{2}} \sqrt{\cos x} dx$

Which of these statements are correct?


- (A) Neither statement      (B) Statement I only  
(C) Statement II only      (D) Both statements

8. A particle is moving along the  $x$ -axis, initially moving to the left from the origin. Its velocity and acceleration are given by  $v^2 = 2 \ln(3 + \cos x)$  and  $\ddot{x} = \frac{-\sin x}{3 + \cos x}$ .

Which of the following describe its subsequent motion?

- (A) Heads only to the left, alternately speeding up and slowing down, without becoming stationary.  
(B) Heads only to the left, alternately slowing to a stop then speeding up.  
(C) Slows to a stop, then heads to the right forever.  
(D) Oscillates between two points.

9. The graph shows a part of the hyperbola  $x = ct, y = \frac{c}{t}$ .




Which pair of parametric equations precisely describe the sections of the hyperbola shown?


- (A)  $x = c(t^2 + 1), y = \frac{c}{t^2 + 1}$       (B)  $x = c(1 - t^2), y = \frac{c}{1 - t^2}$   
(C)  $x = c\sqrt{1 - t^2}, y = \frac{c}{\sqrt{1 - t^2}}$       (D)  $x = c \sin t, y = \frac{c}{\sin t}$

10. After differentiating a relation implicitly, we find that  $\frac{dy}{dx} = \frac{y}{x}$ .  
Which of the following could be a graph of this relation?


(A)




(B)



(C)



(D)



Section II  
Total marks – 90  
Attempt Questions 11 – 16

Free Response Questions

Answer each question on the multiple choice answer sheet provided.

Question 11 (15 marks)

(a) Find the exact value of  $\int_0^1 x e^{-x^2} dx$ . 2

(b) Find  $\int \frac{dx}{x^2 + 6x + 10}$ . 1

(c) Evaluate  $\int_0^1 \sin^{-1} x dx$ . 3

(d) (i) Show that  $\int_0^1 \frac{5 - 5x^2}{(1 + 2x)(1 + x^2)} dx = \frac{1}{2} \left( \pi + \ln \frac{27}{16} \right)$ . 3


(ii) Hence evaluate  $\int_0^{\frac{\pi}{4}} \frac{\cos 2x dx}{1 + 2 \sin 2x + \cos 2x}$  using the substitution  $t = \tan x$ . 3

(e) (i) Show that  $\int_{-a}^a f(x) dx = \int_{-a}^a f(-x) dx$ . 1

(ii) Hence, or otherwise, evaluate  $\int_{-4}^4 (e^x - e^{-x}) \cos x dx$ . 2

**Question 12** (15 marks) Start a new booklet

- (a) Given that  $z = 6i - 8$ , find the square roots of  $z$  in the form  $a + ib$ . 3
- (b) (i) Write  $2 + 2\sqrt{3}i$  in modulus-argument form. 1
- Hence:
- (ii) Express  $(2 + 2\sqrt{3}i)^3$  in the form  $x + iy$ . 1
- (iii) Find all unique solutions to the equation  $z^4 = 2 + 2\sqrt{3}i$ , giving answers in modulus-argument form. 2
- (c) Given  $z$  is a complex number, sketch on a number plane the locus of a point  $P$  representing  $z$  such that  $\arg z = \arg[z - (1 + i)]$ . 2
- (d) In the diagram, a semi-circle has diameter  $OB$  and centre  $A$ , with  $OA = r$ .  $P$  is a point on the semicircle, and the vector  $OP$  represents the complex number  $a \operatorname{cis} \theta$ .




Write in simplest modulus-argument form the complex number represented by the vector

- (i)  $AP$  1
- (ii)  $BP$  2
- (e) In a bank of 12 switches, each switch can be set to one of three positions.
- (i) Write down the total number of ways all the switches in the bank can be arranged. 1
- (ii) Find the probability that if all the switches are set randomly, there will be equal numbers in each position. 2

**Question 13** (15 marks) Start a new booklet

- (a) Drawn below is the graph of  $y = \frac{2x}{1+x^2}$ .




- (i) Find the coordinates of the turning points  $A$  and  $A'$ . (There is no need to test their nature.) 2
- (ii) On separate diagrams draw graphs of the following functions:
1.  $y = \frac{|2x|}{1+x^2}$  1
  2.  $y = \frac{1+x^2}{2x}$  2
  3.  $y^2 = \frac{2x}{1+x^2}$  2
  4.  $y = \ln\left(\frac{2x}{1+x^2}\right)$  2
- (b) (i) The polynomial equation  $P(x) = 0$  has a double root  $x = \alpha$ . Show that  $x = \alpha$  is also a root of the equation  $P'(x) = 0$ . 2
- (ii) You are given that  $y = mx$  is a tangent to the curve  $y = 3 - \frac{1}{x^2}$ . Show that the equation  $mx^3 - 3x^2 + 1 = 0$  has a double root. 1
- (iii) Hence find the equations of any such tangents. 3

Question 14 (15 marks) Start a new booklet

(a) Use the method of cylindrical shells to find the volume of the solid formed when the region bounded by the curve  $y = \frac{1}{2x+1}$ , the  $x$ -axis, the  $y$ -axis and the line  $x=1$  is rotated about the line  $x=1$ .

4

(b) The diagram shows a part of the graph of a function of the form  $y = b \sin nx$ .



(i) Express  $n$  in terms of  $b$ .


1

(ii) Show that the area bounded by the curve and the  $x$ -axis is  $\frac{2b^2}{\pi}$  units<sup>2</sup>.

2

(iii) The base of a solid is the region bounded by the parabola  $x^2 = 4ay$  and its latus rectum.

2




Each slice of width  $\delta y$  taken perpendicular to both the base and the axis of symmetry is half of a sine curve, whose amplitude is equal to its base length.

Find the volume of this solid in terms of  $a$ .

Question 14 continues on the following page

(c) In the diagram,  $A, B$  and  $C$  are three points on a circle.  $P$  is another point on the circle, lying on the minor arc  $BC$ . Points  $L, M$  and  $N$  are the feet of the perpendiculars from  $P$  to the sides  $BC, CA$  and  $AB$  respectively.



(i) Explain why  $P, L, N$  and  $B$  are concyclic.

1

(ii) Explain why  $P, L, C$  and  $M$  are concyclic.

1

Let  $\angle PLM = \alpha$ .

(iii) Show that  $\angle ABP = \alpha$ .

2

(iv) Hence show that  $M, L$  and  $N$  are collinear.

2

End of question 14

Question 15 (15 marks) Start a new booklet

- (a) The point  $P(a\cos\theta, b\sin\theta)$  lies on the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and the ellipse meets the  $x$ -axis at the points  $A$  and  $A'$ .
- (i) Prove that the tangent at  $P$  has the equation  $\frac{x\cos\theta}{a} + \frac{y\sin\theta}{b} = 1$ . 3
- (ii) The tangent at  $P$  meets the tangents from  $A$  and  $A'$  at points  $Q$  and  $Q'$  respectively. Find the coordinates of  $Q$  and  $Q'$ . 2
- (iii) The points  $A, A', Q'$  and  $Q$  form a trapezium. Prove that the product of the lengths of the parallel sides is independent of the position of  $P$ . 2


- (b) Consider the integral  $I_n = \int_0^1 \frac{x^n}{\sqrt{1-x}} dx$ .
- (i) Show that  $I_1 = \frac{4}{3}$ . 2
- (ii) Show that  $I_{n-1} - I_n = \int_0^1 x^{n-1} \sqrt{1-x} dx$ . (No integration is needed.) 1
- (iii) Use integration by parts on the result of part (ii) to show that  $I_n = \frac{2n}{2n+1} I_{n-1}$ . 2

- (c) (i) Show that  $a^2 + b^2 \geq 2ab$  for any values of  $a$  and  $b$ . 1
- (ii) Hence show that  $\tan^2\theta + \cos^2\theta + \operatorname{cosec}^2\theta \geq \sin\theta + \sec\theta + \cot\theta$  for all values of  $\theta$ . 2

Question 16 (15 marks) Start a new booklet

- (a) Consider the function  $f(x) = \frac{3\sin x}{2 + \cos x}$ .
- (i) Show that  $\frac{3\sin x}{2 + \cos x} < x$  for  $x > 0$ . 3

The diagram shows a circle with centre  $O$ , where  $OA = OB = BC$ ,  $\angle POM = \theta$ ,  $\angle PCO = \alpha$ .



- (ii) Show that  $\tan\alpha = \frac{\sin\theta}{2 + \cos\theta}$ . 2
- (iii) Hence show that  $\alpha < \frac{\theta}{3}$ . 2
- (b) The equation  $x^2 + x + 1 = 0$  has roots  $\alpha$  and  $\beta$ .  
A series is defined by  $T_n = \alpha^n + \beta^n$  for  $n = 1, 2, 3, \dots$
- (i) Show that  $T_1 = -1$  and  $T_2 = -1$ . 2
- (ii) Show that  $T_n = -T_{n-1} - T_{n-2}$  for  $n = 3, 4, 5, \dots$  2
- (iii) Hence use Mathematical Induction to show that  $T_n = 2\cos\frac{2n\pi}{3}$  for  $n = 1, 2, 3, \dots$  3
- (iv) Hence write down the value of  $\sum_{k=1}^{2012} T_k$ . 1

End of paper

2012 Extension 2 Trial Solutions

Section I

1. A                      2. B                      3. C                      4. D                      5. B  
 6. B                      7. D                      8. A                      9. D                      10. C

Section II

Question 11

$$\begin{aligned} \text{(a)} \quad \int_0^1 x e^{-x^2} dx &= -\frac{1}{2} \int_0^1 (-2x) e^{-x^2} dx \\ &= -\frac{1}{2} [e^{-x^2}]_0^1 \\ &= -\frac{1}{2} (e^{-1} - 1) \\ &= \frac{e-1}{2e} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \int \frac{dx}{x^2 + 6x + 10} &= \int \frac{dx}{(x+3)^2 + 1} \\ &= \tan^{-1}(x+3) + c \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int_0^1 \sin^{-1} x dx &= \int_0^1 \sin^{-1} x \cdot \frac{d}{dx}(x) dx \\ &= [x \sin^{-1} x]_0^1 - \int_0^1 x \cdot \frac{1}{\sqrt{1-x^2}} dx \\ &= \frac{\pi}{2} - 0 + \frac{1}{2} \int_0^1 (-2x)(1-x^2)^{-\frac{1}{2}} dx \\ &= \frac{\pi}{2} + \frac{1}{2} \cdot 2 \left[ (1-x^2)^{\frac{1}{2}} \right]_0^1 \\ &= \frac{\pi}{2} - 1 \end{aligned}$$

OR

(by subtraction of areas)

$$\begin{aligned} \int_0^1 \sin^{-1} x dx &= 1 \times \frac{\pi}{2} - \int_0^{\frac{\pi}{2}} \sin x dx \\ &= \frac{\pi}{2} + [\cos x]_0^{\frac{\pi}{2}} \\ &= \frac{\pi}{2} + (0-1) \\ &= \frac{\pi}{2} - 1 \end{aligned}$$

$$\begin{aligned} \text{(d) (i)} \quad \text{Let } \frac{5-5x^2}{(1+2x)(1+x^2)} &= \frac{a}{1+2x} + \frac{bx+c}{1+x^2} \\ 5-5x^2 &= a(1+x^2) + (bx+c)(1+2x) \\ \left(x = -\frac{1}{2}\right) \frac{15}{4} &= \frac{5a}{4} & (x=0) \quad 5 &= a+c \\ a &= 3 & c &= 2 \\ (x=1) \quad 0 &= 2a+3(b+c) \\ 0 &= 6+3(b+2) \\ b &= -4 \end{aligned}$$

$$\begin{aligned} \int_0^1 \frac{5-5x^2}{(1+2x)(1+x^2)} dx &= \int_0^1 \left( \frac{3}{1+2x} - \frac{4x}{1+x^2} + \frac{2}{1+x^2} \right) dx \\ &= \left[ \frac{3}{2} \ln(1+2x) - 2 \ln(1+x^2) + 2 \tan^{-1} x \right]_0^1 \\ &= \frac{3}{2} \ln 3 - 2 \ln 2 + \frac{\pi}{2} - 0 \\ &= \frac{1}{2} (3 \ln 3 - 4 \ln 2 + \pi) \\ &= \frac{1}{2} \left( \ln \frac{3^3}{2^4} + \pi \right) \\ &= \frac{1}{2} \ln \left( \pi + \ln \frac{27}{16} \right) \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \int_0^{\frac{\pi}{4}} \frac{\cos 2x dx}{1+2 \sin 2x + \cos 2x} &= \int_0^1 \frac{\frac{1-t^2}{1+t^2} \cdot \frac{dt}{1+t^2}}{1 + \frac{4t}{1+t^2} + \frac{1-t^2}{1+t^2}} \times \frac{(1+t^2)^2}{(1+t^2)^2} \\ &= \int_0^1 \frac{(1-t^2) dt}{(1+t^2)[(1+t^2)+4t+(1-t^2)]} \\ &= \int_0^1 \frac{(1-t^2) dt}{(1+t^2)(4t+2)} \\ &= \frac{1}{10} \int_0^1 \frac{(5-5t^2) dt}{(1+t^2)(2t+1)} \\ &= \frac{1}{20} \left( \pi + \ln \frac{27}{16} \right) \end{aligned}$$

$$\begin{aligned} \text{Let } t &= \tan x \\ dt &= \sec^2 x dx \\ &= (1 + \tan^2 x) dx \\ dx &= \frac{dt}{1+t^2} \\ x=0, \quad t &= 0 \\ x = \frac{\pi}{4}, \quad t &= 1 \end{aligned}$$

$$\begin{aligned} \text{(c) (i)} \quad \int_{-a}^a f(x) dx &= \int_a^{-a} f(-u)(-du) \\ &= \int_{-a}^a f(-x) dx \end{aligned}$$

$$\begin{aligned} \text{let } u &= -x \\ du &= -dx \\ x = -a, \quad u &= a \\ x = a, \quad u &= -a \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \int_{-4}^4 (e^x - e^{-x}) \cos x dx &= \int_{-4}^4 (e^{-x} - e^x) \cos(-x) dx \quad (\text{from part i}) \\ &= - \int_{-4}^4 (e^x - e^{-x}) \cos x dx \end{aligned}$$

$$\begin{aligned} 2 \int_{-4}^4 (e^x - e^{-x}) \cos x dx &= 0 \\ \int_{-4}^4 (e^x - e^{-x}) \cos x dx &= 0 \end{aligned}$$



**Question 12**

(a) Let  $\sqrt{6i-8} = a+ib$  (where  $a$  and  $b$  are real)  
 $(a^2 - b^2) + 2abi = -8 + 6i$

Equating real and imaginary parts:

$$2ab = 6 \qquad a^2 - b^2 = -8$$

$$b = \frac{3}{a} \qquad a^2 - \frac{9}{a^2} = -8$$

$$(\times a^2) \quad a^4 + 8a^2 - 9 = 0$$

$$(a^2 + 9)(a^2 - 1) = 0$$

$$a = \pm 1$$

$$b = \pm 3$$

$$\sqrt{6i-8} = \pm(1+3i)$$

**Alternatively**

As  $|6i-8|=10$ , then  $|a+ib|=\sqrt{10}$ , so  $a^2+b^2=10$ , then solve this with the 2<sup>nd</sup> equation by elimination, substituting the answers in the 1<sup>st</sup> equation to find the second pronomeral.

(b) (i)  $2+2\sqrt{3}i = 4 \operatorname{cis} \frac{\pi}{3}$

(ii)  $(2+2\sqrt{3}i)^3 = \left(4 \operatorname{cis} \frac{\pi}{3}\right)^3$   
 $= 64 \operatorname{cis} \pi$   
 $= -64$


(iii)  $z^4 = 4 \operatorname{cis} \left(\frac{\pi}{3} + 2n\pi\right)$ , where  $n$  is an integer  
 $= 4 \operatorname{cis} \left(\frac{6n+1}{3}\pi\right)$

$$z = \sqrt{2} \operatorname{cis} \left(\frac{6n+1}{12}\pi\right)$$

Taking  $n = -2, -1, 0, 1$ :

$$z = \sqrt{2} \operatorname{cis} \frac{\pi}{12}, \sqrt{2} \operatorname{cis} \frac{7\pi}{12}, \sqrt{2} \operatorname{cis} \left(-\frac{5\pi}{12}\right), \sqrt{2} \operatorname{cis} \left(-\frac{11\pi}{12}\right)$$

(c)



(d) (i)  $\angle BAP = 2\theta$  (angle at centre is twice angle at circumference)

$$\therefore \overline{AP} = r \operatorname{cis} 2\theta$$

(ii)  $\angle OPB = \frac{\pi}{2}$  (angle in semi-circle)

$$\angle PBx = \theta + \frac{\pi}{2} \quad (\text{exterior angle of triangle} = \text{sum of opposite two interior angles})$$

$$PB^2 = (2r)^2 - a^2 \quad (\text{Pythagoras'})$$

$$\therefore \overline{BP} = \sqrt{4r^2 - a^2} \operatorname{cis} \left(\theta + \frac{\pi}{2}\right)$$

(c) (i)  $3^{12}$

(ii)  $\frac{{}^{12}C_4 \times {}^8C_4}{3^{12}} = \frac{3850}{59049}$

NB: We don't divide by 3!, as the 3 groups are considered different, and are enumerated as different cases when the sample space is calculated in part (i).

**OR**

Name the switch positions A, B, C.

The question is the same as forming distinct words from the letters A A A A B B B B C C C C.

$$\text{ie. } \frac{\frac{12!}{(4!)^3}}{3^{12}} = \frac{3850}{59049}$$

Question 13

(a) (i)  $y = \frac{2x}{1+x^2}$


$$\frac{dy}{dx} = \frac{(1+x^2) \cdot 2 - 2x \cdot 2x}{(1+x^2)^2}$$

$$= \frac{2-2x^2}{(1+x^2)^2}$$


Stat Pts:  $\frac{dy}{dx} = 0 \Rightarrow x = \pm 1, y = \pm 1$

ie. stat points at  $(1,1)$  and  $(-1,-1)$


(ii) 1.  $y = \frac{|2x|}{1+x^2}$




2.  $y = \frac{1+x^2}{2x}$



3.  $y^2 = \frac{2x}{1+x^2}$



4.  $y = \ln\left(\frac{2x}{1+x^2}\right)$



(b) (i) Let  $P(x) = (x-\alpha)^2 Q(x)$ .

$$P'(x) = Q(x) \cdot 2(x-\alpha) + (x-\alpha)^2 \cdot Q'(x)$$

$$= (x-\alpha)[2Q(x) + (x-\alpha)Q'(x)]$$

$$\therefore P'(\alpha) = 0$$

So  $x = \alpha$  is a root of  $P(x) = 0$ .

(ii) If  $y = mx$  is a tangent, then  $mx = 3 - \frac{1}{x^2}$  has a double root.

ie.  $mx^3 = 3x^2 - 1$

$$mx^3 - 3x^2 + 1 = 0$$

(iii) Let  $P(x) = mx^3 - 3x^2 + 1$

$$P'(x) = 3mx^2 - 6x$$

By part (i), the double root  $x = \alpha$  must be a root of

$$3mx^2 - 6x = 0$$

$$3x(mx-2) = 0$$

$$x = 0 \text{ or } x = \frac{2}{m}$$

$P(0) \neq 0$ , so  $x = \frac{2}{m}$  must be the double root.

$$P\left(\frac{2}{m}\right) = 0$$

$$m\left(\frac{2}{m}\right)^3 - 3\left(\frac{2}{m}\right)^2 + 1 = 0$$

$$\frac{8}{m^2} - \frac{12}{m^2} + 1 = 0$$


$$\frac{4}{m^2} = 1$$

$$m = \pm 2$$

So the tangents have equation  $y = \pm 2x$ .

Question 14

(a)



Outer radius,  $R = 1 - x$

Inner radius,  $r = 1 - x - \delta x$

Height,  $h = y = \frac{1}{2x+1}$

Volume of typical slice:


$$\delta V \approx \pi(R^2 - r^2)h$$

$$= \pi(R+r)(R-r)h$$

$$= \pi(2 - 2x - \delta x)(\delta x) \cdot \frac{1}{2x+1}$$

$$\approx 2\pi \cdot \frac{1-x}{2x+1} \cdot \delta x \quad \text{when } \delta x \text{ is sufficiently small}$$

OR



$$\delta V \approx 2\pi(1-x) \cdot \frac{1}{2x+1} \cdot \delta x$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^1 2\pi \frac{1-x}{2x+1} \delta x$$

$$= 2\pi \int_0^1 \frac{1-x}{2x+1} dx$$

$$= 2\pi \int_0^1 \left[ \frac{1}{2} - \frac{1}{2} \frac{(2x+1) + \frac{3}{2}}{2x+1} \right] dx$$

$$= 2\pi \int_0^1 \left( -\frac{1}{2} + \frac{3}{2(2x+1)} \right) dx$$

$$= 2\pi \left[ -\frac{x}{2} + \frac{3}{4} \ln(2x+1) \right]_0^1$$

$$= 2\pi \left( -\frac{1}{2} + \frac{3}{4} \ln 3 - 0 \right)$$

$$= \frac{\pi}{2} (3 \ln 3 - 2) \text{ units}^3$$

(b) (i) Period =  $2b$

$$\frac{2\pi}{n} = 2b$$

$$n = \frac{\pi}{b}$$

(ii) Area =  $\int_0^b b \sin \frac{\pi}{b} x dx$

$$= -\frac{b^2}{\pi} \left[ \cos \frac{\pi}{b} x \right]_0^b$$

$$= -\frac{b^2}{\pi} (-1 - 1)$$

$$= \frac{2b^2}{\pi} \text{ units}^2$$

(iii) from part (ii),  $\delta V = \frac{2b^2}{\pi} \delta y$

$$= \frac{2(2x)^2}{\pi} \delta y$$

$$= \frac{8}{\pi} x^2 \delta y$$

$$= \frac{8}{\pi} (4ay) \delta y$$

$$= \frac{32a}{\pi} y \delta y$$

$$V = \lim_{\delta y \rightarrow 0} \sum_{y=0}^a \frac{32a}{\pi} y \delta y$$

$$= \frac{32a}{\pi} \int_0^a y dy$$

$$= \frac{16a}{\pi} [y^2]_0^a$$

$$V = \frac{16}{\pi} a^3 \text{ units}^3$$

(c) (i)  $BP$  subtends equal angles at  $N$  and  $L$  (converse of angles in same segment)

(ii)  $\angle PLC + \angle PMC = 90^\circ + 90^\circ$

$$= 180^\circ$$

$\therefore PLCM$  is cyclic (opposite angles are supplementary)

(iii) Construct  $BP$  and  $PM$ .

$$\angle PCM = \angle PLM = \alpha$$

(angles in same segment of circle  $PLCM$ )

$$\angle ABP = \angle PCM = \alpha$$

(exterior angle or cyclic quad  $BACP$  = opposite interior angle)

(iii) Construct  $NL$ .

$$\angle NLP + \angle NBP = 180^\circ$$

(opposite angles of cyclic quad  $PLNB$  are supplementary)

$$\angle NLP = 180^\circ - \alpha$$

$$\angle NLP + \angle PLM = (180 - \alpha) + \alpha$$

$$\angle MLN = 180^\circ$$

ie.  $M, L,$  and  $N$  are collinear

NB: We can't call  $\angle PLM$  the exterior angle of  $PLNB$  until we know that  $MLN$  is a straight line.

Question 15

(a) (i)  $x = a \cos \theta$        $y = b \sin \theta$        $\frac{dy}{dx} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dx}$   
 $\frac{dx}{d\theta} = -a \sin \theta$        $\frac{dy}{d\theta} = b \cos \theta$        $= b \cos \theta \cdot \frac{-1}{a \sin \theta}$   
 $= -\frac{b \cos \theta}{a \sin \theta}$

$$y - b \sin \theta = -\frac{b \cos \theta}{a \sin \theta} (x - a \cos \theta)$$

$$ay \sin \theta - ab \sin^2 \theta = -bx \cos \theta + ab \cos^2 \theta$$

$$bx \cos \theta + ay \sin \theta = ab (\sin^2 \theta + \cos^2 \theta)$$

$$(+ab) \frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$$

(ii)  $(x = \pm a) \frac{\pm a \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$

$$\frac{y \sin \theta}{b} = 1 \mp \cos \theta$$

$$y = \frac{b(1 \mp \cos \theta)}{\sin \theta}$$

ie.  $Q\left(a, \frac{b(1 - \cos \theta)}{\sin \theta}\right), Q\left(-a, \frac{b(1 + \cos \theta)}{\sin \theta}\right)$

(iii) Product of lengths  $= \frac{b(1 - \cos \theta)}{|\sin \theta|} \cdot \frac{b(1 + \cos \theta)}{|\sin \theta|}$

$$= \frac{b^2(1 - \cos^2 \theta)}{\sin^2 \theta}$$

$$= \frac{b^2 \sin^2 \theta}{\sin^2 \theta}$$

$$= b^2 \quad (\text{which is a constant})$$

(b) (i)  $I_1 = \int_0^1 \frac{x dx}{\sqrt{1-x}}$   
 $= \int_0^1 \frac{-(1-x)+1}{\sqrt{1-x}} dx$   
 $= \int_0^1 \left( -(1-x)^{\frac{1}{2}} + (1-x)^{-\frac{1}{2}} \right) dx$   
 $= \left[ \frac{2}{3}(1-x)^{\frac{3}{2}} - 2(1-x)^{\frac{1}{2}} \right]_0^1$   
 $= (0-0) - \left( \frac{2}{3} - 2 \right)$   
 $= \frac{4}{3}$

Other options:

1. Let  $u = 1-x$

2. Let  $u^2 = 1-x$

3. Let  $x = \sin^2 \theta$  (messy)

4. Integration by parts

(ii)  $I_{n-1} - I_n = \int_0^1 \frac{x^{n-1}}{\sqrt{1-x}} dx - \int_0^1 \frac{x^n}{\sqrt{1-x}} dx$   
 $= \int_0^1 \frac{x^{n-1} - x^n}{\sqrt{1-x}} dx$   
 $= \int_0^1 \frac{x^{n-1}(1-x)}{\sqrt{1-x}} dx$   
 $= \int_0^1 x^{n-1} \sqrt{1-x} dx$

(iii)  $I_{n-1} - I_n = \int_0^1 x^{n-1} \sqrt{1-x} dx$   
 $= \int_0^1 \sqrt{1-x} \cdot \frac{d}{dx} \left( \frac{x^n}{n} \right) dx$   
 $= \frac{1}{n} \left[ x^n \sqrt{1-x} \right]_0^1 - \frac{1}{n} \int_0^1 x^n \cdot \frac{1}{2} (1-x)^{-\frac{1}{2}} \cdot (-1) dx$   
 $= 0 + \frac{1}{2n} \int_0^1 \frac{x^n dx}{\sqrt{1-x}}$

$$I_{n-1} - I_n = \frac{1}{2n} I_n$$

$$2nI_{n-1} - 2nI_n = I_n$$

$$2nI_{n-1} = (2n+1)I_n$$

$$I_n = \frac{2n}{2n+1} I_{n-1}$$

(c) (i)  $(a-b)^2 \geq 0 \quad \forall a, b$   
 $a^2 - 2ab + b^2 \geq 0$   
 $a^2 + b^2 \geq 2ab$

(ii) From (i),  $\tan^2 \theta + \cos^2 \theta \geq 2 \tan \theta \cos \theta = 2 \sin \theta$   
 $\tan^2 \theta + \operatorname{cosec}^2 \theta \geq 2 \tan \theta \operatorname{cosec} \theta = 2 \sec \theta$   
 $\cos^2 \theta + \operatorname{cosec}^2 \theta \geq 2 \cos \theta \operatorname{cosec} \theta = 2 \cot \theta$   
 Adding:  $2(\tan^2 \theta + \cos^2 \theta + \operatorname{cosec}^2 \theta) \geq 2(\sin \theta + \sec \theta + \cot \theta)$   
 $\tan^2 \theta + \cos^2 \theta + \operatorname{cosec}^2 \theta \geq \sin \theta + \sec \theta + \cot \theta$

Question 16

(a) (i) Let  $f(x) = x - \frac{3 \sin x}{2 + \cos x}$ .

$f(0) = 0$

$f'(x) = 1 - \frac{(2 + \cos x)(3 \cos x) - (3 \sin x)(-\sin x)}{(2 + \cos x)^2}$

$= \frac{(2 + \cos x)^2 - 3 \cos x(2 + \cos x) - 3 \sin^2 x}{(2 + \cos x)^2}$

$= \frac{4 + 4 \cos x + \cos^2 x - 6 \cos x - 3 \cos^2 x - 3 + 3 \cos^2 x}{(2 + \cos x)^2}$

$= \frac{1 - 2 \cos x + \cos^2 x}{(2 + \cos x)^2}$

$= \left( \frac{1 - \cos x}{2 + \cos x} \right)^2$

$\geq 0 \quad \forall x$

$\therefore f(x) > 0 \quad \forall x > 0$  (starts at zero, and decreases monotonically)

$\therefore x - \frac{3 \sin x}{2 + \cos x} > 0$

$\therefore \frac{3 \sin x}{2 + \cos x} < x \quad \text{for } x > 0$

(ii) Let  $OB = OP = r$

from  $\triangle MOP$ ,  $OM = r \cos \theta$  and  $PM = r \sin \theta$

In  $\triangle CMP$ ,  $\tan \alpha = \frac{PM}{CM}$

$= \frac{PM}{CO + OM}$

$= \frac{r \sin \theta}{2r + r \cos \theta}$

$\tan \theta = \frac{\sin \theta}{2 + \cos \theta}$

(iii)  $\tan \alpha = \frac{1}{3} \cdot \frac{3 \sin \theta}{2 + \cos \theta}$

$< \frac{1}{3} \theta$  (from part i, since  $\theta > 0$ )

Also, for  $0 < \alpha < \frac{\pi}{2}$ ,  $\alpha < \tan \alpha$ .

$\therefore \alpha < \frac{\theta}{3}$

(b) (i)  $T_1 = \alpha + \beta$

$= -1$

$= 1$

$= -1$

$T_2 = \alpha^2 + \beta^2$

$= (\alpha + \beta)^2 - 2\alpha\beta$

$= (-1)^2 - 2(1)$

$= -1$

(ii) Since  $x^2 + x + 1 = 0$  has roots  $\alpha$  and  $\beta$ , then:

$(\times x^{n-2}) \quad x^n + x^{n-1} + x^{n-2} = 0$  has roots  $\alpha$  and  $\beta$  (and a root 0 of multiplicity  $n-2$ )

$\therefore \alpha^n + \alpha^{n-1} + \alpha^{n-2} = 0$

and  $\beta^n + \beta^{n-1} + \beta^{n-2} = 0$

adding:  $(\alpha^n + \beta^n) + (\alpha^{n-1} + \beta^{n-1}) + (\alpha^{n-2} + \beta^{n-2}) = 0$

$T_n + T_{n-1} + T_{n-2} = 0$

$T_n = -T_{n-1} - T_{n-2}$

OR

RHS  $= -T_{n-1} - T_{n-2}$

$= -(\alpha^{n-1} + \beta^{n-1}) - (\alpha^{n-2} + \beta^{n-2})$

$= -\left[ \left( \frac{\alpha^n}{\alpha} + \frac{\alpha^n}{\alpha^2} \right) + \left( \frac{\beta^n}{\beta} + \frac{\beta^n}{\beta^2} \right) \right]$

$= -\left[ \alpha^n \left( \frac{1+\alpha}{\alpha^2} \right) + \beta^n \left( \frac{1+\beta}{\beta^2} \right) \right]$

$= -[\alpha^n(-1) + \beta^n(-1)]$

$[1 + \alpha + \alpha^2 = 1 + \beta + \beta^2 = 0]$

$= \alpha^n + \beta^n$

$= T_n$

(iii) Test  $n=1$  and  $n=2$ :

RHS  $= 2 \cos \frac{2\pi}{3} = -1 = T_1 = \text{LHS}$

RHS  $= 2 \cos \frac{4\pi}{3} = -1 = T_2 = \text{LHS}$

$\therefore$  true for  $n=1$  and  $n=2$

Assume true for  $n=k$  and  $n=k+1$ :

ie.  $T_k = 2 \cos \frac{2k\pi}{3}$  and  $T_{k+1} = 2 \cos \frac{2(k+1)\pi}{3}$

Prove true for  $n=k+2$ :

ie. Prove  $T_{k+2} = 2 \cos \frac{2(k+2)\pi}{3}$

$T_{k+2} = -T_{k+1} - T_k$  (from part ii)

$= -2 \cos \frac{2(k+1)\pi}{3} - 2 \cos \frac{2k\pi}{3}$  (by assumption)

$= -2 \cos \left( \frac{2k\pi}{3} + \frac{2\pi}{3} \right) - 2 \cos \frac{2k\pi}{3}$

$= -2 \left( \cos \frac{2k\pi}{3} \cos \frac{2\pi}{3} - \sin \frac{2k\pi}{3} \sin \frac{2\pi}{3} + \cos \frac{2k\pi}{3} \right)$

$= -2 \left( -\frac{1}{2} \cos \frac{2k\pi}{3} - \frac{\sqrt{3}}{2} \sin \frac{2k\pi}{3} + \cos \frac{2k\pi}{3} \right)$

$= -2 \left( \frac{1}{2} \cos \frac{2k\pi}{3} - \frac{\sqrt{3}}{2} \sin \frac{2k\pi}{3} \right)$

$$\begin{aligned}
&= 2 \left( -\frac{1}{2} \cos \frac{2k\pi}{3} + \frac{\sqrt{3}}{2} \sin \frac{2k\pi}{3} \right) \\
&= 2 \left( \cos \frac{2k\pi}{3} \cos \frac{4\pi}{3} - \sin \frac{2k\pi}{3} \sin \frac{4\pi}{3} \right) \\
&= 2 \cos \left( \frac{2k\pi}{3} + \frac{4\pi}{3} \right) \\
&= 2 \cos \frac{2(k+2)\pi}{3}
\end{aligned}$$

$\therefore$  True for  $n = k + 2$  when true for  $n = k$  and  $n = k + 1$

$\therefore$  By Mathematical Induction,  $T_n = 2 \cos \frac{2n\pi}{3}$  for  $n = 1, 2, 3, \dots$

$$\begin{aligned}
\text{(iv) } \sum_{k=1}^{2012} T_k &= (-1) + (-1) + 2 + (-1) + (-1) + 2 + \dots \\
&= -2 \quad (\text{since } 2012 \text{ is two more than a multiple of } 3)
\end{aligned}$$