



NORTH SYDNEY GIRLS HIGH SCHOOL
YEAR 12 – TERM 1 ASSESSMENT

2006

MATHEMATICS EXTENSION 1

TIME ALLOWED: 60 minutes
 Plus 2 minutes reading time

INSTRUCTIONS:

- Start each question on a new page
- Hand each question in separately, including a sheet for non-attempts
- Show all necessary working

This task is worth 20% of the HSC Assessment Mark

Question One (9 Marks)

- | | |
|--|---|
| (a) What is the exact value of $\cos\left(\frac{\pi}{6}\right)$? | 1 |
| (b) Differentiate $\cos(x^2 + 1)$ | 1 |
| (c) Find $\int \sec^2 5x \, dx$ | 1 |
| (d) (i) Sketch the curve $y = 4\sin 2x$ for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ | 2 |
| (ii) On your diagram for part (i), sketch the line $y = \frac{1}{3}x$, and shade the region represented by $\int_0^{\frac{\pi}{4}} \left(4\sin 2x - \frac{1}{3}x\right) dx$ | 2 |
| (iii) Find the exact value of the integral in part (ii). | 2 |

Question Two (8 Marks)

- | | |
|--|---|
| (a) A sphere is being heated so that its surface area is increasing at a constant rate of 25 cm^2 per second. Find the rate of increase of the volume when the radius is 5 cm. | 3 |
| (b) One hundred grams of cane sugar in water are being converted into dextrose at a rate which is proportional to the amount unconverted at any time, i.e. if M grams are converted in t minutes, then $\frac{dM}{dt} = k(100 - M)$, where k is a constant. | |
| (i) Show that $M = 100 + A e^{-kt}$, where A is a constant which satisfies the above equation. | 2 |
| (ii) Find the value of A (initially no cane sugar has been converted to dextrose) | 1 |
| (iii) If 40 grams are converted in the first 15 minutes, find how many grams are converted in the first 30 minutes. | 2 |

Question Three (10 Marks)

- | | |
|---|---|
| (a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin\left(\frac{x}{4}\right)}{3x}$ | 2 |
| (b) (i) Express $\cos x - \sin x$ in the form $R \cos(x + \alpha)$, where $R > 0$ and $0 \leq \alpha \leq \frac{\pi}{2}$ | 2 |
| (ii) Hence, or otherwise, solve the equation $\cos x - \sin x = \frac{\sqrt{2}}{2}$ for $0 \leq x \leq 2\pi$ | 2 |
| (c) Prove $\frac{\tan A}{\tan 2A - \tan A} \equiv \cos 2A$ | 4 |

Question Four (10 Marks)

- (a) Find the exact value of $\sin 105^\circ$ 2
- (b) Find the volume of the solid generated when $y = \sin 3x$ is rotated around the x axis from $x = 0$ to $x = \frac{\pi}{3}$. 4
- (c) Differentiate $x \sin 3x$ with respect to x and hence evaluate $\int_0^{\frac{\pi}{2}} x \cos 3x \, dx$ 4

Question Five (9 Marks)

- (a) If $y = \tan 2x$, find the equation of the tangent to the curve at $x = \frac{\pi}{6}$ 3
- (b) Find the acute angle between the lines $4x + y + 5 = 0$ and $6x + 3y - 7 = 0$ correct to the nearest minute. 3
- (c) Solve the equation $\sin 2\theta + \cos \theta = 0$ for $0 \leq \theta \leq 2\pi$ 3

Question Six (10 Marks)

- (a) Solve $5 \sin \theta - 2 \cos \theta = 2$ for $0^\circ \leq \theta \leq 360^\circ$, using the result that $\tan \frac{\theta}{2} = t$ 4
- (b) A particle moves along a straight line so that its displacement, x metres, from a fixed point O is given by $x = 1 - 3 \cos\left(\frac{t}{2}\right)$, where t is measured in seconds.
- (i) Sketch the graph of x as a function of t for $0 \leq t \leq 4\pi$ 2
- (ii) Hence, or otherwise, find when and where the particle first comes to rest after $t = 0$ 2
- (iii) Find a time when the particle reaches its maximum speed. What is this speed? 2

END OF TEST


Question 1

a) $\frac{\sqrt{3}}{2}$ ✓

b) $\frac{d(\cos(x^2+1))}{dx} = -2x \sin(x^2+1)$ ✓

c) $\int \sec^2 5x \, dx = \frac{1}{5} \tan 5x + C$ ✓

d) (i) (ii)



(ii) $\int_0^{\pi/4} (4 \sin 2x - \frac{1}{3}x^2) \, dx = \left[-2 \cos 2x - \frac{1}{6}x^2 \right]_0^{\pi/4}$
 $= -2 \cos \frac{\pi}{2} - \frac{1}{6} \left(\frac{\pi}{4}\right)^2 + 2 \cos 0 + 0$
 $= 2 - \frac{\pi^2}{96}$ ✓

Question 2

a) $\frac{dV}{dt} = 25 \text{ cm}^3/\text{s}$ $V = \frac{4}{3}\pi r^3$ $A = 4\pi r^2$
 $\frac{dV}{dr} = 4\pi r^2$ $\frac{dA}{dr} = 8\pi r$
 $\frac{dV}{dr} = \frac{dV}{dA} \times \frac{dA}{dr}$
 $\frac{dV}{dA} = \frac{dV}{dr} \times \frac{dr}{dA} = \frac{4\pi r^2}{8\pi r} = \frac{r}{2}$
 $\therefore \frac{dV}{dA} = \frac{r}{2} \times 25 = \frac{5}{2} \times 25 \text{ cm}^3/\text{s} = \frac{125}{2} \text{ cm}^3/\text{s} = 62\frac{1}{2} \text{ cm}^3/\text{s}$

Question 2

b) (i) $\frac{dM}{dt} = 100 - Mk$

$\frac{dt}{dM} = \frac{1}{100 - Mk}$ ✓

$t = -\frac{1}{k} \ln(100 - M) + C$

$t = 0 \Rightarrow -\frac{1}{k} \ln(100 - M) = C$

$k(C - t) = \ln(100 - M)$

$e^{k(C-t)} = 100 - M$

let $A = e^{kC}$

$100 - Ae^{-kt} = M$

(ii) $0 = 100 - A$
 $\therefore A = 100$

(iii) $40 = 100 - 100e^{-15k}$

$60 = 100e^{-15k}$

$\ln \frac{60}{100} = -15k$
 $k = \frac{\ln \frac{60}{100}}{-15}$

$t = 30$

$M = 100 - 100e^{-k \cdot 30}$

$= 64$

64 gms are converted.

Ques 3

$$\begin{aligned} a) \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{3})}{3x} &= \frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{3})}{\frac{x}{3}} \\ &= \frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{3})}{\frac{x}{3}} \\ &= \frac{1}{3} \end{aligned}$$

b) (i) $\cos x - \sin x \Rightarrow \cos x = 1 \quad \therefore R = \sqrt{2}$
 $\sin x = 1$
 $\therefore \cos x - \sin x = \sqrt{2} \cos(x + \frac{\pi}{4})$

(ii) $\cos x - \sin x = \frac{\sqrt{2}}{2}$
 $\sqrt{2} \cos(x + \frac{\pi}{4}) = \frac{\sqrt{2}}{2}$
 $\cos(x + \frac{\pi}{4}) = \frac{1}{2}$

$-\frac{\pi}{4} \leq x \leq \frac{7\pi}{4}$
 $x + \frac{\pi}{4} = \frac{\pi}{3}, \frac{5\pi}{3} \quad \therefore 0 \leq x + \frac{\pi}{4} \leq 2\pi$
 $x = -\frac{\pi}{12}, \frac{17\pi}{12}$

c) LHS = $\frac{\tan A}{\tan 2A - \tan A} = \frac{\tan A}{2 \tan A - \tan A} = \frac{\tan A}{\tan A} = 1$
 $\frac{\tan A}{2 \tan A - \tan A} = \frac{\tan A}{\tan A} = 1$
 $\frac{\tan A}{2 \tan A - \tan A} = \frac{\tan A}{\tan A} = 1$
 $\frac{\tan A}{2 \tan A - \tan A} = \frac{\tan A}{\tan A} = 1$
 $\frac{\tan A}{2 \tan A - \tan A} = \frac{\tan A}{\tan A} = 1$

Ques 4

a) $\sin(105^\circ) = \sin(60 + 45)$
 $= \sin 60 \cos 45 + \sin 45 \cos 60$
 $= \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \times \frac{1}{2}$
 $= \frac{\sqrt{3} + 1}{2\sqrt{2}}$
 $= \frac{\sqrt{6} + \sqrt{2}}{4}$

b) $\int_0^{\frac{\pi}{3}} \sin^2 3x \, dx$
 $= \int_0^{\frac{\pi}{3}} \frac{1 - \cos 6x}{2} \, dx$
 $= \frac{\pi}{6} \left[x - \frac{\sin 6x}{6} \right]_0^{\frac{\pi}{3}}$
 $= \frac{\pi^2}{6}$

c) $\int_0^{\frac{\pi}{2}} x \cos 3x \, dx$
 $= \frac{1}{3} \int_0^{\frac{\pi}{2}} 3x \cos 3x \, dx + \sin 3x - \sin 3x$
 $= \frac{1}{3} \left[x \sin 3x + \frac{1}{3} \cos 3x \right]_0^{\frac{\pi}{2}}$
 $= \frac{1}{3} \left[-\frac{\pi}{2} - \frac{1}{3} \right]$
 $= -\left(\frac{\pi}{6} + \frac{1}{9} \right)$

Question 5

a) $y = \tan 2x$
 $y' = 2 \sec^2 2x$

at $x = (\frac{\pi}{6}, \sqrt{3})$

$\therefore y' = 8 \checkmark$

$\therefore y - \sqrt{3} = 8(x - \frac{\pi}{6})$

$8x - y - \frac{\pi}{6} + \sqrt{3} = 0 \checkmark$

b) $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$
 $m_1 = -4$
 $m_2 = -2$
 $= \left| \frac{-2}{9} \right|$

$\theta \approx \cancel{150^\circ 57'} 12' 32'$

c) $\sin 2\theta + \cos \theta = 0$

$2 \sin \theta \cos \theta + \cos \theta = 0$

$\cos \theta (\sin \theta + 1) = 0$

$\cos \theta = 0$

$\theta = \frac{\pi}{2}, \frac{3\pi}{2} \checkmark$

$\sin \theta = -\frac{1}{2}$

$\theta = \frac{7\pi}{6}, \frac{11\pi}{6} \checkmark$

$\therefore \theta = \frac{\pi}{2}, \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6} \checkmark$

Question 6

a) $5 \sin \theta - 2 \cos \theta = 2$ let $\tan \frac{\theta}{2} = t \therefore \sin \theta = \frac{2t}{1+t^2}$

$\frac{10t}{1+t^2} - \frac{2(1-t^2)}{1+t^2} = 2$

$\frac{10t - 2 + 2t^2}{1+t^2} = 2$

Question 6

c) cont'd. $\frac{2t^2 + 10t - 2}{1+t^2} = 2$

$2t^2 + 10t - 2 = 2 + 2t^2$ $10t = 4$

$2t^2 - 10t + 4 = 0$ $t = \frac{1}{5}$

$t = \frac{10 \pm \sqrt{100 - 32}}{4}$

$= \frac{10 \pm \sqrt{68}}{4}$

$= \frac{10 \pm 2\sqrt{17}}{4}$

$= \frac{5 \pm \sqrt{17}}{2}$

$\therefore \tan \frac{\theta}{2} = \frac{1}{5}$ $0 \leq \frac{\theta}{2} \leq 180^\circ$

$\frac{\theta}{2} = 11^\circ 19'$

$\theta = 22^\circ 38'$

Also need to test for $\theta = 180^\circ$

$\therefore 5 \sin \theta - 2 \cos \theta = 2$

$0 - 2(-1) = 2$

is true


$\therefore \theta = 180^\circ$ is also a soln.

$\therefore \theta \approx 155^\circ 16'$

$47^\circ 21'$

180°

b) (i)



(ii) $x = 1 - 3 \cos(\frac{t}{2})$

$x = \frac{3}{2} \sin(\frac{t}{2})$

\therefore parallel 1st stationary at $t = 2\pi$

$n = 4$

For $x = 0$, $t = 2\pi$

Prob 6

b) (iii) max speed at $\ddot{x}=0$ i.e. point of inflection
at $t=\pi$ or 3π .

$$\ddot{x} = \frac{3}{4} \cos\left(\frac{t}{2}\right) = 0$$

$$\therefore \underline{\underline{t = \pi}} \checkmark \text{ the speed is } \underline{\underline{\frac{3}{2} \text{ m/s}}} \checkmark$$