



NORTH SYDNEY GIRLS HIGH SCHOOL  
YEAR 12 – TERM 1 ASSESSMENT

2006

MATHEMATICS EXTENSION 1

TIME ALLOWED: 60 minutes  
Plus 2 minutes reading time

INSTRUCTIONS:

- Start each question on a new page
- Hand each question in separately, including a sheet for non-attempts
- Show all necessary working

This task is worth 20% of the HSC Assessment Mark

Question One (9 Marks)

- |  | Marks |
|--|-------|
| (a) What is the exact value of $\cos\left(\frac{\pi}{6}\right)$ ?  | 1     |
| (b) Differentiate $\cos(x^2 + 1)$  | 1     |
| (c) Find $\int \sec^2 5x \, dx$  | 1     |
| (d) (i) Sketch the curve $y = 4\sin 2x$ for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$   | 2     |
| (ii) On your diagram for part (i), sketch the line $y = \frac{1}{3}x$ , and shade the region represented by $\int_0^{\frac{\pi}{4}} \left(4\sin 2x - \frac{1}{3}x\right) dx$ | 2     |
| (iii) Find the exact value of the integral in part (ii).   | 2     |

Question Two (8 Marks)

- |  |   |
|--|---|
| (a) A sphere is being heated so that its surface area is increasing at a constant rate of $25 \text{ cm}^2$ per second. Find the rate of increase of the volume when the radius is 5 cm.   | 3 |
| (b) One hundred grams of cane sugar in water are being converted into dextrose at a rate which is proportional to the amount unconverted at any time, i.e. if $M$ grams are converted in $t$ minutes, then $\frac{dM}{dt} = k(100 - M)$ , where $k$ is a constant. |   |
| (i) Show that $M = 100 + A e^{-kt}$ , where $A$ is a constant which satisfies the above equation.  | 2 |
| (ii) Find the value of $A$ (initially no cane sugar has been converted to dextrose)  | 1 |
| (iii) If 40 grams are converted in the first 15 minutes, find how many grams are converted in the first 30 minutes.  | 2 |

Question Three (10 Marks)

- |   |   |
|---|---|
| (a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin\left(\frac{x}{4}\right)}{3x}$   | 2 |
| (b) (i) Express $\cos x - \sin x$ in the form $R \cos(x + \alpha)$ , where $R > 0$ and $0 \leq \alpha \leq \frac{\pi}{2}$ | 2 |
| (ii) Hence, or otherwise, solve the equation $\cos x - \sin x = \frac{\sqrt{2}}{2}$ for $0 \leq x \leq 2\pi$              | 2 |
| (c) Prove $\frac{\tan A}{\tan 2A - \tan A} \equiv \cos 2A$  | 4 |

**Question Four** (10 Marks)

- (a) Find the exact value of  $\sin 105^\circ$  2
- (b) Find the volume of the solid generated when  $y = \sin 3x$  is rotated around the  $x$  axis from  $x = 0$  to  $x = \frac{\pi}{3}$ . 4
- (c) Differentiate  $x \sin 3x$  with respect to  $x$  and hence evaluate  $\int_0^{\frac{\pi}{2}} x \cos 3x \, dx$  4

**Question Five** (9 Marks)

- (a) If  $y = \tan 2x$ , find the equation of the tangent to the curve at  $x = \frac{\pi}{6}$  3
- (b) Find the acute angle between the lines  $4x + y + 5 = 0$  and  $6x + 3y - 7 = 0$  correct to the nearest minute. 3
- (c) Solve the equation  $\sin 2\theta + \cos \theta = 0$  for  $0 \leq \theta \leq 2\pi$  3

**Question Six** (10 Marks)

- (a) Solve  $5 \sin \theta - 2 \cos \theta = 2$  for  $0^\circ \leq \theta \leq 360^\circ$ , using the result that  $\tan \frac{\theta}{2} = t$  4
- (b) A particle moves along a straight line so that its displacement,  $x$  metres, from a fixed point  $O$  is given by  $x = 1 - 3 \cos\left(\frac{t}{2}\right)$ , where  $t$  is measured in seconds.
- (i) Sketch the graph of  $x$  as a function of  $t$  for  $0 \leq t \leq 4\pi$  2
- (ii) Hence, or otherwise, find when and where the particle first comes to rest after  $t = 0$  2
- (iii) Find a time when the particle reaches its maximum speed. What is this speed? 2

END OF TEST


Question 1

a)  $\frac{\sqrt{3}}{2} \checkmark$

b)  $\frac{d(\cos(x^2+1))}{dx} = -2x \sin(x^2+1) \checkmark$

c)  $\int \sec^2 5x \, dx = \frac{1}{5} \tan 5x + C \checkmark$

d) (i) (ii)



(ii)  $\int_0^{\pi/4} (4 \sin 2x - \frac{1}{3}x) \, dx = \left[ -2 \cos 2x - \frac{1}{6}x^2 \right]_0^{\pi/4}$

$= -\frac{\pi^2}{96} + 2$

$= 2 - \frac{\pi^2}{96} \checkmark$

Question 2

a)  $\frac{dV}{dt} = 25 \text{ cm}^3/\text{s}$

$V = \frac{4}{3}\pi r^3$       $A = 4\pi r^2$

$\frac{dV}{dr} = 4\pi r^2$       $\frac{dA}{dr} = 8\pi r$

$\frac{dV}{dr} = \frac{dV}{dt} \times \frac{dt}{dr}$

$\frac{dV}{dA} = \frac{dV}{dr} \times \frac{dr}{dA}$

$= 4\pi r^2 \times \frac{1}{8\pi r}$

$\therefore \frac{dV}{dA} = \frac{dV}{dr} \times \frac{dr}{dA} = \frac{dV}{dA}$

$= \frac{r}{2}$

$= \frac{r}{2} \times 25 = \frac{5}{2} \times 25 \text{ cm}^3/\text{s} = \frac{125}{2} \text{ cm}^3/\text{s}$

$= 62\frac{1}{2} \text{ cm}^3/\text{s}$

Quest 2

b) (i)  $\frac{dM}{dt} = 100 - Mk$

$\frac{dt}{dM} = \frac{1}{100 - Mk}$

$t = -\frac{1}{k} \ln(100 - M) + C$

$t = 0 \Rightarrow -\frac{1}{k} \ln(100 - M)$

$k(t - C) = \ln(100 - M)$

$e^{k(t-C)} = 100 - M$

let  $A = e^{kt}$

$100 - Ae^{kt} = M$

(ii)  $0 = 100 - A$

$A = 100$

(iii)  $40 = 100 - 100e^{-15t}$

$60 = 100e^{-15t}$

$\ln \frac{60}{100} = -15t$

$t = 30$

$M = 100 - 100e^{-15 \times 30}$

$= 64$

Quest 3

$$a) \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{4})}{3x} = \frac{-1}{3} \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{4})}{\frac{x}{4}}$$

$$= \frac{-1}{3} \lim_{x \rightarrow 0} \frac{\sin(\frac{x}{4})}{\frac{x}{4}} \checkmark$$

$$= \frac{-1}{3} \checkmark$$

b) (i)  $\cos x - \sin x \Rightarrow \cos x = 1 \quad \therefore R = \sqrt{2}$   
 $\sin x = 1$   
 $\therefore \cos x - \sin x = \sqrt{2} \cos(x + \frac{\pi}{4}) \checkmark$

(ii)  $\cos x - \sin x = \frac{\sqrt{2}}{2}$   
 $\sqrt{2} \cos(x + \frac{\pi}{4}) = \frac{\sqrt{2}}{2}$

$$\cos(x + \frac{\pi}{4}) = \frac{1}{2} \checkmark$$

$$-\frac{\pi}{4} \leq x + \frac{\pi}{4} \leq \frac{7\pi}{4}$$

$$x + \frac{\pi}{4} = \frac{\pi}{3}, \frac{5\pi}{3} \quad \therefore 0 \leq x + \frac{\pi}{4} \leq 2\pi$$

$$x = -\frac{\pi}{12}, \frac{17\pi}{12} \checkmark$$

c)  $\lim_{A \rightarrow 0} \frac{\tan A}{\tan 2A - \tan A} = \frac{\tan A}{2 \tan A - \tan A} - \tan A$   
 $= \frac{\tan A}{1 - \tan^2 A} - \tan A$   
 $= \frac{\tan A}{1 - \tan^2 A} - \frac{\tan A(1 - \tan^2 A)}{1 - \tan^2 A}$   
 $= \frac{1 - 1 + \tan^2 A}{1 - \tan^2 A} \checkmark$

Quest 4

a)  $\sin(105^\circ) = \sin(60 + 45)$   
 $= \sin 60 \cos 45 + \sin 45 \cos 60$   
 $= \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \times \frac{1}{2}$   
 $= \frac{\sqrt{3} + 1}{2\sqrt{2}}$   
 $= \frac{\sqrt{6} + \sqrt{2}}{4}$

b)  $\int_0^{\frac{\pi}{2}} \sin^2 3x \, dx$   
 $= \int_0^{\frac{\pi}{2}} \frac{1 - \cos 6x}{2} \, dx$   
 $= \frac{\pi}{2} \left[ x - \frac{\sin 6x}{6} \right]_0^{\frac{\pi}{2}}$   
 $= \frac{\pi^2}{6} \checkmark$

c)  $\int_0^{\frac{\pi}{2}} x \cos 3x \, dx = \frac{1}{3} \int_0^{\frac{\pi}{2}} 3x \cos 3x + \sin 3x - \sin 3x$   
 $= \frac{1}{3} \left[ x \sin 3x + \frac{1}{3} \cos 3x \right]_0^{\frac{\pi}{2}}$   
 $= \frac{1}{3} \left[ -\frac{\pi}{2} - \frac{1}{3} \right]$   
 $= -\left( \frac{\pi}{6} + \frac{1}{9} \right) \checkmark$

Question 5

a)  $y = \tan 2x$   
 $y' = 2 \sec^2 2x$

$\frac{dy}{dx} = 8$   
 $\frac{dy}{dx} = 2 \sec^2 2x$   
 $\therefore \frac{8}{2} = \sec^2 2x$   
 $4 = \sec^2 2x$   
 $\sec 2x = \pm 2$   
 $2x = \arccos(\pm \frac{1}{2})$   
 $x = \frac{1}{2} \arccos(\pm \frac{1}{2})$   
 $x = \frac{1}{2} (\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3})$   
 $x = \frac{\pi}{6}, \frac{\pi}{3}, \frac{2\pi}{3}, \frac{5\pi}{6}$

$y - \sqrt{3} = 8(x - \frac{\pi}{6})$   
 $8x - y - \frac{\pi}{6} + \sqrt{3} = 0$

b)  $\tan \theta = \frac{m_1 - m_2}{1 + m_1 m_2}$   
 $m_1 = -4$   
 $m_2 = -2$   
 $\tan \theta = \frac{-4 - (-2)}{1 + (-4)(-2)} = \frac{-2}{9}$

$\theta \approx 150^\circ 57' 12''$

c)  $\sin 2\theta + \cos \theta = 0$   
 $2 \sin \theta \cos \theta + \cos \theta = 0$   
 $\cos \theta (2 \sin \theta + 1) = 0$

$\cos \theta = 0$

$\theta = \frac{\pi}{2}, \frac{3\pi}{2}$

$\sin \theta = -\frac{1}{2}$

$\theta = \frac{7\pi}{6}, \frac{11\pi}{6}$

$\therefore \theta = \frac{\pi}{2}, \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6}$

Question 6

a)  $5 \sin \theta - 2 \cos \theta = 2$  let  $\tan \frac{\theta}{2} = t \therefore \sin \theta = \frac{2t}{1+t^2}$

$\frac{10t}{1+t^2} - \frac{2(1-t^2)}{1+t^2} = 2$

$10t - 2 + 2t^2 = 2 + 2t^2$

$\cos \theta = \frac{1-t^2}{1+t^2}$

Question 6

a) cont'd.

$\frac{2t^2 + 10t - 2}{1+t^2} = 2$

$2t^2 + 10t - 2 = 2 + 2t^2$   
 $2t^2 - 10t + 4 = 0$

$10t = 2$

$t = \frac{1}{5}$

$t = \frac{10 \pm \sqrt{100 - 32}}{4}$

$\therefore \tan \frac{\theta}{2} = \frac{1}{5} \quad 0 \leq \frac{\theta}{2} \leq \frac{\pi}{2}$

$\frac{\theta}{2} = 11^\circ 19'$

$\theta = 22^\circ 38'$

Also need to test for  $\theta = 180^\circ$

$5 \sin \theta - 2 \cos \theta = 2$

$5(0) - 2(-1) = 2$

is true


$\therefore \theta = 180^\circ$  is also a solution.

$\therefore \theta \approx 155^\circ 16'$

$47^\circ 21'$

$180^\circ$

b) (i)



(ii)  $x = 1 - 3 \cos(\frac{t}{2})$

$\dot{x} = \frac{3}{2} \sin(\frac{t}{2})$

particle is at rest at  $t = 2\pi$

$n = 4$

For  $x = 0, t = 2\pi$

Ques 6

b) (iii) max speed at  $\ddot{x} = 0$  i.e. at point of inflection  
at  $t = \pi$  or  $3\pi$ .

$$\dot{x} = \frac{3}{4} \cos\left(\frac{t}{2}\right) = 0$$

$$\therefore t = \pi \checkmark \text{ the speed is } \frac{3}{2} \text{ m/s.} \checkmark$$