

# C.E.M.TUITION

Student Name : \_\_\_\_\_

**Review Topic : Double Angle Formulae**

**(Preliminary - Paper 1)**

**Year 12 - 3 Unit**

1. Prove that  $\frac{1-\cos 2x}{1+\cos 2x} = \tan^2 x$  and hence show that the exact value of  $\tan\left(\frac{\pi}{8}\right)$  is  $\sqrt{2} - 1$ .

2. Express  $\sin x$  and  $\cos x$  in terms of  $t = \tan\left(\frac{x}{2}\right)$ . Hence prove

$$\text{that: } \frac{1 + \sin x - \cos x}{1 + \sin x + \cos x} = \tan\left(\frac{x}{2}\right)$$

3. If  $\cos x = \frac{7}{9}$  and  $\sin y = \frac{1}{3}$ , where angles  $x$  and  $y$  are acute,

- (a) Show that  $x = 2y$
- (b) Find the exact value of  $\tan(x + y)$

4. Given  $\sin x = \frac{1}{\sqrt{3}}$ ,  $\sin y = \frac{1}{\sqrt{2}}$ , and  $0 < x, y < \frac{\pi}{2}$ , find the exact value of  $\sin(x + y)$ .

5. Given that  $\frac{\cos(A - B)}{\cos(A + B)} = \frac{7}{3}$ , show that  $5\tan A = 2\cot B$ .

Further, given that  $\tan B = 2$ , and A is acute, find in exact terms:

- (a)  $\tan(A + B)$       (b)  $\sin A$       (c)  $\cos 2A$

1.  $\frac{1-\cos 2x}{1+\cos 2x}$

$$= \frac{1-(1-2\sin^2 x)}{1+2\cos^2 x-1}$$

$$= \sin^2 x + \cos^2 x$$

$$= \tan^2 x$$

Put  $x = \frac{\pi}{8}$ , then :

$$\tan^2\left(\frac{\pi}{8}\right)$$

$$= \left(1 - \frac{1}{\sqrt{2}}\right) + \left(1 + \frac{1}{\sqrt{2}}\right)$$

$$= (\sqrt{2}-1) + (\sqrt{2}+1)$$

$$= \frac{(\sqrt{2}-1)^2}{(\sqrt{2}-1)(\sqrt{2}+1)}$$

$$\therefore \tan\left(\frac{\pi}{8}\right) = \sqrt{2}-1$$

2.  $t = \tan\left(\frac{x}{2}\right), \sin x = \frac{2t}{1+t^2}$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$\frac{1+\sin x - \cos x}{1+\sin x + \cos x}$$

$$= \frac{1 + \frac{2t}{1+t^2} - \frac{1-t^2}{1+t^2}}{1 + \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2}}$$

$$= \frac{2t(1+t)}{2(1+t)}$$

$$= t, \text{ i.e. } \tan\left(\frac{x}{2}\right)$$

3.  $\cos x = \frac{7}{9}, \sin x = \frac{4\sqrt{2}}{9}$

$$\sin y = \frac{1}{3}, \cos y = \frac{2\sqrt{2}}{3}$$

(a)  $\sin 2y = 2 \sin y \cos y$

$$= \frac{4\sqrt{2}}{9}$$

$$\therefore \sin 2y = \sin x$$

$x = 2y$  ( $x$  is acute)

(b)  $\tan x = \frac{4\sqrt{2}}{7},$

$$\tan y = \frac{1}{2\sqrt{2}}$$

$$\tan(x+y) = \frac{\frac{4\sqrt{2}}{7} + \frac{1}{2\sqrt{2}}}{1 - \frac{4\sqrt{2}}{14\sqrt{2}}}$$

$$\therefore \tan(x+y) = \frac{23}{2\sqrt{2}}$$

4.  $x, y$  acute angles.

$$\sin x = \frac{1}{\sqrt{3}}, \cos x = \frac{\sqrt{2}}{\sqrt{3}}$$

$$\sin y = \frac{1}{\sqrt{2}}, \cos y = \frac{1}{\sqrt{2}}$$

$$\sin(x+y) = \sin x \cos y$$

$$+ \cos x \sin y$$

$$= \frac{1+\sqrt{2}}{\sqrt{6}}$$

5.  $\frac{\cos(A-B)}{\cos(A+B)} = \frac{7}{3}$

$$\frac{\cos A \cos B + \sin A \sin B}{\cos A \cos B - \sin A \sin B} = \frac{7}{3}$$

Dividing both numerator and denominator by  $\cos A \cos B$

$$\frac{1 + \tan A \tan B}{1 - \tan A \tan B} = \frac{7}{3}$$

$$3 + 3 \tan A \tan B$$

$$= 7 - 7 \tan A \tan B$$

$$5 \tan A \tan B = 2$$

$$\therefore 5 \tan A = 2 \cot B$$

Given  $\tan B = 2, \cot B = \frac{1}{2}$


then  $\tan A = \frac{1}{5}$

(a)  $\tan(A+B)$

$$= \left(\frac{1}{5} + 2\right) + \left(1 - \frac{2}{5}\right)$$

$$= \frac{11}{3}$$

(b)  $\sin A = \frac{1}{\sqrt{26}}$



(c)  $\cos 2A = 1 - 2 \sin^2 A$

$$= 1 - \frac{2}{26} \text{ or } \frac{12}{13}.$$