



SYDNEY BOYS HIGH
SCHOOL
MOORE PARK, SURRY HILLS

2006
YEAR 11
YEARLY EXAM

Mathematics Extension

General Instructions

- Reading Time – 5 Minutes
- Working time – 90 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- Each Section is to be returned in a separate bundle.
- Marks may NOT be awarded for messy or badly arranged work.
- All necessary working should be shown in every question.

Total Marks – 60

- Attempt questions 1-3
- Hand up in 3 sections clearly marked A,B & C

Examiner: *A.M.Gainford*

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, x > 0$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}, a \neq 0$$

$$\int \cos ax dx = \frac{1}{a}\sin ax, a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a}\cos ax, a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a}\tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a}\sec ax, a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a}\tan^{-1} \frac{x}{a}, a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, a > 0, -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, x > 0$

SECTION A

Question 1 (20 marks)	Marks
a) Find the remainder when the polynomial $P(x) = x^3 + 3x - 2$ is divided by $x - 3$.	1
b) If $\operatorname{cosec} \theta = \frac{2}{\sqrt{3}}$, find the possible values of $\tan \theta$.	2
c) The equation $x^2 - (1 - 2k)x + k + 3 = 0$ has consecutive integral roots. Find the possible values of k .	3
d) i) In how many ways can 6 committee members be selected from 10 people?	4
ii) In how many ways can this be done if two particular people will only serve together?	
e) Express $\sin 3\theta$ as an expression in powers of $\sin \theta$ only.	2
f) i) Write the equation of the tangent to the curve $y = x^3$ at the point on the curve where $x = 1$.	3
ii) Find the co-ordinates of the point where this tangent crosses the curve.	
g) Solve $ 2x + 6 < 4$	2
h) Solve for x : $\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 12 = 0$	3

SECTION B

Question 2 (20 marks)	Marks
Start a New Booklet	
a) Find the coordinates of the point which divides AB with $A(1,4)$ and $B(5,2)$ externally in the ratio 1:3.	2
b) Given the curve with equation $y = \frac{x}{x^2 + 1}$:	7
i) Find the first and second derivatives.	
ii) Identify and determine the nature of any turning points and points of inflexion.	
iii) Make a neat sketch of the curve.	3
c) Solve the inequality $\frac{4}{5-x} \geq 1$.	4
d) Find the general solution to the equation $\cos\left(\theta + \frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}$.	4
e) i) Express $\sin \theta + \sqrt{3} \cos \theta$ in the form $R \sin(\theta + \alpha)$.	
ii) Hence or otherwise sketch the graph of $y = \sin x + \sqrt{3} \cos x$ in the domain $0 \leq x \leq \pi$.	

SECTION C

Question 3 (20 marks)

Marks


Start a New Booklet

a) In the diagram $\angle\alpha > \angle\beta > 0$ and $\frac{\cos(\alpha + \beta)}{\cos(\alpha - \beta)} = \frac{4}{5}$

i) Show that $\tan\alpha \cdot \tan\beta = \frac{1}{9}$

ii) If $PR = QR$, show that $9(\tan\alpha + \tan\beta) = 8$

iii) Hence find α and β .



8

4

b) i) The polynomial equation $P(x) = 0$ has a double root at $x = 9$. By writing $P(x) = (x - a)^2 Q(x)$, where $Q(x)$ is a polynomial, show $P'(a) = 0$.

ii) Hence or otherwise find the values of a and b if $x = 1$ is a double root of $x^4 + ax^3 + bx^2 - 5x + 1 = 0$

c) Two points $P(2ap, ap^2)$ and $Q(2aq, aq^2)$ lie on the parabola $x^2 = 4ay$.

8

i) Derive the equation of the tangent to the parabola at P .

ii) Find the coordinates of the point of intersection T of the tangents to the parabola at P and Q .

iii) Given that the tangents at P and Q in (ii) intersect at an angle of 45° , show that $p - q = 1 + pq$.

iv) By evaluating the expression $x^2 = 4ay$ at T , or otherwise, find the locus of T .

SECTION A. (Q1).

(a) $R = P(3) = 27 + 9 - 2$
 $= 184$ [1]

(b) $\operatorname{cosec} \theta = \frac{2}{\sqrt{3}}$

$\therefore \sin \theta = \frac{\sqrt{3}}{2}$

$\theta = 60^\circ, 120^\circ \text{ etc.}$

$\therefore \tan \theta = \pm \sqrt{3}$ [2]

(c) Let the roots be α and $\alpha+1$.

Now $\alpha + (\alpha+1) = -\frac{b}{a} = 1-2k$ ie $2\alpha+1 = 1-2k$
 ie $\alpha = -k$.

And $\alpha(\alpha+1) = \frac{c}{a} = k+3$ ie $-k(-k+1) = k+3$
 $k^2 - k = k+3$

$k^2 - 2k - 3 = 0$

$(k+1)(k-3) = 0$

$k = 3, -1$ [3]

(d) (i) $\binom{10}{6} = 210$ [2]

(ii) If they (ie 2 particular people) will only ~~see~~ together, they are either both on or both off.

$\therefore \binom{8}{4} + \binom{8}{6} = 70 + 28 = 98$ [2]

(e) $\sin 3\theta = \sin(2\theta + \theta)$
 $= \sin 2\theta \cos \theta + \cos 2\theta \sin \theta$
 $= 2\sin \theta \cos \theta \cos \theta + (1 - 2\sin^2 \theta) \sin \theta$
 $= 2\sin \theta \cos^2 \theta + \sin \theta - 2\sin^3 \theta$
 $= 2\sin \theta (1 - \sin^2 \theta) + \sin \theta - 2\sin^3 \theta$
 $= 2\sin \theta - 2\sin^3 \theta + \sin \theta - 2\sin^3 \theta$

$\therefore \sin 3\theta = 3\sin \theta - 4\sin^3 \theta$ [2]

(f) (i) $y = x^3$
 $y' = 3x^2$ \therefore slope of tangent at $x=1$,
 is 3.

\therefore eqn. of tangent at $(1, 1)$

is $y-1 = 3(x-1)$

$y-1 = 3x-3$

$y = 3x-2$ [1 1/2]

(ii) Solving $y = x^3$ and $y = 3x-2$.

ie $x^3 = 3x-2$

$x^3 - 3x + 2 = 0$ — (A)

Now (A) will have roots 1, 1 and α

Now $1+1+\alpha = -\frac{b}{a} = 0$.

$\therefore \alpha = -2$.

\therefore the other point is $(-2, -8)$ [1 1/2]

$$(g) |2x+6| < 4$$

$$\begin{aligned} \therefore -4 < 2x+6 < 4 \\ -10 < 2x < -2 \\ \underline{-5 < x < -1} \end{aligned}$$

2

$$(h) \text{ let } x + \frac{1}{x} = u.$$

\therefore equation becomes

$$\begin{aligned} u^2 - u - 12 &= 0 \\ (u-4)(u+3) &= 0 \\ u &= 4, -3 \end{aligned}$$

$$\text{If } u=4$$

$$\begin{aligned} x + \frac{1}{x} &= 4 \\ x^2 + 1 &= 4x \\ x^2 - 4x + 1 &= 0 \\ x &= \frac{4 \pm \sqrt{16-4}}{2} \\ &= \frac{4 \pm \sqrt{12}}{2} \\ &= \frac{4 \pm 2\sqrt{3}}{2} \\ \underline{x = 2 \pm \sqrt{3}} \end{aligned}$$

$$\text{If } u=-3$$

$$\begin{aligned} x + \frac{1}{x} &= -3 \\ x^2 + 1 &= -3x \\ x^2 + 3x + 1 &= 0 \\ x &= \frac{-3 \pm \sqrt{9-4}}{2} \\ \underline{x = \frac{-3 \pm \sqrt{5}}{2}} \end{aligned}$$

3

YR11 Sept 2006 ext 1 paper

/20

Section B

$$\text{Q2. (a) } A(x_1, y_1) \quad B(x_2, y_2) \quad -1:3.$$

$$\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right)$$

$$\Rightarrow \left(\frac{-1 \times 5 + 3 \times 1}{-1+3}, \frac{-1 \times 2 + 3 \times 4}{-1+3} \right) = (-1, 5) \quad \text{②}$$

$$(b) y = \frac{x}{x^2+1}$$

$$(i) y' = \frac{(x^2+1) \times 1 - x \times 2x}{(x^2+1)^2}$$

$$= \frac{x^2+1-2x^2}{(x^2+1)^2} = \frac{1-x^2}{(x^2+1)^2} \quad \text{①/2}$$

$$y'' = \frac{(x^2+1)^2 \times -2x - (1-x^2) \times 2(x^2+1) \times 2x}{(x^2+1)^4}$$

$$= \frac{-2x(x^2+1)^2 - 4x(1-x^2)(x^2+1)}{(x^2+1)^4}$$

$$= \frac{-2x(x^2+1) - 4x(1-x^2)}{(x^2+1)^3}$$

$$= \frac{-2x^3 - 2x - 4x + 4x^3}{(x^2+1)^3} = \frac{2x^3 - 6x}{(x^2+1)^3}$$

$$= \frac{2x(x^2-3)}{(x^2+1)^3} \quad \text{①/2}$$

(b) (ii) Stat pts exist when $y' = 0$
 Since $(x^2+1)^2 \neq 0$, $1-x^2 = 0$

$$x^2 = 1$$

$$x = \pm 1$$

When $x = +1$, $y = \frac{1}{2}$ $(1, \frac{1}{2})$

When $x = -1$, $y = \frac{-1}{2}$ $(-1, -\frac{1}{2})$

At $(1, \frac{1}{2})$ $y'' = \frac{2(1-3)}{8+} = \frac{-2}{4} < 0$ MAX. stat pt. ①

At $(-1, -\frac{1}{2})$ $y'' = \frac{-2(1-3)}{8+} = \frac{2}{4} > 0$ min stat pt. ①

Inflexions occur when $y'' = 0$ and \exists a sign change
 $2x(x^2-3) = 0$ because $(x^2+1)^3 \neq 0$

$$x = 0, x^2 - 3 = 0$$

$$x^2 = 3$$

$$x = \pm \sqrt{3}$$

At $x = 0$, $y = \frac{0}{0+1} = 0 \Rightarrow (0, 0)$ ①

At $x = +\sqrt{3}$, $y = \frac{\sqrt{3}}{4} \Rightarrow (\sqrt{3}, \frac{\sqrt{3}}{4})$ ①


At $x = -\sqrt{3}$, $y = \frac{-\sqrt{3}}{4} \Rightarrow (-\sqrt{3}, -\frac{\sqrt{3}}{4})$ ①

At $x = 0 - \epsilon$ $y'' > 0$
 At $x = 0 + \epsilon$ $y'' < 0$ } sign change

At $x = \sqrt{3} - \epsilon$ $y'' < 0$
 At $x = \sqrt{3} + \epsilon$ $y'' > 0$ } sign change

At $x = -\sqrt{3} - \epsilon$ $y'' < 0$
 At $x = -\sqrt{3} + \epsilon$ $y'' > 0$ } sign change

b) (iii)



$(1, \frac{1}{2})$ MAX

$(0, 0)$ infl.

$(-1, -\frac{1}{2})$ MIN

$(\sqrt{3}, \frac{\sqrt{3}}{4}) \doteq (1.7, .4)$ infl

$(-\sqrt{3}, -\frac{\sqrt{3}}{4}) \doteq (-1.7, -.4)$ infl

(c) $\frac{4}{5-x} \geq 1$ now $x \neq 5$.

$$\frac{4(5-x)^2}{(5-x)^2} \geq 1 \cdot (5-x)^2$$


$$4(5-x) \geq (5-x)^2$$

$$4(5-x) - (5-x)^2 \geq 0$$

$$(5-x)[4 - (5-x)] \geq 0$$

$$(5-x)[-1+x] \geq 0$$

$$\text{So } (x-1)(5-x) \geq 0$$



above or touching $1 \leq x \leq 5$

but $x \neq 5$, so $\{x: 1 \leq x < 5\}$ // ③

2 (d) in radians

(4)

If $\cos \theta = \cos \alpha$

$\theta = 2\pi \times n \pm \alpha$

So $\cos(\theta + \frac{\pi}{6}) = \frac{\sqrt{3}}{2}$

$\theta + \frac{\pi}{6} = 2\pi \times n \pm \frac{\pi}{6}$

$\theta = 2\pi n \pm \frac{\pi}{6} - \frac{\pi}{6}$

So $\theta = 2\pi n$, $\theta = 2\pi n - \frac{\pi}{3}$

(4)

2 (e) (i) $\sin \theta + \sqrt{3} \cos \theta = 2(\frac{1}{2} \sin \theta + \frac{\sqrt{3}}{2} \cos \theta)$ $R = \sqrt{1 + \sqrt{3}^2} = 2$

no $2(\frac{1}{2} \sin \theta + \frac{\sqrt{3}}{2} \cos \theta) = R(\sin \theta \cos \alpha + \cos \theta \sin \alpha)$

now $R=2$, $\cos \alpha = \frac{1}{2}$ quad 1, 4 } quad 1.
 $\sin \alpha = \frac{\sqrt{3}}{2}$ quad 1, 2 }

So $\alpha = \frac{\pi}{3}$ (radian measure)


So $2 \sin(\theta + \frac{\pi}{3}) = \sin \theta + \sqrt{3} \cos \theta$

(3)

(ii)

$y = 2 \sin(\theta + \frac{\pi}{3})$

$\theta = 0, y = 2 \sin \frac{\pi}{3} = \sqrt{3}$
 $\theta = \frac{\pi}{2}, y = 2 \sin \frac{5\pi}{6} = 1$
 $\theta = \pi, y = 2 \sin \frac{4\pi}{3} = -\sqrt{3}$



Question [3] Solution

$\frac{\cos(\alpha+\beta)}{\cos(\alpha-\beta)} = \frac{4}{5}$

(8)

$5[\cos \alpha \cos \beta - \sin \alpha \sin \beta] = 4[\cos \alpha \cos \beta + \sin \alpha \sin \beta]$

$\cos \alpha \cos \beta = 9 \sin \alpha \sin \beta$ [2]

$\therefore \tan \alpha \tan \beta = \frac{1}{9}$ — (1)

If $P R = Q R \Rightarrow \alpha + \beta = 45^\circ$

$\therefore \tan(\alpha + \beta) = 1$ [1]

Now, $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = 1$

$\therefore \tan \alpha + \tan \beta = 1 - \tan \alpha \tan \beta = 1 - \frac{1}{9} = \frac{8}{9}$ [2]

$\Rightarrow q(\tan \alpha + \tan \beta) = 8$ — (2)

Let $\tan \alpha, \tan \beta$ be the roots to $x^2 - \frac{8}{q}x + \frac{1}{q} = 0$ [1]

$\therefore 9x^2 - 8x + 1 = 0$
 $x = \frac{8 \pm \sqrt{64 - 36}}{18}$ [2]
 $\therefore \tan \alpha = \frac{8 + 2\sqrt{7}}{18}$ $\alpha = 36^\circ 27'$
 $\therefore \tan \beta = \frac{8 - 2\sqrt{7}}{18}$ $\beta = 8^\circ 33'$

(b) $P(x) = (x-9)^2 Q(x)$ (4)

(i) $P'(x) = 2Q(x)(x-9) + (x-9)^2 Q'(x)$
 $\therefore P'(9) = 0$ [2]

(ii) $P(1) = 0 \Rightarrow 1 + a + b - 4 = 0$
 $\therefore a + b = 3$ — (1) [2]

$P'(1) = 0 \Rightarrow 4 + 3a + 2b - 5 = 0$


i.e. $3a + 2b = 5$ — (2)

(1) $x = -2 \Rightarrow -2a + 2b = -6$ — (3)

$a = -1$
$b = 4$

(c)

(8)



$\frac{dy}{dx} = \frac{2x}{4a} = \frac{x}{2a}$

$\frac{dy}{dx} \Big|_{x=2ap} = \frac{2ap}{2a} = p$

(i) $y - ap^2 = p(x - 2ap)$ [1]

$y = px - ap^2$ — (1)

Equation of QT is

(ii) $y = qx - aq^2$ — (2)

Solve (1) & (2) [2]

$(p-q)x = a(p+q)(p-q)$

$\therefore x = a(p+q)$
 $y = apq$ — (4)

$\tan \alpha = 1$
 $\therefore \tan \alpha = \frac{p-q}{1+pq}$ [2]

$\therefore p-q = (1+pq) \tan \alpha$ — (5)

$\frac{x}{a} = p+q$
 $\frac{y}{a} = pq$ [3]

$(p-q)^2 = (p+q)^2 - 4pq$

$\therefore (1+pq)^2 = (p+q)^2 - 4pq$

$\therefore (1 + \frac{y}{a})^2 = \frac{x^2}{a^2} - \frac{4y}{a}$

$1 + \frac{2y}{a} + \frac{y^2}{a^2} = \frac{x^2}{a^2} - \frac{4y}{a}$

$a^2 + 2ay + y^2 = x^2 - 4ay$ (6)

Locus of T

$x^2 - y^2 - 6ay - a^2 = 0$