



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS

SEPTEMBER 2007

Yearly Examination

YEAR 11

Mathematics Extension (Continuers)

General Instructions

- Reading Time – 5 Minutes.
- Working time – 60 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for messy or badly arranged work.
- Start each NEW section in a separate answer booklet.

Total Marks – 60

- Attempt questions 1 – 4

Examiner: R. Boros

Total marks 60

Attempt questions 1 to 4

Answer each Section in a Separate writing booklet

Section A (Use a SEPARATE writing booklet)

Question 1 (14 marks)

- (a) Find the acute angle between the lines $y = 2x + 1$ and $y = -x + 1$, correct to the nearest minute. 2
- (b) Consider the polynomial $K(x) = 4x^3 + tx^2 + 2x - 1$. Given that $x + 1$ is a factor of $K(x)$, find the value of t . 2
- (c) The parametric equations of a curve are $x = \frac{2}{t}$ and $y = 2t^2$. What is the cartesian equation for the curve? 2
- (d) For the parabola $(x - 3)^2 = 6y + 12$, find the:
- i. coordinates of the vertex 1
 - ii. coordinates of the focus 1
 - iii. equation of the directrix 1
- (e) Find the coordinates of the point Q which divides the interval joining $A(2, -3)$ and $B(-4, 1)$ externally in the ratio $1 : 3$. 3
- (f) Sketch the graph of $y = x^2(x - 2)^3$ without the use of calculus. 2

Question 2 (14 marks)

- (a) Differentiate $f(x) = 5 - x^2$ by using first principles. 2
- (b) Given that $\sin \alpha = \frac{7}{25}$ and $\cos \beta = -\frac{3}{5}$ where α and β are obtuse angles, find the exact value of:
- i. $\sin 2\alpha$ 1
- ii. $\cos(\alpha + \beta)$ 2
- (c) In a class of 30 students, 22 study Chemistry, 18 study Physics and 13 study both Chemistry and Physics. If a student is chosen at random, what is the probability that the student studies Chemistry or Physics? 2
- (d) i. Express $\sin x - \sqrt{3} \cos x$ in the form $A \sin(x - \alpha)$, with $A > 0$ and $0 < \alpha < \frac{\pi}{2}$. 2
- ii. Find the solutions to $\sin x - \sqrt{3} \cos x = \frac{2}{\sqrt{2}}$ for $0 \leq x \leq 2\pi$. 2
- (e) Solve the inequation $\frac{1}{x+2} \leq \frac{1}{x+3}$. 3

Section B (Use a SEPARATE writing booklet)

Question 3 (15 marks)

- (a) Solve $\log_{27} 16 = x \log_3 2$. 2
- (b) Prove the trigonometric identity $\frac{\cos 2x}{(\cos x + \sin x)^3} = \frac{\cos x - \sin x}{1 + \sin 2x}$. 3
- (c) A committee of three is to be chosen from a group of four males and five females. The committee must include at least one male and at least one female. How many different committees can be formed? 2
- (d) A maths teacher pays \$1000 into a superannuation fund at the beginning of each year. Compound interest is paid at 9%p.a. on the investment.
- i. Show that the first \$1000 invested becomes \$20413.97 to the nearest cent after 35 years. 1
- ii. What will be the value of the investment at the end of 35 years? Answer correct to the nearest dollar. 3
- (e) Given that the cubic equation $2x^3 + 6x - 1 = 0$ has real roots α , β and γ . Evaluate:
- i. $\alpha^3 \beta^3 \gamma^3 + \alpha^3 \beta^2 \gamma^3 + \alpha^2 \beta^3 \gamma^3$. 2
- ii. $\frac{\alpha}{\beta\gamma} + \frac{\beta}{\alpha\gamma} + \frac{\gamma}{\alpha\beta}$. 2

Question 4 (17 marks)

(a) The derivative of $x\sqrt{x^2+3}$ is $\frac{ax^2+b}{\sqrt{x^2+3}}$, where a and b are constants. Find the value of a and b .

3

(b) The student council at a local school consists of 4 boys and 2 girls. In how many ways can they sit next to each other around a circular table for a meeting if:

- there are no restrictions.
- the girls are not to sit next to each other.


1

2

(c) Find the general solution of the equation $\tan 2\theta = \tan \theta$ in radians.

3

(d)



The point $A(3at, -at^2)$ is a variable point on the parabola $x^2 = -9ay$. The normal at A meets the line $x = -at$ at the point B .

i. Show that the equation of the normal to the parabola at A is


2

$$3x - 2ty = 2at^3 + 9at.$$

ii. Find the coordinates of B .

1

(e)



A building AB of height $2h$ metres has a flag pole of height h metres on top of it. From a point C , due south of the building, the angle of elevation of the top of the building is 40° . From a point D , due west of the building, the angle of elevation of the top of the flagpole is 50° . The points C and D are on the same level as A and they are 40 metres apart.

i. Find expressions for AC and AD in terms of h .

2

ii. Show that $h = \frac{40}{\sqrt{4 \cot^2 40^\circ + 9 \cot^2 50^\circ}}$.

2

iii. Find to the nearest degree, the true bearing of D from C .

1

End of paper

Section A

QUESTION 1

$$(a) \tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$= \left| \frac{2 - (-1)}{1 + (2)(-1)} \right|$$

$$= 3.$$

$$\tan \alpha = 3$$

$$\alpha = 71^\circ 34'$$

$$(b) K(-1) = -4 + t - 2 - 1; = 0.$$

$$t = 7.$$

$$(c) x = \frac{t}{2} \Rightarrow t = \frac{2}{x}$$

So

$$y = 2 \left(\frac{2}{x} \right)^2$$

$$y = \frac{8}{x^2}$$

$$(d) (x-3)^2 = 4 \left(\frac{3}{2} \right) (y+2).$$

(i) Vertex $(3, -2)$.

(ii) Focus $(3, -\frac{1}{2})$.

(iii) Directrix $y = -\frac{7}{2}$ or $-3\frac{1}{2}$


$$(e) \left(\frac{mx + my_2}{m+n}, \frac{ny_1 + my_2}{m+n} \right)$$

External division $m:1$

$$\left(\frac{-3 \times 2 + 1 \times -4}{1-3}, \frac{-3 \times -3 + 1 \times 1}{1-3} \right)$$

$$= \left(\frac{-6-4}{-2}, \frac{9+1}{-2} \right)$$

$$= (-1, -5)$$



QUESTION 2.

$$(a) f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{5 - (x+h)^2 - (5 - x^2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{5 - x^2 - 2xh - h^2 - 5 + x^2}{h}$$

$$= \lim_{h \rightarrow 0} -2x + h$$

$$= -2x$$

$$(b) \sin \alpha = \frac{7}{25} \Rightarrow \cos \alpha = -\frac{24}{25}$$

$$\cos \beta = -\frac{3}{5} \Rightarrow \sin \beta = \frac{4}{5}$$

$$(i) \sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$= 2 \times \frac{7}{25} \times -\frac{24}{25}$$


$$= -\frac{336}{625}$$

$$(ii) \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$= -\frac{24}{25} \times -\frac{3}{5} - \frac{7}{25} \times \frac{4}{5}$$

$$= \frac{44}{125}$$

(c)



$$\frac{27}{30} = \frac{9}{10}$$

$$(d)(i) A \sin(x - \alpha) = A \cos \alpha \sin x - A \sin \alpha \cos x$$

$$\text{So } A \cos \alpha = 1$$

$$A \sin \alpha = \sqrt{3}$$

$$\text{Thus } \tan \alpha = \sqrt{3} \quad \text{and} \quad A^2 \cos^2 \alpha + A^2 \sin^2 \alpha = 1 + 3$$

$$A^2 = 4$$

$$A = 2$$

$$\sin x - \sqrt{3} \cos x = 2 \sin(x - \frac{\pi}{3})$$

$$(ii) 2 \sin(x - \frac{\pi}{3}) = \frac{2}{\sqrt{2}}$$

$$\sin(x - \frac{\pi}{3}) = \frac{1}{\sqrt{2}}$$

$$x - \frac{\pi}{3} = \frac{\pi}{4}, \frac{3\pi}{4}$$

$$x = \frac{7\pi}{12}, \frac{13\pi}{12}$$


$$(e) \frac{1}{x+2} - \frac{1}{x+3} \leq 0$$

$$\frac{(x+3) - (x+2)}{(x+3)(x+2)} \leq 0$$

$$\frac{1}{(x+3)(x+2)} \leq 0$$

$$(x+3)(x+2) \leq 0$$

$$-3 \leq x \leq -2$$



B:1) 2007 Yr II Yearly - Continuers SECTION B

QUESTION 3

a) $\log_{27} 16 = x \log_3 2$

$$\begin{aligned} \text{LHS} &= \log_{27} 16 \\ &= \frac{\log_3 16}{\log_3 27} \\ &= \frac{\log_3 2^4}{3} \\ &= \frac{4}{3} \log_3 2 \end{aligned}$$

$\therefore x = \frac{4}{3}$

b) $\frac{\cos 2x}{(\cos x + \sin x)^3} = \frac{\cos x - \sin x}{1 + \sin 2x}$

$$\begin{aligned} \text{LHS} &= \frac{\cos 2x}{(\cos x + \sin x)^3} \\ &= \frac{\cos^2 x - \sin^2 x}{(\cos x + \sin x)^3} \\ &= \frac{(\cos x + \sin x)(\cos x - \sin x)}{(\cos x + \sin x)^3} \\ &= \frac{\cos x - \sin x}{(\cos x + \sin x)^2} \\ &= \frac{\cos x - \sin x}{\cos^2 x + 2\sin x \cos x + \sin^2 x} \end{aligned}$$

b) continued.

$$\begin{aligned} &= \frac{\cos x - \sin x}{1 + \sin 2x} \\ &= \text{RHS} \end{aligned}$$

c) 2 women 1 man = ${}^5C_2 \times {}^4C_1$
 + 2 men 1 woman = ${}^4C_2 \times {}^5C_1$
 $= 10 \times 4 + 6 \times 5$
 $= \underline{\underline{70 \text{ ways}}}$

d)

i) $1000(1.09)^{35} = \underline{\underline{20413.97}}$

ii) 1st yr = $1000(1.09)$
 2nd year = $1000(1.09) + (1000)(1.09)^2$
 3rd yr = $1000(1.09) + 1000(1.09)^2 + 1000(1.09)^3$
 35^{th} yr = $1000(1.09 + 1.09^2 + \dots + 1.09^{35})$
 $= 1000 \left(\frac{a(r^n - 1)}{r - 1} \right)$ $a = 1.09$
 $= 1000 \left(\frac{1.09(1.09^{35} - 1)}{1.09 - 1} \right)$ $r = 1.09$
 $= \underline{\underline{\$235,124.72}}$ $n = 35$

B:2) 2007 Yr II Yearly - Continuers: SECTION B.

QUESTION 3 (CONTINUED)

e) ii) CONTINUED

e) $2x^3 + 0x^2 + 6x - 1 = 0$

$= 0 + 3$

i) $\alpha^3 \beta^3 \gamma^3 + \alpha^3 \beta^2 \gamma^3 + \alpha^2 \beta^3 \gamma^3$
 $= \alpha^2 \beta^2 \gamma^2 (\alpha\beta + \alpha\gamma + \beta\gamma)$

$\frac{1}{2}$

$= (\alpha\beta\gamma)^2 (\alpha\beta + \alpha\gamma + \beta\gamma)$

$= \underline{\underline{-6}}$

$\alpha\beta\gamma = \frac{-d}{a} = \frac{1}{2}$

$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a} = \frac{6}{2} = 3$

$\therefore (\alpha\beta\gamma)^2 (\alpha\beta + \alpha\gamma + \beta\gamma) = \left(\frac{1}{2}\right)^2 \times 3$
 $= \frac{3}{4}$

ii) $\frac{\alpha}{\beta\gamma} + \frac{\beta}{\alpha\gamma} + \frac{\gamma}{\alpha\beta}$

$= \frac{\alpha^2 + \beta^2 + \gamma^2}{\alpha\beta\gamma}$

$= \frac{(\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \alpha\gamma)}{\alpha\beta\gamma}$

$\Rightarrow \alpha + \beta + \gamma = \frac{-b}{a} = 0$

$\alpha\beta + \beta\gamma + \alpha\gamma = \frac{c}{a} = 3$

$\alpha\beta\gamma = \frac{-d}{a} = \frac{1}{2}$

B:3 2007 Yr11 - Yearly - Continuers - SECTION B.

QUESTION 4.

Let $y = x\sqrt{x^2+3}$.

$$\frac{dy}{dx} = vu' + uv'$$

where $u = x$ $u' = 1$.

$$v = (x^2+3)^{\frac{1}{2}} \quad v' = \frac{1}{2}(2x)(x^2+3)^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = (x^2+3)^{\frac{1}{2}}(1) + (x)(x)(x^2+3)^{-\frac{1}{2}}$$

$$= \sqrt{x^2+3} + \frac{x^2}{\sqrt{x^2+3}}$$

$$= \frac{x^2+3+x^2}{\sqrt{x^2+3}}$$

$$= \frac{2x^2+3}{\sqrt{x^2+3}}$$

$$= \frac{ax^2+b}{\sqrt{x^2+3}}$$

$a=2$ $b=3$.

D) No of ways around a table
 $= (n-1)!$
 $= 5!$
 $= 120$ ways

b) (continued)

ii) $\underline{1} \quad \underline{^4C_1} \quad \underline{^3C_1} \quad \underline{^2C_1} \quad \underline{^1C_1} \quad \underline{1}$

↑
fix.

$$= 1 \times 4 \times 3 \times 3 \times 2 \times 1$$

$$= \underline{72 \text{ ways}}$$

© $\tan 2\theta = \tan \theta$.

$$\frac{2 \tan \theta}{1 - \tan^2 \theta} = \tan \theta$$

$$2 \tan \theta = \tan \theta - \tan^3 \theta$$

$$\tan^3 \theta + \tan \theta = 0$$

$$\tan \theta (\tan^2 \theta + 1) = 0$$

either:

$\tan^2 \theta + 1 = 0$ which can't happen so

$$\tan \theta = 0$$

thus this happens when

$$\sin \theta = 0$$

∴ General solution is.

$$\theta = n\pi \quad n \in \mathbb{Z}$$

B:4 2007 Yr11 - Yearly - Continuers - SECTION B.

QUESTION 4 (continued)

d) ii)

① i) $x = -at$ $A(3at, -at^2)$

$$x^2 = -9ay$$

$$y =$$

gradient of tangent

$$\frac{dy}{dx} = \frac{2x}{-9a} = m$$

gradient of normal is.

$$m_2 = \frac{1}{m_1}$$

$$= \frac{9a}{2x} \quad @ (3at, -at^2)$$

Point gradient formula.

$$y - y_1 = m(x - x_1)$$

$$y + at^2 = \frac{9a}{2(3at)} (x - 3at)$$

$$baty + 6a^2t^3 = 9ax - 27a^2t$$

$$6ty + 6at^3 = 9x - 27at$$

$$2ty + 2at^3 = 3x - 9at$$

$$\underline{3x - 2ty = 2at^3 + 9at}$$

$$3x - 2ty = 2at^3 + 9at$$

$$\text{at } x = -at$$

$$-3(at) - 2ty = 2at^3 + 9at$$

$$-2ty = 2at^3 + 9at + 3at$$

$$y = \frac{2at^3 + 12at}{-2t}$$

$$y = -at^2 - 6a$$

Co-ordinates of B

$$\underline{(-at, -at^2 - 6a)}$$

④ e) i) $AC \Rightarrow$

$$\tan 40 = \frac{2h}{AC}$$

$$AC = \frac{2h}{\tan 40}$$

$AD \Rightarrow$

$$\tan 50 = \frac{2h}{AD}$$

$$AD = \frac{2h}{\tan 50}$$

$$\underline{\underline{\tan 50}}$$

3.5) 2007 Yr11 Yearly - Continuers - SECTION B.

QUESTION 4 (CONTINUED)

e) continued.

ii). By Pythagoras.

$$40^2 = \left(\frac{2h}{\tan 40}\right)^2 + \left(\frac{3h}{\tan 50}\right)^2$$

$$40^2 = \frac{4h^2}{\tan^2 40} + \frac{9h^2}{\tan^2 50}$$

$$40^2 = h^2 (4 \cot^2 40 + 9 \cot^2 50)$$

$$h^2 = \frac{40^2}{4 \cot^2 40 + 9 \cot^2 50}$$


$$h = \frac{40}{\sqrt{4 \cot^2 40 + 9 \cot^2 50}}$$

$$\tan \theta = 1.0561.$$

$$\theta = 46.56.$$

∴ Bearing of D from C
is $360^\circ - 46.56^\circ$

$$= \underline{313^\circ \text{ (nearest degree)}}$$



$$\begin{aligned} \tan \theta &= \frac{3h}{\tan 50} \cdot \frac{2h}{\tan 40} \\ &= \frac{3h}{\tan 50} \times \frac{\tan 40}{2h} \\ &= \frac{3}{2} \frac{\tan 40}{\tan 50} \end{aligned}$$