



SYDNEY BOYS HIGH
SCHOOL
MOORE PARK, SURRY HILLS

2003
HIGHER SCHOOL CERTIFICATE
ASSESSMENT TASK # 3

Mathematics Extension 1

General Instructions

- Reading time – 5 minutes.
- Working time – 2 hours.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for messy or badly arranged work.
- Hand in your answer booklets in 3 sections. Section A (Questions 1 and 2), Section B (Questions 3 and 4) and Section C (Questions 5 and 6).
- Start each **NEW** section in a separate answer booklet.

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

Total Marks - 84 Marks

- All questions are of equal value.

Examiner: *B. Opferkuch*

QUESTION 1.

Use a *separate* Writing Booklet

Marks

- (a) Find a primitive of $\frac{1}{\sqrt{1-x^2}}$. 1
- (b) Evaluate $\int_0^{\ln 5} e^{-x} dx$ 2
- (c) Find $\lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{h}$ 1
- (d) A box contains 5 red marbles and 4 white marbles. Three marbles are drawn in succession without replacement, What is the probability that any two marbles are red and one is white? 2
- (e) The sector OAB of a circle centre O and radius r has an area of $\frac{3\pi}{4} \text{ cm}^2$. If the arc AB subtends an angle of $\frac{\pi}{6}$ at O , find the length of the arc AB . 2
- (f) If $\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \tan^{-1}(1) = k\pi$ (where k is rational) Find the value of k . 2
- (g) Find $\int \frac{\ln 3x}{x} dx$, using the substitution $u = \ln 3x$. 4

QUESTION 2.

Use a *separate* Writing Booklet.

Marks

(a) Differentiate the following with respect to x .

8

(i) $x \cos x$

(ii) $\log_e(\cos 5x)$

(iii) $\tan^{-1}\left(\frac{x}{3}\right)$

(iv) $e^{\tan x}$

(v) $\{(1 + \cos^{-1}(3x))\}^3$

(b) Find a primitive of:

6

(i) $x^2 e^{2x^3+1}$

(ii) $\frac{5}{4+x^2}$

(iii) $\frac{x^2}{x^3+7}$

(iv) $-\sin(\pi - x)$

QUESTION 3.

Use a *separate* Writing Booklet.

Marks

(a) In choosing three letters from the word PROBING, and assuming each choice is equally likely, what is the probability of choosing just one vowel? 2

(b) (i) In how many ways can the numbers 1, 2, 3, 4, 5, 6 be arranged around a circle? 3

(ii) How many of these arrangements have at least two even numbers together?

(c) (i) Show that the function $f(x) = x^2 + e^{-\frac{1}{2}x} - 5$ has a root between -2 and -1 . 4(ii) Taking $x = -2$ as a first approximation, apply Newton's method once, to show that the root of $f(x) = 0$ is approximately $-\frac{18}{e+8}$.(d) (i) Show that $\frac{1+x}{1-x} = -1 + \frac{2}{1-x}$. 3(ii) Hence find $\int \frac{1+x}{1-x} dx$ (e) Write down the general solution of $\sqrt{3} \tan \theta - 1 = 0$. Leave answer in exact radian form. 2

QUESTION 4.

Use a *separate* Writing Booklet

Marks

- (a) Consider the curves $y = \sin x$ and $y = \cos 2x$. 5
- (i) Sketch the graphs of the two curves on the same axes, in the domain $-\frac{\pi}{2} \leq x \leq \frac{\pi}{6}$.
- (ii) Show that the curves intersect at $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{6}$.
- (iii) Hence find the exact area bounded by the two curves.

- (b) Consider the function $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$. 3

- (i) Evaluate $f(2)$.
- (ii) State the domain and range of $y = f(x)$.
- (iii) Sketch the graph of $y = f(x)$.

- (c) Find the exact value of $\tan\left[2 \tan^{-1}\left(-\frac{1}{2}\right)\right]$. 3

- (d) The function $g(x)$ is given by $g(x) = \cos^{-1} x + \sin^{-1} x$, $0 \leq x \leq 1$. 3
- (i) Find $g'(x)$.
- (ii) Sketch the graph of $y = g(x)$.

QUESTION 5.

Use a *separate* Writing Booklet

Marks

- (a) Consider the function $f(x) = x \sin^{-1}(x^2)$. 8
- (i) State the domain and range of $f(x)$.
- (ii) Find $f'(x)$.
- (iii) Show that there is a horizontal inflexion at the origin.
- (iv) What is the slope of the tangent at $x = 1$.
- (v) Sketch the curve which represents $f(x)$.

- (b) (i) Show that $\int \frac{x}{\sqrt{x^2 + 2x - 3}} dx = \int \frac{x}{\sqrt{(x+1)^2 - 4}} dx$. 3

- (ii) Using the substitution $2 \sec u = x + 1$, find $\int \frac{x}{\sqrt{x^2 + 2x - 3}} dx$

- (c) The curve $y = \frac{1}{\sqrt{1+x^2}}$ is rotated about the x -axis. 3

Find the volume of the solid enclosed between $x = \frac{1}{\sqrt{3}}$ and $x = \sqrt{3}$.

QUESTION 6.

Use a separate Writing Booklet

Marks

(a) Evaluate $\int_0^{\sqrt{2}} x \sqrt[3]{x^2+1} dx$ using the substitution $u = x^2 + 1$.

3

Answer in exact form.

(b) Consider the equation $\cos 3x = \sin x$

3

(i) Show that $\cos\left(\frac{\pi}{2} - A\right) = \sin A$

(ii) Hence, or otherwise, find the general solution of the equation $\cos 3x = \sin x$.

(c) (i) Show that $\frac{d}{dx} \tan^3 x = 3 \sec^4 x - 3 \sec^2 x$.

4

(ii) Using (i) or otherwise, evaluate $\int_0^{\frac{\pi}{4}} \sec^4 x dx$

(d) Show that the exact value of $\int_0^{\frac{\pi}{6}} \sin^2 x dx = \frac{2\pi - 3\sqrt{3}}{24}$

4

THIS IS THE END OF THE PAPER

2003 Ext 1 #3

(1) (a) $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$

(b) $\int_0^{\ln 5} e^{-x} dx = -e^{-x} \Big|_0^{\ln 5} = -[e^{-\ln 5} - e^0] = -[e^{-\frac{\ln 5}{1}} - 1] = -[e^{-\frac{1}{5}} - 1] = 1 - \frac{1}{5} = \frac{4}{5}$

$e^{\ln x} = x$

(c) $\lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}} = \frac{1}{2} \lim_{h \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}} = \frac{1}{2} \times 1 = \frac{1}{2}$

$\left[\lim_{u \rightarrow 0} \frac{\sin u}{u} = 1 \right]$

(d) 5R 4W


method 1:

$\frac{\binom{5}{2} \times \binom{4}{1}}{\binom{9}{3}} = \frac{10}{21}$

method 2: RRW

$\left(\frac{5}{9} \times \frac{4}{8} \times \frac{4}{7} \right) \times 3 = \frac{10}{21}$

(e)



$\frac{1}{2} r^2 \theta = \frac{3\pi}{4}$
 $\frac{1}{2} r^2 \times \frac{\pi}{6} = \frac{3\pi}{4}$
 $\therefore r^2 = \frac{3}{4} \times 12 = 9$
 $\therefore r = 3$

$\therefore l = r\theta = 3 \times \frac{\pi}{6} = \frac{\pi}{2} \text{ cm}$

$$(f) \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \tan^{-1}(1) = \frac{\pi}{6} + \frac{\pi}{4} = \frac{10\pi}{24} = \frac{5\pi}{12}$$

$$k = \frac{5}{12}$$

$$(g) \int \frac{\ln 3x}{x} dx \quad u = \ln 3x$$

$$\therefore du = \frac{3}{3x} dx = \frac{dx}{x}$$

$$= \int \ln 3x \times \frac{dx}{x}$$

$$= \int u du = \frac{1}{2} u^2 + C$$

$$= \frac{1}{2} \ln^2 3x + C$$

$$(2) (a) (i) \frac{d(x \cos x)}{dx} = x(-\sin x) + (1) \cos x$$

$$= \cos x - x \sin x$$

$$(ii) \frac{d(\ln(\cos 5x))}{dx} = \frac{1}{\cos 5x} \times -5 \sin 5x$$

$$= -\frac{5 \sin 5x}{\cos 5x} = -5 \tan 5x$$

$$(iii) \frac{d\left(\tan^{-1}\left(\frac{x}{3}\right)\right)}{dx} = \frac{1}{3} \times \frac{1}{1 + \left(\frac{x}{3}\right)^2} = \frac{1}{3} \times \frac{9}{9 + x^2}$$

$$= \frac{3}{9 + x^2}$$

$$(iv) \frac{d(e^{\tan x})}{dx} = \sec^2 x e^{\tan x}$$

$$(v) \frac{d\left(\left[1 + \cos^{-1}(3x)\right]^3\right)}{dx} = 3\left[1 + \cos^{-1}(3x)\right]^2 \times \frac{-3}{\sqrt{1-9x^2}} = -\frac{9\left[1 + \cos^{-1}(3x)\right]^2}{\sqrt{1-9x^2}}$$

$$2(b) (i) \int x^2 e^{2x^3+1} dx$$

$$= \frac{1}{6} \int (6x^2) e^{2x^3+1} dx$$

$$= \frac{1}{6} e^{2x^3+1} + C$$

$$[N.B. \frac{d(2x^3+1)}{dx} = 6x^2]$$

$$(ii) \int \frac{5}{4+x^2} dx = \frac{5}{2} \int \frac{2}{4+x^2} dx$$

$$= \frac{5}{2} \tan^{-1}\left(\frac{x}{2}\right) + C$$

$$(iii) \int \frac{x^2}{x^3+7} dx = \frac{1}{3} \int \frac{3x^2}{x^3+7} dx = \frac{1}{3} \ln|x^3+7| + C$$

$$(iv) -\sin(\pi-x) = -\sin x$$

$$\int -\sin x dx = \cos x + C$$

$$(3) (a) \text{ PRBN IG}$$

$$3 \text{ letters} = \binom{6}{3} = 20 \quad \therefore \frac{12}{20} = \frac{3}{5}$$

$$1 \text{ vowel} = \binom{4}{2} \times \binom{2}{1} = 6 \times 2$$

$$(b) (i) 6 \text{ numbers} \Rightarrow 5! = 120$$

$$(ii) 135 \quad 246$$

No even numbers together means alternating



place an even number first

$$\therefore 2! \times 3! = 2 \times 3 = 12 \text{ ways}$$

$$\therefore \text{Prob} = \frac{12}{120} = \frac{1}{10}$$

$$\therefore \text{At least 2} = 1 - \frac{1}{10} = \frac{9}{10}$$

(c) (i) $f(x) = x^2 + e^{-\frac{1}{2}x} - 5$ is continuous

$f(-2) = 4 + e^{-5} = e^{-1} > 0$ ($\because e > 3$)

$f(-1) = 1 + e^{\frac{1}{2}} - 5 = e^{\frac{1}{2}} - 4 < 0$

$\therefore f(-2), f(-1) < 0$ and with f continuous
 $\exists c$ s.t. $f(c) = 0, -2 < c < -1$

(ii) $x_0 = -2$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} \quad \left[\begin{array}{l} f(-2) = e^{-1} \\ f'(x) = 2x - \frac{1}{2}e^{-\frac{1}{2}x} \\ f'(-2) = -4 - \frac{1}{2}e \\ = \frac{-8 - e}{2} = -\frac{e+8}{2} \end{array} \right.$$

$\therefore x_1 = -2 - \frac{e-1}{-\frac{e+8}{2}}$

$$= -2 + \frac{2(e-1)}{e+8} = \frac{-2(e+8) + 2(e-1)}{e+8} = \frac{-2e - 16 + 2e - 2}{e+8} = -\frac{18}{e+8}$$


(d) (i) $\frac{1+x}{1-x} = \frac{-(1-x) + 2}{1-x}$
 $= -\frac{(1-x)}{1-x} + \frac{2}{1-x}$
 $= -1 + \frac{2}{1-x}$

(ii) $\int \frac{1+x}{1-x} dx = \int \left(-1 + \frac{2}{1-x} \right) dx$
 $= -x - 2 \ln|1-x| + C$
 $= -x - 2 \ln|1-x| + C$

3(e) $\sqrt{3} \tan \theta = 1$
 $\tan \theta = \frac{1}{\sqrt{3}}$

$\therefore \theta = n\pi + \frac{\pi}{6}$

(4) (a) (i)



(ii) $\sin(-\frac{\pi}{2}) = -1$
 $\cos(2 \times -\frac{\pi}{2}) = \cos(-\pi) = -1$
 $\sin \frac{\pi}{6} = \frac{1}{2}$
 $\cos(2 \times \frac{\pi}{6}) = \cos \frac{\pi}{3} = \frac{1}{2}$


(iii) Area = $\int_{-\frac{\pi}{2}}^{\frac{\pi}{6}} (\cos x - \sin x) dx = \left[\frac{1}{2} \sin 2x + \cos x \right]_{-\frac{\pi}{2}}^{\frac{\pi}{6}}$
 $= \left(\frac{1}{2} \sin \frac{\pi}{3} + \cos \frac{\pi}{6} \right) - (0)$
 $= \frac{1}{2} \times \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2}$
 $= \frac{3}{2} \times \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{4}$

(b) $f(x) = 3 \sin^{-1}(\frac{x}{2})$

(i) $f(2) = 3 \sin^{-1}(1) = 3 \times \frac{\pi}{2} = \frac{3\pi}{2}$

(ii) $-1 \leq \frac{x}{2} \leq 1 \Rightarrow -2 \leq x \leq 2$

$-\frac{\pi}{2} \leq \frac{y}{3} \leq \frac{\pi}{2} \Rightarrow -\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$



4(c) $\tan \left[2 \tan^{-1} \left(-\frac{1}{2} \right) \right]$ $\left\| \begin{array}{l} -\frac{\pi}{2} < \tan^{-1} x < \frac{\pi}{2} \end{array} \right.$

let $\alpha = \tan^{-1} \left(-\frac{1}{2} \right)$


$\therefore \tan \alpha = -\frac{1}{2}$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{2 \times (-\frac{1}{2})}{1 - (-\frac{1}{2})^2} = \frac{-1}{1 - \frac{1}{4}} = \frac{-1}{\frac{3}{4}} = -\frac{4}{3}$$

(d) $g(x) = \cos^{-1} x + \sin^{-1} x$

(i) $g'(x) = \frac{-1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-x^2}} = 0$

(ii) $\therefore g(x) = \text{constant}$ for $-1 \leq x \leq 1$
 $g(0) = \cos^{-1}(0) + \sin^{-1}(0) = \frac{\pi}{2} + 0 = \frac{\pi}{2}$



(5) (a) $f(x) = x \sin^{-1}(x^2)$

(i) $-1 \leq x^2 \leq 1 \Rightarrow -1 \leq x \leq 1$
 $\Rightarrow -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

(ii) $f'(x) = \sin^{-1}(x^2) + x \left(\frac{2x}{\sqrt{1-x^4}} \right)$
 $= \sin^{-1}(x^2) + \frac{2x^2}{\sqrt{1-x^4}}$
 $= \text{even function}$

(iii)

x	0	0	0
f'(x)	+	0	+

\therefore HPOI


\nearrow
 \searrow
 \nearrow
 \searrow

(iv) $f'(1) = \sin^{-1} 1 + \frac{2}{\sqrt{0}}$

\therefore vertical tangent

[OR: $\sin^{-1}(x^2)$ has a double root at $x=0$ \therefore even
 $\therefore x \sin^{-1}(x^2)$ has a triple root
 \therefore HPOI]

5 (a) (v) $f(x)$ is an odd function




(b) (i) $x^2 + 2x - 3 = (x^2 + 2x + 1) - 4$
 $= (x+1)^2 - 4$

$\therefore \int \frac{x}{\sqrt{x^2+2x-3}} dx = \int \frac{x dx}{\sqrt{(x+1)^2 - 4}}$

(ii) $2 \sec u = x+1$
 $\therefore 2 \sec u + \tan u du = dx$

$\tan^2 u + 1 = \sec^2 u$
 $[4 \sec^2 u - 4 = 4(\sec^2 u - 1) = 4 \tan^2 u]$



$\therefore \tan u = \frac{\sqrt{x^2+2x-3}}{2}$

$$\begin{aligned} & \int \frac{x dx}{\sqrt{(x+1)^2 - 4}} \\ &= \int \frac{(2 \sec u - 1) 2 \sec u \tan u du}{\sqrt{4 \sec^2 u - 4}} \\ &= \int \frac{2(2 \sec u - 1) \sec u \tan u du}{2 + 2 \tan^2 u} \\ &= \int (2 \sec^2 u - \sec u) du \\ &= 2 \tan u - \int \sec u du \\ &= \sqrt{x^2+2x-3} - \int \sec u du \end{aligned}$$

[could be done reversing the substitution BUT worth more than 2 marks!]

not an Ext 1 integral
 $= \sqrt{x^2+2x-3} - \ln |\sec u + \tan u| + C$
 $= \sqrt{x^2+2x-3} - \ln \left| \frac{x+1}{2} + \frac{\sqrt{x^2+2x-3}}{2} \right| + C$
 $= \sqrt{x^2+2x-3} - \ln |x+1 + \sqrt{x^2+2x-3}| + K$

(5) (c) $y = \frac{1}{\sqrt{1+x^2}}$

$$\begin{aligned} \therefore V &= \pi \int_{\frac{1}{\sqrt{2}}}^{\sqrt{3}} y^2 dx \\ &= \pi \int_{\frac{1}{\sqrt{2}}}^{\sqrt{3}} \frac{1}{1+x^2} dx \\ &= \pi \left[\tan^{-1}(x) \right]_{\frac{1}{\sqrt{2}}}^{\sqrt{3}} \\ &= \pi \left[\tan^{-1}(\sqrt{3}) - \tan^{-1}\left(\frac{1}{\sqrt{2}}\right) \right] \\ &= \pi \left[\frac{\pi}{3} - \frac{\pi}{6} \right] \\ &= \pi \left[\frac{\pi}{6} \right] \\ &= \frac{\pi^2}{6} \text{ c.u.} \end{aligned}$$

(6) a) $\int_0^{\sqrt{2}} x^2 \sqrt{x^2+1} dx \equiv \frac{1}{2} \int_0^{\sqrt{2}} \sqrt{x^2+1} (2x dx)$

$$\begin{array}{l} u = x^2 + 1 \quad \therefore du = 2x dx \\ x = 0 \Rightarrow u = 1 \\ x = \sqrt{2} \Rightarrow u = 3 \end{array} \left| \begin{array}{l} = \frac{1}{2} \int_1^3 u^{\frac{1}{2}} du \\ = \frac{1}{2} \times \frac{2}{\frac{3}{2}} u^{\frac{3}{2}} \Big|_1^3 \\ = \frac{3}{8} [3^{\frac{3}{2}} - 1] \\ = \frac{3}{8} [3\sqrt{3} - 1] \\ = \frac{3}{8} [3^2\sqrt{3} - 1] \end{array} \right.$$

$27 \times 3 = 81$

6(b) $\cos 3x = \sin x$

(i) $\cos\left(\frac{\pi}{2} - A\right)$
 $= \cos \frac{\pi}{2} \cos A + \sin \frac{\pi}{2} \sin A$
 $= 0 \times \cos A + 1 \times \sin A$
 $= \sin A$

(ii) $\cos 3x = \sin x$
 $\Rightarrow \cos 3x = \cos\left(\frac{\pi}{2} - x\right)$

$\therefore 3x = 2n\pi \pm \left(\frac{\pi}{2} - x\right)$

$$\begin{array}{l} 3x = 2n\pi + \frac{\pi}{2} - x \\ 4x = (4n+1)\pi \\ x = \frac{(4n+1)\pi}{4} \end{array} \left| \begin{array}{l} 3x = 2n\pi - \frac{\pi}{2} + x \\ 2x = \frac{(4n-1)\pi}{2} \\ x = \frac{(4n-1)\pi}{4} \end{array} \right.$$

$\therefore x = \frac{(4n+1)\pi}{4}$

(c) (i) $\frac{d(\tan^3 x)}{dx} = 3 \tan^2 x \cdot \sec^2 x$
 $= 3(\sec^2 x - 1) \sec^2 x$
 $= 3 \sec^4 x - 3 \sec^2 x$

(ii) $\therefore \tan^3 x = \int (3 \sec^4 x - 3 \sec^2 x) dx$

$\therefore \tan^3 x \Big|_0^{\pi/4} = 3 \int_0^{\pi/4} \sec^4 x dx - 3 \int_0^{\pi/4} \sec^2 x dx$

$\therefore 3 \int_0^{\pi/4} \sec^4 x dx = [\tan^3 \frac{\pi}{4} - \tan^3(0)] + 3 \tan x \Big|_0^{\pi/4}$
 $= (1-0) + 3(1-0)$
 $= 4$

$\therefore \int_0^{\pi/4} \sec^4 x dx = \frac{4}{3}$

$$(6)(d) \int_0^{\pi/6} \sin^2 x \, dx$$

$$= \frac{1}{2} \int_0^{\pi/6} 2 \sin^2 x \, dx$$

$$= \frac{1}{2} \int_0^{\pi/6} (1 - \cos 2x) \, dx$$

$$= \frac{1}{2} \left[x - \frac{1}{2} \sin 2x \right]_0^{\pi/6}$$

$$= \frac{1}{2} \left[\frac{\pi}{6} - \frac{1}{2} \sin \frac{\pi}{3} \right]$$

$$= \frac{\pi}{12} - \frac{1}{4} \times \frac{\sqrt{3}}{2}$$

$$= \frac{\pi}{12} - \frac{\sqrt{3}}{8}$$

$$= \frac{2\pi - 3\sqrt{3}}{24}$$

$$\cos 2A = 1 - 2\sin^2 A$$