



**SYDNEY BOYS HIGH SCHOOL**  
**MOORE PARK, SURRY HILLS**

**2006**  
**HIGHER SCHOOL CERTIFICATE**  
**ASSESSMENT TASK #2**

# Mathematics

## General Instructions

- Reading Time – 5 Minutes
- Working time – 90 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- Each Section is to be returned in a separate bundle.
- All necessary working should be shown in every question.

## Total Marks – 85

- Attempt questions 1 – 6
- Hand up in 3 separate booklets clearly labelled Section A, Section B and Section C

Examiner: *A. Fuller*

## SECTION A

### Question 1 (13 marks)

- (a) Convert  $80^\circ$  to radians in exact form. [1]
- (b) Convert  $\frac{17\pi}{12}$  radians to degrees. [1]
- (c) Differentiate the following:
- (i)  $4x^2 + 5$  [1]
- (ii)  $(2x^3 - 5)^4$  [2]
- (iii)  $\frac{3x+1}{2x-1}$  [2]
- (iv)  $x\sqrt{1-2x}$  [2]
- (d) Find a primitive of:
- (i)  $\frac{2}{x^4}$  [1]
- (ii)  $\sqrt{x}$  [1]
- (e) Find  $f''(2)$  if  $f(x) = x^5$  [2]

**Question 2 (15 marks)**


- (a) A(-1,5), B(2,1) and C(4, k) are collinear. Find the value of k. [3]
- (b) Find  $\int \frac{x^2 + 1}{x^2} dx$  [2]
- (c) Evaluate  $\int_{-1}^1 (x-1)(x+1) dx$  [3]
- (d) A die is tossed twice. The sum of the numbers which appear on the upmost face of the die is calculated. Using a table or otherwise:
- (i) Find the probability that the sum is greater than 8. [2]
- (ii) It is known that a 4 appears on the die at least once in the two throws. Find the probability that the sum is greater than 8. [2]
- (e) The vertices of a triangle are A(1,3), B(8,2) and C(4,-1).
- (i) Find the coordinates of D and E, the midpoints of AC and AB respectively. [1]
- (ii) Hence, show that DE is parallel to CB. [2]

## SECTION B

### Question 3 (14 marks)

(a) If  $y = (x^2 + 4)(x - 3)$ , solve  $\frac{dy}{dx} = 4$ . [3]

(b) The diagram below is the graph of  $y = 3 + 3 \cos x$



(i) Copy this graph onto your answer sheet.

(ii) State the amplitude and period of  $y = 3 \sin 2x$ . [1]


(iii) On the same graph as (i), sketch  $y = 3 \sin 2x$  for  $0 \leq x \leq 2\pi$ . [2]

(iv) How many solutions are there to the equation  $3 + 3 \cos x = 3 \sin 2x$ ? [1]

(c) A class of 30 students contains 18 students who play cricket, 13 who play tennis and 5 who play both cricket and ~~softball~~<sup>tennis</sup>. If one student is chosen at random find the probability that this student plays neither cricket nor tennis. [2]

- (d) The diagram below shows a paddock with one side bounded by a river. AE is a boundary fence 40 metres in length. AO, BP, CQ, DR, ES are offsets measured from the fence to the river with lengths as shown.  $AB = BC = CD = DE$ .

[3]



Use Simpson's rule with 5 function values as shown on the diagram to approximate the area of the paddock.

- (e) The gradient function of a curve is  $3x^2 - 1$  and the curve passes through the point (4,1). Find the equation of the curve.

[2]

**Question 4 (13 marks)**

(a) A girl has 5 tickets in a raffle where 100 tickets are sold.

First prize is drawn discarded and then the second prize is drawn.

Find the probability that she wins:

(i) first prize [1]

(ii) second prize [2]

(b) Consider the curve  $y = x^4 - 4x^3$

(i) Find the coordinates of the stationary points. [2]

(ii) Determine the nature of these stationary points. [2]

(iii) For what values of  $x$  is the curve concave up? [1]

(iv) For what values of  $x$  is the curve decreasing? [1]


(v) Hence sketch the curve. [2]

(c) If the probability that an event E occurs is  $\frac{1}{x}$ , express the probability that E does not occur as a single fraction. [2]

### SECTION C


#### Question 5 (15 marks)

- (a) A flower bed is made in the shape of a minor sector with angle  $\theta$  and radius 2 metres.




- (i) If the area of the flower bed is  $1\frac{1}{6}m^2$ , find the angle  $\theta$  to the nearest minute. [2]
- (ii) Find the perimeter of the flower bed to the nearest cm. [2]

- (b) ABCD is a parallelogram. The bisectors of angles ADC and BCD meet at P on the side AB. Prove that:




- (i)  $\angle DPC$  is a right angle. [2]
- (ii)  $\triangle ADP$  is isosceles. [2]
- (iii)  $AB = 2BC$  [2]

- (c) The graph below of the function  $f$  consists of a quarter circle AB and two line segments BC and CE.




- (i) Evaluate  $\int_0^8 f(x) dx$ . [2]
- (i) For what value(s) of  $x$  satisfying  $0 < x < 8$  is the function  $f$  not differentiable? [1]
- (d) The diagram below shows the graph of the gradient function of a curve. For what value(s) of  $x$  does  $f(x)$  have a local maximum? Justify your answer. [2]






**Question 6 (15 marks)**

- (a) The diagram below shows the graphs of the functions  $y = -x^2 + 7x - 6$  and  $y = x + 2$ .



- (i) Show that the value of A and B is 2 and 4 respectively. [2]
- (ii) Calculate the area of the shaded region. [2]
- (iii) Write down a pair of inequalities that specify the shaded region. [1]


(b)



(i) Find the equation of the tangent and the equation of the normal to the curve  $y = -2x^2 + 6x + 3$  at the point  $A(2,7)$ . [4]

(ii) The tangent cuts the  $x$  axis at  $B$ . The normal cuts the  $x$  axis at  $C$  as shown in the diagram. Find the values of  $B$  and  $C$ . Hence, find the area of  $\Delta ABC$ . [2]

(c) In the diagram below  $AC$  is parallel to  $FD$  and  $BF$  is parallel to  $CE$ .  $B$  lies on  $AC$ ,  $D$  lies on  $CE$  and  $F$  lies on  $AE$ .  $AF = 6\text{cm}$ ,  $FE = 14\text{cm}$  and  $AB = 7\text{cm}$ .



(i) Find  $BC$ . [2]

(ii) Find the ratio of  $BF$  to  $DE$ . [2]

**END OF TEST**

Ⓐ

$$80 \times \frac{\pi}{180} = \frac{4\pi}{9} \text{ c}$$

Q11

2006  
2U  
TASK 2

Ⓑ

$$\frac{17\pi}{12} = 255^\circ$$

Ⓒ

$$i) \frac{d}{dx}(4x^2+5) = 8x$$

$$ii) \frac{d}{dx}(-(2x^3-5)^4) = 4 \times 6x^2 \times (2x^3-5)^3 \\ = 24x^2(2x^3-5)^3$$

$$iii) \frac{d}{dx} \left( \frac{3x+1}{2x-1} \right) = \frac{v u' - u v'}{v^2} \\ = \frac{(2x-1)(3) - (3x+1)(2)}{(2x-1)^2} \\ = \frac{\cancel{6x} - 3 - \cancel{6x} - 2}{(2x-1)^2} \\ = \frac{-5}{(2x-1)^2}$$

~~~~~

Q1 (c) (iv).

$$\begin{aligned}\frac{d}{dx} \left( x \sqrt{1-2x} \right) &= v u' + u v' \\ &= \sqrt{1-2x} + x \left( \frac{1}{2} (1-2x)^{-\frac{1}{2}} (-2) \right) \\ &= \sqrt{1-2x} - \frac{x}{\sqrt{1-2x}} \\ &= \frac{1-2x-x}{\sqrt{1-2x}} \\ &= \frac{1-3x}{\sqrt{1-2x}}.\end{aligned}$$

$$\begin{aligned}d) \ i) \int \frac{2}{x^4} \cdot dx &= \int 2x^{-4} \cdot dx \\ &= \frac{2}{-3} x^{-3} + C \\ &= -\frac{2}{3x^3} + C.\end{aligned}$$

$$\text{Q1) d) ii) } \int \sqrt{x} \cdot dx = \int x^{\frac{1}{2}} dx.$$

$$= \frac{2}{3} x^{\frac{3}{2}} + C$$

e)

~~$$f(x) = x^5.$$~~

~~$$f'(x) = \frac{x^6}{6}$$~~

~~$$f''(x) = \frac{x^7}{7 \times 6}$$~~
~~$$= \frac{x^7}{42}.$$~~

~~$$f''(2) = \frac{2^7}{42}$$~~

~~$$= 3.05 \text{ (2 dp)}$$~~

$$f(x) = x^5,$$

$$f'(x) = 5x^4$$

$$f''(x) = 20x^3.$$

$$f''(2) = 20(2)^3$$

$$= 160$$

## QUESTION(2)

a)  $A(1,5) B(2,1)$ .

Two pt.

~~$m_1 = \frac{5-1}{1-2}$~~

~~$= \frac{4}{-3}$~~

~~$= \frac{4}{-3}$~~

$$\frac{y-1}{x-2} = \frac{5-1}{-1-2}$$

$$-3(y-1) = 4(x-2)$$

$$-3y+3 = 4x-8$$

~~$4x-3y$~~   $y = \frac{4x-11}{-3}$

$C(4, k)$

$$y = \frac{4(4)-11}{-3}$$

$$= \frac{-3}{-3} + \frac{5}{-3} = \boxed{\frac{5}{3}}$$

$$= 1.$$

$\therefore k$  is 1 and  $C(4, \frac{5}{3})$ .

∴ Question 2.

$$b). \int \frac{x^2+1}{x^2} \cdot dx.$$

$$= \int 1 + \frac{1}{x^2} \cdot dx.$$

$$= \int 1 + x^{-2} \cdot dx.$$

$$= x + \frac{x^{-1}}{-1} \cdot dx.$$

$$= x - \frac{1}{x} + C$$

$$c). \int_{-1}^1 (x-1)(x+1) \cdot dx.$$

$$= \int_{-1}^1 x^2 - 1 \cdot dx.$$

$$= \left[ \frac{x^3}{3} - x \right]_{-1}^1$$

$$= \left[ \frac{1}{3} - 1 \right] - \left[ -\frac{1}{3} + 1 \right].$$

$$= -\frac{2}{3} - \frac{2}{3}.$$

$$= -\frac{4}{3} = \underline{\underline{-1\frac{1}{3}}}$$

# QUESTION (2) (d).

i)

|   | 1 | 2 | 3 | 4  | 5  | 6  |
|---|---|---|---|----|----|----|
| 1 | 2 | 3 | 4 | 5  | 6  | 7  |
| 2 | 3 | 4 | 5 | 6  | 7  | 8  |
| 3 | 4 | 5 | 6 | 7  | 8  | 9  |
| 4 | 5 | 6 | 7 | 8  | 9  | 10 |
| 5 | 6 | 7 | 8 | 9  | 10 | 11 |
| 6 | 7 | 8 | 9 | 10 | 11 | 12 |

$$P(\text{sum} > 8) = \frac{10}{36} = \frac{5}{18}$$

$$P(1,4 \& \text{sum} > 8) = \frac{4}{36} = \frac{1}{9}$$

(e) A(1,3) B(8,2) C(4,-1).

$$\text{Midpoint AB} = \frac{E}{2} = \left(\frac{9}{2}, \frac{5}{2}\right)$$

$$\begin{aligned} \text{Midpoint AC} = D &= \left(\frac{4+1}{2}, \frac{3-1}{2}\right) \\ &= \left(\frac{5}{2}, 1\right). \end{aligned}$$

$$\text{grad DE} = \frac{5/2 - 1}{9/2 - 5/2} = \frac{3/2}{2} = \frac{3}{4}$$

$$\text{grad CB} = \frac{2 - -1}{8 - 4} = \frac{3}{4}$$

∴ CB // DE.



### Question 3

a) If  $y = (x^2 + 4)(x - 3)$ , solve  $\frac{dy}{dx} = 4$ .

$$y = x^3 - 3x^2 + 4x - 12$$

$$\frac{dy}{dx} = 3x^2 - 6x + 4$$

$$\therefore 4 = 3x^2 - 6x + 4$$

$$0 = 3x^2 - 6x$$


$$0 = 3x(x - 2)$$

$$3x = 0, \quad x - 2 = 0$$

$$x = 0, \quad x = 2$$

b) i)

iii)




ii)  $y = 3\sin 2x$ . amplitude = 3

$$\text{period} = \frac{2\pi}{2} = \pi$$

iv) There are 2 solutions.

c) class.



$$\begin{aligned} P(\text{neither}) &= 4/30 \\ &= 2/15 \end{aligned}$$

|    |        |    |    |    |    |    |
|----|--------|----|----|----|----|----|
| d) | $x$    | 0  | 10 | 20 | 30 | 40 |
|    | $f(x)$ | 20 | 14 | 12 | 7  | 8  |

$$h = \frac{40-0}{4} = \frac{40}{4} = 10$$

$$\begin{aligned} \int_0^{40} f(x) dx &\doteq \frac{10}{3} [(20+8) + 4(14+7) + 2(12)] \\ &\doteq \frac{10}{3} [28 + 4(21) + 2(12)] \\ &= 453 \frac{1}{3} \text{ m}^2 \end{aligned}$$

e)  $f(x) = 3x^2 - 1$  P.t. point (4, 1)

$$\begin{aligned} f(x) &= \int f'(x) \\ &= \int 3x^2 - 1 \\ &= x^3 - x + C \end{aligned}$$

$$\begin{aligned} f(4) = 1 \quad \therefore \quad 1 &= 4^3 - 4 + C \\ 1 &= 64 - 4 + C \\ 1 &= 60 + C \\ C &= -59 \end{aligned}$$

$$\therefore f(x) = x^3 - x - 59$$

## Question 4

a) i)  $P(\text{1st prize}) = \frac{5}{100}$   
 $= \frac{1}{20}$

ii)  $P(\text{2nd prize}) = P(\text{1st \& 2nd prize}) + P(\text{not 1st \& 2nd prize})$   
 $= (\frac{5}{100} \times \frac{4}{99}) + (\frac{95}{100} \times \frac{5}{99})$   
 $= \frac{1}{20}$

b)  $y = x^4 - 4x^3$   
 $y' = 4x^3 - 12x^2$

$$y'' = 12x^2 - 24x$$

i) SP @  $y' = 0$   
 $0 = 4x^3 - 12x^2$   
 $0 = 4x^2(x - 3)$   
 $4x^2 = 0, x - 3 = 0$   
 $x = 0, x = 3$

when  $x = 0, y = 0$   
 $\therefore$  SP @  $(0, 0)$   
when  $x = 3, y = 3^4 - 4 \times 3^3$   
 $= -27$   
 $\therefore$  SP @  $(3, -27)$

ii)  $y'' = 0$  helps determine nature of st. pts.

$$y''(0) = 0 - 0 = 0$$

$$y''(3) = 12 \times 3^2 - 24 \times 3 = 36$$

- possible pt of inflexion  $\therefore (3, -27)$  is a local minimum

check  $y'$  for sign of derivative.

|      |     |   |    |
|------|-----|---|----|
| $x$  | -1  | 0 | 1  |
| $y'$ | -16 | 0 | -8 |

$\therefore (0, 0)$  is a pt of inflexion.

iii)  $y'' > 0$  for concave up;  $\therefore 12x^2 - 24x > 0$   
 $x^2 - 2x > 0$   
 $x(x - 2) > 0$   
 $\therefore x > 2, x < 0$   
 $\therefore$  the curve is concave up for  $x > 2, x < 0$ .

iv)  $f'(x) < 0$  for curve decreasing.  
 $4x^3 - 12x^2 < 0$   
 $x^2(x - 3) < 0$   
 $x^2 < 0, x < 3$   
 $\therefore$  the curve is decreasing for  $x < 3, x \neq 0$

( )


$$c) P(E) = 1/x$$

( )

$$P(\text{not } E) = 1 - 1/x$$
$$= \frac{x-1}{x}$$


( )

Q4b) v)



## Solution to Section (c)

Question (5). (a)


(i)   $A = \frac{1}{2} r^2 \theta$   
 $1.6 = 2\theta$   
 $\therefore \theta = 0.8$  \*

[2] i.e.  $0.8^c = \frac{180}{\pi} \times 0.8$   
 $\doteq 45^\circ 50'$  \*

(ii)  $l = r\theta$   
 $= 2 \times 0.8$   
 $= 1.6$  \*

$\therefore p = l + 2r = 5.6 \text{ m}$   
 $= 560 \text{ cm.}$  \*

[2]



(i) In parm. ABCD.  
 $\angle ADC + \angle BCD = 180^\circ$   
 (Co-interior  $\angle$ s are supplementary,  $AD \parallel BC$ )

$\therefore 2a + 2x = 180$   
 $a + x = 90^\circ$

$\angle DPC = 180^\circ - (a+x)$   
 (Angle sum of  $\triangle DPC$ )

i.e.  $\angle DPC = 180^\circ - 90^\circ$   
 $= 90^\circ$  [2]

(ii) In  $\triangle APD$ . [2]

$\angle PDC = \angle APD = a^\circ$   
 (alternate  $\angle$ s,  $AP \parallel DC$ )

$\therefore \angle ADP = \angle APD = a^\circ$   
 i.e.  $\triangle ADP$  is isosceles.

(iii) Similarity (from (ii))  
 •  $BP = BC$ . ( $\triangle BPC$  is isosceles)  
 and  $AD = AP$  (proven).

• but  $AD = BC$   
 (opposite sides of a parm equal). [2]

$AB$   
 $\therefore = AP + PB$   
 $= AD + BC.$   
 $= 2BC.$

(c)

(i)  $\int_0^8 f(x) dx$   
 $= \frac{1}{4}(\pi \times 2^2) + 2^2 + \frac{1}{2}(2) \times 2$   
 $- \frac{1}{2}(2) \times 2$  [2]  
 $= \pi + 4$  (7.142)

(ii) At  $x=4$   $f$  is [1]  
 not differentiable.

(d) At  $x=5$  [2]

$f'(5) = 0$

$f'(5-\epsilon) > 0$  (for small positive  $\epsilon$ )

$f'(5+\epsilon) < 0$

$\therefore$  By the 1<sup>st</sup> derivative test  $f(x)$  has a local maximum at  $(5, 0)$ .

Question (6)

(a)  $y = -x^2 + 7x - 6$   
 $y = x + 2$

$-x^2 + 7x - 6 = x + 2$

(i)  $x^2 - 6x + 8 = 0$

$(x - 4)(x - 2) = 0$

$\therefore x = 2, y = 4$   
 $x = 4, y = 6$

$(2, 4) (4, 6)$  [2]

$\therefore$  A = 2 and B = 4.

(ii)  $A = \int_2^4 [(-x^2 + 7x - 6) - x - 2] dx$

$= \int_2^4 (-x^2 + 6x - 8) dx$

$= \left[ -\frac{x^3}{3} + 3x^2 - 8x \right]_2^4$


$= \left[ -\frac{64}{3} + 48 - 32 \right] - \left[ -\frac{8}{3} + 12 - 16 \right]$

$= -\frac{56}{3} + 16 + 4$   
 $= \frac{4}{3}$  [2]

(iii)  $-x^2 + 6x - 8 > 0$

$x^2 - 6x + 8 < 0$

$(x - 4)(x - 2) < 0$



$2 < x < 4$  [2]

(b)  $y = -2x^2 + 6x + 3$

A (2, 7)

$\frac{dy}{dx} = -4x + 6$

$\frac{dy}{dx} \Big|_{x=2} = -2$


$\therefore y - 7 = -2(x - 2)$

$2x + y - 11 = 0$  [tg+]

$y - 7 = \frac{1}{2}(x - 2)$

$2y - 14 = x - 2$

$\therefore x + 2y + 12 = 0$   
 (normal) [4]



$2x = 11 \quad x = \frac{11}{2}$  [2]

$x = -12$

B  $(\frac{11}{2}, 0)$  C  $(-12, 0)$

$\therefore$  Area =  $\frac{1}{2} \times 17\frac{1}{2} \times 7$

(c)

Let BC = x.

(i)  $\frac{7}{7+x} = \frac{6}{30}$

$70 = 21 + 3x$  [2]

$9 = 3x$

$x = 49/3$

(ii) BF : DE

$7 = \frac{49}{3}$

$21 = 49$

$3 = 7$  [2]