



SYDNEY BOYS HIGH  
MOORE PARK, SURRY HILLS

APRIL 2010

TASK #2  
YEAR 12

## Mathematics Ext 1

**General Instructions:**

- Reading time—5 minutes.
- Working time—90 minutes.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may NOT be awarded for messy or badly arranged work.
- Answer in simplest exact form unless otherwise stated.

Total marks—72 Marks

- Attempt questions 1–6.
- The mark-value of each question is boxed in the right margin.
- Start each NEW section in a separate answer booklet.
- Hand in your answer booklets in 3 sections:  
Section A (Questions 1 and 2),  
Section B (Questions 3 and 4),  
Section C (Questions 5 and 6),

Examiner: Mr D. Hespe

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

### Section A

**Question 1 (12 points)**

- (a) (i) How many four-letter arrangements can be made from the letters IOLS? [1]
- (ii) In the Herald's *Target* competition, arrangements ending in *S* are not allowed. How many four-letter *Target* arrangements can be made from the letters IOLS? [1]
- (b) Prove that a line from the centre of a circle to the midpoint of a chord is perpendicular to the chord. [3]
- (c) Use the graph of  $y = \sin x$  to illustrate why [2]

$$\int_{-1}^1 \sin x \, dx = 0.$$

- (d) Evaluate  $\lim_{x \rightarrow 0} \left\{ \frac{\tan 4x}{7x} \right\}$ . [1]
- (e) (i) Differentiate  $x \log_e x$ . [1]
- (ii) Hence integrate  $\log_e x$ . [2]
- (f) On a certain railway line, there are eleven railway stations at which a train can stop. The rail authority needs to print tickets for travel between every possible pair of stations on the line. How many different one-way tickets must be printed if the ticket specifies which direction the passenger is travelling? [1]

**Question 2 (12 points)**

(a) Find

(i)  $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \tan^2 x dx,$

Marks

[2]

(ii)  $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \sin^2 x dx.$

[2]

(b) (i) Prove by mathematical induction that

$$\sum_{r=1}^n \frac{1}{r(r+1)} = \frac{n}{n+1}.$$

[4]

(ii) What can you say about

[1]

$$\lim_{n \rightarrow \infty} \left( \sum_{r=1}^n \frac{1}{r(r+1)} \right) ?$$

(c) Using one iteration of Newton's Method and a first approximation of  $x_0 = 0.7$ , find, correct to three decimal places, a second approximation to

$$y = \sin^3 x - 0.25.$$

[3]

**Section B**

(Use a separate writing booklet.)

Marks

**Question 3 (12 points)**(a) By writing  $\cot x$  as  $\frac{\cos x}{\sin x}$ , evaluate

$$\int_{\pi/6}^{\pi/3} \cot x dx.$$

[2]

(b) Given that a team of five players is to be selected from a group of ten boys, find the number of teams that contain

(i) at least one of the two best players,

[2]

(ii) no more than one of the three youngest players.


[3]

(c) (i) Show that  $e^x = 3x + 2$  has a solution between 2 and 2.5.

[2]

(ii) Hence use "halving the interval" to find, correct to one decimal place, a solution in the interval [2, 2.5].

[3]

**Question 4 (12 points)**

Find the value of  $x$ , giving reasons.

- (b) Prove that  $7^{n+1} + 3^n$  is divisible by 4 for all positive integers  $n$ .


Marks

[3]

- (c) A team of three people is to be chosen from six men and five women by putting the eleven names in a hat and drawing out three simultaneously at random. Find the probability that the team will be of mixed sex.

[4]

(d)



[2]

Consider the above graph of  $y = f(x)$ . The value  $a$  shown on the axis is taken as the first approximation to the solution  $b$  of  $f(x) = 0$ .

Is the second approximation obtained by Newton's method a better approximation to  $b$  than  $a$  is? Give a reason for your answer.

**Section C**

(Use a separate writing booklet.)

Marks

**Question 5 (12 points)**

- (a) Differentiate with respect to  $x$

$$(i) \log_e(\cos x),$$

$$(ii) (x+1)e^{-x}.$$

[2]

[2]

- (b) (i) Differentiate  $x + \log_e x$ .

[2]

$$(ii) \text{ Hence or otherwise, find a primitive of } \frac{x+1}{x^2+x\log_e x}.$$

[3]

- (c)  $AKB, CKD$  are two chords of a circle (meeting at an internal point  $K$ ).

Given the following lengths

$AB = 10 \text{ cm}$ ,  $CD = 6 \text{ cm}$ ,  $AK = 1 \text{ cm}$ ,  
calculate the ratio  $AC : BD$ .

[3]

**Question 6 (12 points)**

- | Marks   | STANDARD INTEGRALS   |
|---|--|
| (a) (i) How many three-figure numbers can be formed from the nine digits 1, 2, 3, ..., 9, there being no repetitions? <span style="border: 1px solid black; padding: 2px;">[1]</span>   | $\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$ |
| (ii) From a pack of nine cards numbered 1, ..., 9, three cards are drawn at random and laid on a table from left to right.<br>(α) What is the probability that the number formed by the three digits drawn should exceed 500? <span style="border: 1px solid black; padding: 2px;">[1]</span> | $\int \frac{1}{x} dx = \ln x, \quad x > 0$   |
| (β) What is the probability that the digits should be drawn in ascending order, not necessarily consecutive? <span style="border: 1px solid black; padding: 2px;">[3]</span>  | $\int e^{ax} dx = \frac{1}{a}e^{ax}, \quad a \neq 0$                                     |
| (b) (i) Prove that the graph of $y = \ln x$ is concave down for all $x > 0$ . <span style="border: 1px solid black; padding: 2px;">[2]</span>   | $\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$                                  |
| (ii) Sketch the graph of $y = \ln x$ . <span style="border: 1px solid black; padding: 2px;">[1]</span>  | $\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$                                 |
| (iii) Suppose $1 < a < b$ and consider the points $A(a, \ln a)$ and $B(b, \ln b)$ on the graph of $y = \ln x$ .<br>Find the coördinates of the point $P$ that divides the line segment $AB$ in the ratio 2 : 1. <span style="border: 1px solid black; padding: 2px;">[2]</span>               | $\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$                                |
| (iv) By using (ii) and (iii), deduce that $\frac{1}{3} \ln a + \frac{2}{3} \ln b < \ln(\frac{1}{3}a + \frac{2}{3}b)$ . <span style="border: 1px solid black; padding: 2px;">[2]</span>  | $\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$                          |

**STANDARD INTEGRALS**

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$


$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE:  $\ln x = \log_e x, \quad x > 0$

End of Paper

$$\textcircled{1} \textcircled{a} 4 \times 3 \times 1 \times 1 = 4! = 24 \quad \textcircled{1}$$

$$\textcircled{b} 3 \times 2 \times 1 \times 3 = 18 \quad \textcircled{1}$$



Join AO and BO  
In  $\triangle AOM$  and  $\triangle BOM$

$AO = BO$  radii

$AM = MB$  data

on common side

$\therefore \triangle AOM \cong \triangle BOM$  (SSS)


Hence  $\angle AMO + \angle BMO = 180^\circ$

But  $\angle AMO = \angle BMO$  corresponding angles in congruent triangles

So  $\angle AMO = \angle BMO = 90^\circ$

Hence line from centre of circle to the midpoint of a chord is  $\perp$  to the chord.  $\textcircled{3}$

(4)



area under curve (shaded) cancels out area above curve (shaded)  
 $\sin x$  is an odd function  $\textcircled{1}$

(2)

$$\textcircled{1} \textcircled{a} \textcircled{ii} \lim_{x \rightarrow 0} \frac{\tan 4x}{7x}$$

$$\frac{4}{7} \lim_{x \rightarrow 0} \frac{\tan 4x}{4x} = \frac{4}{7} \times 1 = \frac{4}{7} \quad \textcircled{1}$$

$$\textcircled{1} \textcircled{b} \textcircled{i} \frac{d}{dx}(x \ln x) = x \times \frac{1}{x} + \ln x \times 1 \\ = 1 + \ln x \quad \textcircled{1}$$

$$\textcircled{2} \frac{d}{dx}(x \ln x) - 1 = \ln x$$

$$\textcircled{1} \textcircled{ii} \int \left( \frac{d}{dx}(x \ln x) - 1 \right) dx = \int \ln x dx$$

$$\therefore \int \ln x dx = x \ln x - x + C \quad \textcircled{2}$$

$$\textcircled{1} \textcircled{iii} 2 \times (10+9+8+7+6+5+4+3+2) = 10 \quad \textcircled{1} \textcircled{ii} P_2$$

(2)

$$\textcircled{1} \textcircled{i} \int \frac{1}{3} \tan^2 x dx \quad \text{using } \tan^2 x = \sec^2 x - 1$$

$$= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} (\sec^2 x - 1) dx$$

$$= \left[ (\tan x - x) \right]_{\frac{\pi}{6}}^{\frac{\pi}{3}}$$

$$= \left( \tan \frac{\pi}{3} - \frac{\pi}{3} \right) - \left( \tan \frac{\pi}{6} - \frac{\pi}{6} \right)$$

$$= \sqrt{3} - \frac{\pi}{3} - \frac{1}{\sqrt{3}} + \frac{\pi}{6} = \frac{2}{\sqrt{3}} - \frac{\pi}{6} = \left( \frac{2\sqrt{3}}{3} - \frac{\pi}{6} \right)$$

12.

(2)

(3)

$$\begin{aligned}
 2. (a) (ii) & \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \sin^2 x dx \\
 &= \left( \frac{1}{2}x - \frac{1}{4}\sin 2x \right) \Big|_{\frac{\pi}{6}}^{\frac{\pi}{3}} \\
 &= \left( \frac{\pi}{6} - \frac{1}{4}\sin \frac{2\pi}{3} \right) - \left( \frac{\pi}{12} - \frac{1}{4}\sin \frac{\pi}{3} \right) \\
 &= \frac{\pi}{6} - \frac{\pi}{12} - \frac{1}{4} \times \frac{\sqrt{3}}{2} + \frac{1}{4} \times \frac{\sqrt{3}}{2} \\
 &= \frac{\pi}{12} \quad (2)
 \end{aligned}$$

(b) (i) m. Induction  $\sum_{r=1}^n \frac{1}{r(r+1)} = \frac{n}{n+1}$

LHS means:  $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}$   
sum.

Step 1 let  $n=1$ , LHS =  $\frac{1}{1 \times 2} = \frac{1}{2}$ .

RHS,  $n=1$  term  $\frac{1}{1+1} = \frac{1}{2}$ .  
True for  $n=1$ .

Step 2 let there be a value of  $n=k$ , ( $k \leq n$ ) such that  $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{k(k+1)} = \frac{k}{k+1}$  is true

and we must prove that when  $n=k+1$

$$\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{k(k+1)} + \frac{1}{(k+1)(k+2)} = \frac{k+1}{k+2}.$$

LHS.  $\frac{k}{(k+1)} + \frac{1}{(k+1)(k+2)} = \frac{k(k+2)+1}{(k+1)(k+2)} = \frac{(k+1)^2}{(k+1)(k+2)}$

(4)

$$\begin{aligned}
 \text{LHS } & \frac{(k+1)}{(k+2)} \\
 &= \text{RHS} \\
 \therefore n=k+1 & \text{ is true.}
 \end{aligned} \quad (4)$$

Step 3 By the principle of math. induction  
if it is true for  $n=k$ , it is true for  
 $n=k+1$  etc

(b) (ii) as  $n \rightarrow \infty$ ,  $\lim_{n \rightarrow \infty} \left( \sum_{r=1}^n \frac{1}{r(r+1)} \right) = 1$  (1)

2. (c)  $y = \sin^3 x - 0.25$   
Using  $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$

$$\begin{aligned}
 y &= \sin^3 0.7 - 0.25 = 0.01736 \\
 y' &= 3 \sin^2 x \cos x = 3 \sin^2 0.7 \times \cos 0.7 = 0.95227
 \end{aligned}$$

$$\begin{aligned}
 x_1 &= 0.7 - 0.01736 \\
 &\quad \text{MATERIALS} \quad 0.95227 \\
 &= 0.6827 \quad (3DP). \\
 &\quad \text{ANSWER} \quad 0.682
 \end{aligned} \quad (3)$$

# Extension 1 Solutions

## Section B

Q3 (a)  $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \cot x dx = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos x}{\sin x} dx$

$$\begin{aligned} &= \left[ \ln |\sin x| \right]_{\frac{\pi}{6}}^{\frac{\pi}{3}} \\ &= \ln \sin \frac{\pi}{3} - \ln \sin \frac{\pi}{6} \\ &= \ln \frac{\sqrt{3}}{2} - \ln \frac{1}{2} \\ &= \ln \left( \frac{\sqrt{3}}{2} = \frac{1}{2} \right) \quad [\text{Using log laws}] \\ &= \ln \sqrt{3} \\ &= 0.549 \text{ correct to 3 decimal places} \end{aligned}$$

(b) (i) Having at least one of the two best players means having one of the two or both of them  
 $\therefore$  No. teams  $= {}^2C_1 \times {}^8C_4 + {}^2C_2 \times {}^8C_3$   
 $= 140 + 56$   
 $= 196.$

(ii) Having no more than one of the three youngest players means having none of the youngest players or having just one of them  
 $\therefore$  No. teams  $= {}^7C_5 + {}^3C_1 \times {}^7C_4$   
 $= 21 + 105$   
 $= 126.$

13 (c) (i) Let  $f(x) = e^x - 3x - 2$  and consider  $f(x)=0$ .  
 $\text{Now } f(2) = e^2 - 6 - 2 = 7.39 - 6.2 = 1.19$   
 $\text{and } f(2.5) = e^{2.5} - 7.5 - 2 = 12.18 - 9.5 = 2.68$

Now  $f(x)$  is continuous for  $2 \leq x \leq 2.5$  and changes sign in that interval.  
 $\therefore f(x)$  has a zero between  $x=2$  and  $x=2.5$ .

(ii) Consider  $f\left(\frac{2+2.5}{2}\right) = f(2.25) = 9.488 - 8.75 = 0.734$   
 $\therefore$  There is a root between  $x=2$  and  $x=2.25$

Consider  $f\left(\frac{2+2.25}{2}\right) = f(2.125) = e^{2.125} - 8.375 = -0.001$

$\therefore$  There is a root between  $x=2.125$  and  $x=2.25$

Consider  $f\left(\frac{2.125+2.25}{2}\right) = f(2.188) \doteq 0.349$   
 $\therefore$  There is a root between  $x=2.188$  and  $x=2.125$

Consider  $f\left(\frac{2.188+2.125}{2}\right) = f(2.157) \doteq 0.173$


$\therefore$  There is a root between  $x=2.157$  and  $x=2.125$

Consider  $f\left(\frac{2.157+2.125}{2}\right) = f(2.141) \doteq 0.081$

$\therefore$  There is a root between  $x=2.125$  and  $x=2.141$   
 And  $2.125$  and  $2.141$  are both 2.1 correct to one decimal place.

$\therefore$  No further application of halving the interval is required to find the required root  
 i.e. The root is 2.1.

Q4(a)



From the figure as labelled.  
 $\angle COD = 90^\circ$  [tangent  $\perp$  to radius  
drawn to point of contact]

$$\therefore \angle COD = 180^\circ - (90^\circ + 25^\circ) \quad [\text{sum of } \triangle \text{ is } 180^\circ]$$

$$\therefore \angle CAB = \frac{65^\circ}{2} \quad [\text{angle at centre } 2 \times \text{angle at circumference on same arc}]$$

(b) Consider  $S(n) = 7^{n+1} + 3^n$ Step 1: Let  $n=1$ , then  $S(1) = 7^{1+1} + 3^1 = 49 + 3 = 52$ 

And  $52 = 4 \times 13$   
 $\therefore$  The statement is true for  $n=1$ . — (A)

Step 2: Assume that the statement is true for  $n=k$   
 ie Assume that  $7^{k+1} + 3^k = 4A$  (A an integer)Now consider the statement for  $n=k+1$ .

$$\begin{aligned} \text{Then } S(k+1) &= 7^{k+2} + 3^{k+1} \\ &= 7 \cdot 7^{k+1} + 3^{k+1} \\ &= 4 \cdot 7^{k+1} + 3 \cdot 7^{k+1} + 3 \cdot 3^k \\ &= 4 \cdot 7^{k+1} + 3(7^{k+1} + 3^k) \\ &= 4 + 3 \cdot 4A \text{ using the assumption} \\ &= 4(1 + 3A) \text{ and } 1+3A \text{ is integral.} \end{aligned}$$

$\therefore$  The statement is true for  $n=k+1$  if it is true for  $n=k$  as assumed.


Step 3: But the statement is true for  $n=1$  (from A)

$\therefore$  True for  $n=1+1 = 2$   $\rightarrow$  True for  $2+1 = 3$   
 and so on for all integers  $n$ .

(c) No. of teams possible  
 without restriction =  ${}^6C_3$   
 $= 165$

$$\begin{aligned} \text{No. of teams with mixed sex} &= \text{No. teams } 2M \& 1W + \text{No. teams } 1M \& 2W \\ &= {}^6C_2 \times {}^5C_1 + {}^6C_1 \times {}^5C_2 \\ &= 75 + 60 \\ &= 135. \end{aligned}$$

$$\therefore P(E) = \frac{135}{165} = \frac{9}{11}.$$



Newton's method would not give a better approximation because there is a turning point between  $a$  and  $b$ .

Thus the tangent to the curve at  $a$  will cut the  $x$ -axis further away from  $b$  than  $a$  is.

Section C

5) a) i)  $y = \log_e(\cos x)$

$$y' = \frac{-\sin x}{\cos x}$$

$$y' = -\tan x$$

(ii)  $y = (x+1)e^{-x}$        $u = x+1$        $v = e^{-x}$   
 $u' = 1$        $v' = -e^{-x}$

$$y' = -(x+1)e^{-x} + e^{-x}$$


$$y' = -xe^{-x} - e^{-x} + e^{-x}$$

$$y' = -xe^{-x}$$

(b) i)  $\frac{d(x + \log_e x)}{dx} = 1 + \frac{1}{x}$   
 $= \frac{x+1}{x}$

(iii)  $\int \frac{x+1}{x^2+x\log_e x} dx = \int \frac{x+1}{x(x+\log_e x)} dx$   
 $= \int \frac{\left(\frac{x+1}{x}\right)}{x+\log_e x} dx$   
 in form  $\int \frac{f'(x)}{f(x)} dx$   
 $= \log_e(x + \log_e x) + C$

(c)



$$CD = 6 \text{ cm.}$$

$$AK \cdot KB = CK \cdot KD$$

(product of intercepts)  
 intersecting chords

$$\text{let } CK = x$$

$$1 \times 9 = x(6-x)$$

$$6x - x^2 = 9$$

$$x^2 - 6x + 9 = 0$$

$$(x-3)^2 = 0$$

$$x = 3.$$

In A's  $\triangle AKC \triangle DKB$

$\angle AKC = \angle DKB$  (vertically opposite angles)

$\angle CAK = \angle BDK$  (angles in same segment)

$\therefore \triangle AKC \sim \triangle DKB$  (extriangular)

$\frac{AC}{BD} = \frac{AK}{DK}$  (corresponding sides of similar triangles  
 in same ratio)

$$\frac{AC}{BD} = \frac{1}{3}$$

$$\therefore AC : BD = 1 : 3$$

6) a) i)  $9 \times 8 \times 7 = 504$  or  ${}^3P_3$

(ii)  $\omega$ )  $5 \times 8 \times 7 = 280$

$$\text{Probability} = \frac{280}{504}$$

$$= \frac{5}{9}$$

(B) Given a 3 digit number (of different digits not including zero), only one arrangement will be in ascending order.

$$\text{ie } \frac{1}{3!} = \frac{1}{6} \quad \text{or } \frac{^9C_3}{9P_3}$$

(b) i)  $y = \ln x, x > 0$

$$y' = \frac{1}{x} = x^{-1}$$


$$y'' = -x^{-2}$$

$$= -\frac{1}{x^2}$$

$$y'' < 0 \quad \text{for all } x > 0$$

$\therefore y = \ln x$  is concave down for all  $x > 0$ .

(ii)



(iii)  $A(a, \ln a) \quad B(b, \ln b)$

~~$2 : 1$~~

$$P = \left( \frac{a+2b}{2+1}, \frac{\ln a + 2\ln b}{2+1} \right)$$

$$P = \left( \frac{1}{3}a + \frac{2}{3}b, \frac{1}{3}\ln a + \frac{2}{3}\ln b \right)$$

(iv) Consider the point Q which lies on  $y = \ln x$  with the same x-coordinate as P.

Clearly from the graph in (ii)

$$y_p < y_q$$

$$\frac{1}{3}\ln a + \frac{2}{3}\ln b < \ln \left( \frac{1}{3}a + \frac{2}{3}b \right)$$