



THE SCOTS COLLEGE
2011

PRE TRIAL EXAMINATION
YEAR 12 MATHEMATICS

General Instructions

- Eight questions of equal value
- 5 minutes reading time
- Working time - 2 hours
- Write using blue or black pen
- Board approved calculators may be used
- Start a new booklet for each question
- All necessary working should be shown in every question
- Standard Integrals Table is attached

TOTAL MARKS: 80

WEIGHTING: 30 %

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE: $\ln x = \log_e x, \quad x > 0$

Question 1: 10 Marks

START A NEW PAGE

Marks

a) Find the value of:

i. $\log_3 9$

1

ii. $\log_e 1$

1

b) Using $\log_a 2 = 0.387$ and $\log_a 3 = 0.613$, find the value of $\log_a 12$.

2

c) Differentiate with respect to x :

i. $x^2 e^{2x}$

2

ii. $(e^x - e^{-x})^2$

2

d) By first apply the log laws, differentiate $y = \ln\left(\frac{x+2}{x-1}\right)$

2

END OF QUESTION 1

Question 2: 10 Marks

START A NEW PAGE

a) The diagram below shows the points $A(2, -2)$, $B(-2, -3)$ and $C(0, 2)$ which are the vertices of a triangle ABC.

Marks

i. Show the equation of the line AC in the general form is $2x + y - 2 = 0$


2

ii. Calculate the perpendicular distance of B from the side AC.

2

iii. Find the area of $\triangle ABC$.

3



b) For the parabola $x^2 = 16y$ state:

i. the focal length

1

ii. the coordinates of the focus

1

iii. the equation of the directrix

1

END OF QUESTION 2

Question 3: 10 Marks

START A NEW PAGE


Marks

- a)
 i. Write down the discriminant of $7x^2 + 5x + k$.
 ii. For what values of k does $7x^2 + 5x + k = 0$ have equal roots?

1

1

- b) ABCD is a parallelogram. $\angle AEB = \angle DFC = 90^\circ$



- i) Show that $\triangle ABE \cong \triangle DCF$.
 ii) Show that ABEF is a rectangle.

2

3

- c) Solve for x :

2

$$|1+2x|=7$$

- d) Express $\frac{1}{4+\sqrt{3}}$ with a rational denominator.

1

END OF QUESTION 3

Question 4: 10 Marks

START A NEW PAGE

- a) For what value of x is the tangent to the curve $y = e^{3x}$ parallel to the line $y = 6x$

3

- b) Evaluate:

i. $\int_2^3 e^{2x-4} dx$.

2

ii. $\int_1^3 \frac{x}{x^2+1} dx$.

2

- c) Find $\frac{d}{dx}(\ln x)^2$ and hence evaluate $\int_1^2 \frac{\ln x}{x} dx$.

3

END OF QUESTION 4


Question 5: 10 Marks

START A NEW PAGE

Marks

- a) In the diagram below, XY is an arc of a circle of radius 10 cm and $\angle XOY = \frac{3\pi}{8}$. Find the area of the shaded region correct to the nearest cm^2 .

2



- b) Draw a neat sketch of $y = 3\cos 2x$ for $0 \leq x \leq \pi$. State period, amplitude and label all x intercepts.
- c) State the exact value of $\cos \frac{\pi}{6}$.
- d) A straight road was constructed to cut a dangerous bend on a country road. It was found that the bend was part of an arc of radius 170 metres and the straight road was 250 metres long.

3

1




DIAGRAM NOT DRAWN TO SCALE

- i. Use the cosine rule to find the size of θ correct to the nearest degree.
- ii. Find the distance by which the old road was shortened. Answer correct to the nearest metre.

2

2

END OF QUESTION 5

Question 6: 10 Marks

START A NEW PAGE

- a) For the curve $y = \frac{1}{x}$ in the domain $x > 0$

i. write down the range of this function;

1

ii. use calculus to show that it:

$\alpha.$ has no stationary points;

2

$\beta.$ is always decreasing;

1

$\gamma.$ is always concave up.

2

- b) Calculate the following limit:

$$\lim_{x \rightarrow 3} \frac{2x^4 - 6x^3}{x^2 - 5x + 6}$$

2

- c) Find the exact area of an equilateral triangle that has side lengths 10cm.

2


END OF QUESTION 6

Question 7: 10 Marks

START A NEW PAGE

Marks

- a) The diagram shows the area bounded by the graph $y = \ln x$, the co-ordinate axes and the line $y = \ln 3$.



- i. Find the shaded area. 3
- ii. Hence find the exact value of $\int_1^3 \ln x dx$. 1
- b) Find the co-ordinates of the stationary point on the curve $y = e^{-x^2}$ and determine its nature. 3
- c) Find the volume of the solid formed when the area bounded by the lines $x = 0$ and $x = 1$ and the curve $y = e^x$ is rotated about the x axis. Leave your answer in exact form. 3

END OF QUESTION 7

Question 8: 10 Marks

START A NEW PAGE

Marks

- a) For the curve $y = xe^x$
- i) Find any stationary points and determine their nature. 2
- ii) Find any asymptotes and intercepts. 2
- iii) Sketch the curve. 1
- b) A normal is drawn to the curve $y = e^{2x}$ at the point $P(\log_e 2, 4)$. The normal cuts the x -axis at Q which has co-ordinates $(32 + \log_e 2, 0)$.
- i. Show that the equation of the normal at P is $x + 8y = 32 + \log_e 2$. 2
- ii. Find the area of the region bounded by the curve $y = e^{2x}$, the normal at P and the co-ordinate axes. 3

END OF EXAM

Question 1

$$a) i) \log_3 9 = \frac{\log_e 9}{\log_e 3}$$

$$= 2 \quad \checkmark$$

$$ii) \log_e 1 = 0$$

$$b) \log_a 12 = \log_a(2^2 \times 3)$$

$$= \log_a(2^2 \times 3)$$

$$= \log_a 2^2 + \log_a 3$$

$$= 2 \log_a 2 + \log_a 3$$

$$= 2 \times 0.387 + 0.613$$

$$= 1.387 \quad \checkmark$$

$$c) i) y = x^2 e^{2x} \quad u = x^2 \quad v = e^{2x}$$

$$\frac{dy}{dx} = vu' + uv' \quad u' = 2x \quad v' = 2e^{2x}$$

$$= e^{2x} \times 2x + x^2 \times 2e^{2x}$$

$$= 2xe^{2x} + 2x^2 e^{2x}$$

$$= 2xe^{2x}(1+x) \quad \checkmark$$

$$ii) y = (e^x - e^{-x})^2 \quad \frac{dy}{dx} = \frac{vu' - uv'}{v^2} \quad u = e^{4x} - 2e^{2x} + 1 \quad v = e^{2x}$$

$$= (e^x - \frac{1}{e^x})^2 \quad u' = 4e^{4x} - 4e^{2x} \quad v' = 2e^2$$

$$= \frac{(e^{2x} - 1)^2}{e^x} \quad = \frac{e^{2x} \times (4e^{4x} - 4e^{2x}) - (e^{4x} - 2e^{2x} + 1) \times 2e^2}{(e^{2x})^2}$$

$$= \frac{e^{4x} - 2e^{2x} + 1}{e^{2x}} \quad = \frac{4e^{4x}(e^{2x} - 1) - 2e^{2x}(e^{2x} - 1)^2}{e^{4x}}$$

$$= \frac{e^{2x}(e^{2x} - 1)[4e^{2x} - 2(e^{2x} - 1)]}{e^{4x}}$$

$$= \frac{e^{2x}(e^{2x} - 1)[4e^{2x} - 2e^{2x} + 2]}{e^{4x}}$$

$$= \frac{(e^{2x} - 1)(2e^{2x} + 2)}{e^{2x}}$$

$$= \frac{2(e^{2x} - 1)(e^{2x} + 1)}{e^{2x}}$$

$$= \frac{2(e^{4x} - 1)}{e^{2x}} \quad \checkmark \checkmark$$

$$d) y = \ln \left[\frac{x+2}{x-1} \right] \quad \checkmark$$

$$= \ln(x+2) - \ln(x-1) \quad \checkmark$$

$$\frac{dy}{dx} = \frac{1}{x+2} - \frac{1}{x-1} \quad \checkmark$$

$$= \frac{x-1 - (x+2)}{(x+2)(x-1)}$$

$$= \frac{x-1-x-2}{x^2 - x + 2x - 2}$$

$$= \frac{-3}{x^2 - x - 2}$$

$$= \frac{-3}{(x+2)(x-1)}$$

Question 2

d) i) m of AC = $\frac{y_2 - y_1}{x_2 - x_1}$ A(2, -2) C(0, 2)

$$= \frac{2 - (-2)}{0 - 2}$$

$$= \frac{2 + 2}{-2}$$

$$= \frac{4}{-2}$$

$$= -2$$

Equation of line AC is:

$$y - y_1 = m(x - x_1)$$

$$y - 2 = -2(x - 0)$$

$$y - 2 = -2x$$

$$2x + y - 2 = 0$$

∴ equation of line AC in general form is

$$2x + y - 2 = 0$$

$$2x + y - 2 = 0 \quad \begin{matrix} a=2 \\ b=1 \\ c=-2 \end{matrix}$$

ii) $DI = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$ B(-2, -3)

$$= \frac{|2x - 2 + |x - 3 + -2|}{\sqrt{2^2 + 1^2}}$$

$$= \frac{|-4 - 3 - 2|}{\sqrt{4 + 1}}$$

$$= \frac{|-9|}{\sqrt{5}}$$

$$= \frac{9}{\sqrt{5}} \text{ units or } \frac{9\sqrt{5}}{5} \text{ units}$$

iii) Area of $\triangle ABC = \frac{1}{2} \times \text{base} \times \text{height}$

$$= \frac{1}{2} \times \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \times \frac{9\sqrt{5}}{5}$$

$$= \frac{9\sqrt{5}}{10} \times \sqrt{(2 - 0)^2 + (-2 - 2)^2}$$

$$= \frac{9\sqrt{5}}{10} \times \sqrt{(2^2) + (0)^2}$$

$$= \frac{9\sqrt{5}}{10} \times \sqrt{4}$$

$$= \frac{9\sqrt{5}}{10} \times 2$$

$$= \frac{18\sqrt{5}}{10}$$

$$= \frac{9\sqrt{5}}{5} \text{ units}^2$$

b) i) $x^2 = 16y$ $x^2 = 4ay$

$$4a = 16$$

$$a = \frac{16}{4}$$

$$\therefore a = 4$$

$$\therefore \text{total length} = a = 4$$

ii) focus S is $(0, a) = (0, 4)$

$$\therefore \text{focus } (0, 4)$$

iii) directrix is $y = -a$

$$\therefore \text{directrix } y = -4$$

Question 3

a) i) $7x^2 + 5x + k$

$$\begin{aligned} \text{discriminant } \Delta &= b^2 - 4ac \\ &= 5^2 - 4 \times 7 \times k \\ &= 25 - 28k \end{aligned}$$

ii) for $7x^2 + 5x + k = 0$ to have equal roots

$$\begin{aligned} \Delta &= 0 \\ \Delta &= 25 - 28k \\ 0 &= 25 - 28k \\ 28k &= 25 \\ k &= \frac{25}{28} \end{aligned}$$

\therefore for $7x^2 + 5x + k = 0$ to have equal roots $k = \frac{25}{28}$

b) i) In $\triangle ABE$ & $\triangle DCF$

$AB = DC$ (opp sides of parallelogram =)

$\angle ABE = \angle DCF$ (transversal FB passes through
 2 lines AB & $DC \therefore$ corresponding

$\angle AEB = \angle DCF$ (Both = 90° given)

$\therefore \triangle ABE \cong \triangle DCF$ (ASA)

ii) $AE = DF$ (corresponding sides in $\cong \triangle$'s =)

$BE = CF$ (corresponding sides in $\cong \triangle$'s =)

Let $BE = x \therefore CF = x$, let $EC = y$

$$BE + EC = BC$$

$BC = AD$ (opp sides of parallelogram =)

$$\therefore AD = x + y$$

$$CF + EC = EF$$

$$= x + y$$

however AD also = $x + y$

$$\therefore EF = AD$$

$$\begin{aligned} \angle AEF &= 180^\circ - 90^\circ \text{ (}\angle \text{ sum at straight line} = 180^\circ\text{)} \\ &= 90^\circ \end{aligned}$$

$BC \parallel AD$ (opp sides in parallelogram =)

$\therefore BF \parallel AD$ (BC lies on line BF)

$$\begin{aligned} \therefore \angle ADF &= 180^\circ - 90^\circ \text{ (co-interior } \angle\text{'s} = 180^\circ\text{)} \\ &= 90^\circ \end{aligned}$$

$$\begin{aligned} \angle DAE &= 360^\circ - 90^\circ - 90^\circ - 90^\circ \text{ (}\angle \text{ sum at quadrilateral} \\ &\quad \text{4 sided polygon} = 360^\circ\text{)} \\ &= 90^\circ \end{aligned}$$

$\therefore AEFD$ is a rectangle (opp sides = as shown above
 Call \angle 's = 90° as shown above)

$$c) |1+2x| = 7$$

$$1+2x = 7 \quad \text{or} \quad 1+2x = -7$$

$$2x = 7-1$$

$$2x = -7-1$$

$$x = \frac{6}{2}$$

$$x = \frac{-8}{2}$$

$$x = 3 \checkmark$$

$$x = -4 \checkmark$$

$$\therefore x = 3 \quad \text{or} \quad x = -4$$

$$d) \frac{1}{4+\sqrt{3}} = \frac{1}{4+\sqrt{3}} \times \frac{4-\sqrt{3}}{4-\sqrt{3}}$$

$$= \frac{4-\sqrt{3}}{4^2 - (\sqrt{3})^2}$$

$$= \frac{4-\sqrt{3}}{16-3}$$

$$= \frac{4-\sqrt{3}}{13} \checkmark$$

Question 4

$$a) y = 6x \quad \textcircled{1}$$

m_1 of $\textcircled{1}$ is 6

$$y = e^{3x} \quad \textcircled{2}$$

$$\frac{dy}{dx} = 3e^{3x} \checkmark$$

for the tangent at the curve $y = e^{3x}$ to be parallel to $y = 6x$ m_1 must equal m_2

$$\text{If } m_1 = m_2$$

$$6 = \frac{dy}{dx}$$

$$6 = 3e^{3x} \checkmark$$

$$\frac{6}{3} = e^{3x}$$

$$2 = e^{3x}$$

$$\log_e 2 = 3x$$

$$x = \frac{\ln 2}{3} \checkmark$$

$\therefore x = \frac{\ln 2}{3}$ for $y = e^{3x}$ to be parallel to $y = 6x$

$$b) i) \int_2^3 e^{2x-4} dx = \int_2^3 e^{2x} \cdot e^{-4} dx$$

$$= e^{-4} \int_2^3 e^{2x} dx \checkmark$$

$$= e^{-4} \left[\frac{1}{2} e^{2x} \right]_2^3$$

$$= \frac{1}{2} e^{-4} \left[e^{2x} \right]_2^3$$

$$= \frac{1}{2} e^{-4} \left[e^{2 \times 3} - e^{2 \times 2} \right]$$

over the
area

$$e^6 = (e^3)^2$$

$$= \frac{1}{2} e^{-4} [e^6 - e^4]$$

~~$$= \frac{1}{2} e^{-4} \times e^4 (e^2 - 1)$$~~

$$= \frac{1}{2} e^{-4} (e^6 - e^4) = \frac{1}{2} (e^2 - 1)$$

× Please simplify

$$\text{ii) } \int_1^3 \frac{x}{x^2+1} dx = \int_1^3 \frac{x}{x^2+1} \times \frac{2}{2} dx$$

$$= \frac{1}{2} \int_1^3 \frac{2x}{x^2+1} dx$$

$$= \frac{1}{2} [\ln(x^2+1)]_1^3$$

$$= \frac{1}{2} [\ln(3^2+1) - \ln(1^2+1)]$$

$$= \frac{1}{2} (\ln 10 - \ln 2)$$

$$= \frac{1}{2} (\ln(5 \times 2) - \ln 2)$$

$$= \frac{1}{2} (\ln 5 + \ln 2 - \ln 2)$$

$$= \frac{1}{2} (\ln 5 + 0)$$

$$= \frac{1}{2} \ln 5$$

$$= \frac{\ln 5}{2} \checkmark$$

$$\text{c) } y = (\ln x)^2$$

$$\text{Let } u = \ln x$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$\frac{du}{dx} = \frac{1}{x}$$

$$y = u^2$$

$$\frac{dy}{du} = 2u$$

$$= \frac{1}{x} \times 2u$$

$$= \frac{1}{x} \times 2 \ln x$$

$$= \frac{2 \ln x}{x} \checkmark$$

$$\int_1^2 \frac{\ln x}{x} dx = \int_1^2 \frac{\ln x}{x} \times \frac{2}{2} dx$$

$$= \frac{1}{2} \int_1^2 \frac{2 \ln x}{x} dx$$

$$= \frac{1}{2} [(\ln x)^2]_1^2$$

$$= \frac{1}{2} [(\ln 2)^2 - (\ln 1)^2]$$

$$= \frac{1}{2} [(\ln 2)^2 - 0]$$

$$= \frac{1}{2} (\ln 2)^2$$

$$= \frac{(\ln 2)^2}{2} \checkmark$$

Question 5

a) Area of sector = $\frac{\theta}{2\pi} \times \pi r^2$

$$= \frac{\theta}{2} r^2$$


$$= \frac{3\pi}{8} \times 10^2 \quad \theta = \frac{3\pi}{8}, r = 10$$

$$= \frac{3\pi}{16} \times 100$$

$$= \frac{300\pi}{16}$$

$$\approx 58.904 \dots$$

$$\approx 59 \text{ cm}^2$$



period = $\frac{2\pi}{2}$

$$= \pi$$

\therefore period = π Amplitude = 3

c) $\cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$

d) i) $a^2 = b^2 + c^2 - 2bc \cos A$

$$2bc \cos A = b^2 + c^2 - a^2$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$A = \cos^{-1} \left(\frac{b^2 + c^2 - a^2}{2bc} \right)$$

$$= \cos^{-1} \left(\frac{170^2 + 170^2 - 250^2}{2 \times 170 \times 170} \right)$$

$$= \cos^{-1} (-0.0813145 \dots)$$

$$= 94^\circ 39' 50.94''$$

$$\approx 95^\circ$$

ii) arc length = $\frac{2\pi r \times \theta}{360}$

$$= 2\pi r \times \frac{95^\circ}{360^\circ}$$

$$= 2 \times \pi \times 170 \times \frac{95}{360}$$

$$\approx 280.874 \dots \text{ m}$$

Distance old road was shortened:

$$= \text{Distance old road} - \text{distance new road}$$

$$= 280.874 - 250$$

$$= 30.87418 \dots \text{ m}$$

$$\approx 31 \text{ m}$$

\therefore old road was shortened by

Question 6

a) $y = \frac{1}{x}, x > 0$ X

i) R: $y \neq 0$ as $0 \neq \frac{1}{x}$
~~# $y > 0$~~ ~~$x = \frac{1}{0}$ undefined~~

ii) a) $y = \frac{1}{x}$
 $y = x^{-1}$
 $\frac{dy}{dx} = -x^{-2}$

when $\frac{dy}{dx} = 0$

$0 = -x^{-2}$

$0 = -\frac{1}{x^2}$ //

$x^2 = \sqrt{-\frac{1}{0}}$, undefined

\therefore no stationary points

B) $y = \frac{1}{x}, \frac{dy}{dx} = -x^{-2}$

$\frac{d^2y}{dx^2} = - -2x^{-3}$

~~#~~ $= 2x^{-3}$

$\therefore y'' = \frac{2}{x^3}$ \therefore for all $x > 0$, y'' will get smaller as the denominator x^3 gets bigger

i.e. $y'' = \frac{2}{x^3} \quad y'' = \frac{2}{(1)^3} = 2$

$\therefore y = \frac{1}{x}$ is always decreasing $y'' = \frac{2}{(2)^3} = 0.25$

as $y'' \rightarrow 0$ as $x \rightarrow \infty$ in the $y'' = \frac{2}{(10)^3} = 0.01$

Y) for a function to be concave up

$y'' > 0$

In $y = \frac{1}{x}$

$y'' = \frac{2}{x^3}$


as $x \rightarrow \infty, y'' \rightarrow 0$

since $x > 0$, y will always be > 0

$\therefore y'' > 0$ for all x in the domain

$\therefore y = \frac{1}{x}$ is always concave up.

b) $\lim_{x \rightarrow 3} \frac{2x^4 - 6x^3}{x^2 - 5x + 6} = \lim_{x \rightarrow 3} \frac{2x^4 - 6x^3}{x^2 - 5x + 6}$ (divide by the highest power)
 ~~$= \lim_{x \rightarrow 3} \frac{2x^3(x-3)}{(x-2)(x-3)} = \lim_{x \rightarrow 3} \frac{2 - \frac{6}{x}}{\frac{1}{x^2} - \frac{5}{x} + \frac{6}{x^2}}$~~
 ~~$= \frac{2(3^3)}{3-2} = 54$~~ ~~$= 2$~~ X



~~$\sin 60^\circ = \frac{h}{10}$~~

$h = 10 \sin 60^\circ$

$= 10 \times \frac{\sqrt{3}}{2}$ //

$= 5\sqrt{3}$

\therefore Area of equilateral Δ is ~~$\frac{1}{2}bh$~~

$A = \frac{1}{2}bh$

$= \frac{1}{2} \times 10 \times 5\sqrt{3} = 25\sqrt{3} \text{ cm}^2$

Question 7

d) i) $y = \ln x$
 $x = e^y$

$$\therefore \text{shaded area} = \int_0^{\ln 3} e^y dy$$

$$= [e^y]_0^{\ln 3}$$

$$= [e^{\ln 3} - e^0]$$

$$= [3 - 1]$$

$$= 2 \text{ units}^2 \quad \text{///}$$

ii) $\int_1^3 \ln x dx = \text{Area rectangle } (\ln 3 \times 3) - \text{shaded area}$

$$= \ln 3 \times 3 - 2$$

$$= 3 \ln 3 - 2 \quad \checkmark$$

b) $y = e^{-x^2}$ (1) Stationary points when $\frac{dy}{dx} = 0$

$$\frac{dy}{dx} = -2x e^{-x^2} \quad \checkmark$$

$$= \frac{-2x}{e^{2x^2}}$$

$$\frac{dy}{dx} = \frac{-2x}{e^{2x^2}}$$

$$0 = \frac{-2x}{e^{2x^2}} \quad \checkmark$$


$$0 = -2x$$

$$x = 0 \text{ sub } x = 0 \text{ into (1)}$$

$$y = e^{-0^2} = 1 \quad \checkmark$$

x	0 ⁻	0	0 ⁺
$\frac{dy}{dx}$	+	0	-

\therefore local max at $(0, 1)$



$$y = e^x$$

$$V = \pi \int_0^1 (e^x)^2 dx$$

$$= \pi \int_0^1 e^{2x} dx$$

$$= \pi \int_0^1 e^{2x} dx$$

$$= \pi \left[\frac{1}{2} e^{2x} \right]_0^1$$

$$= \pi \left(\frac{1}{2} e^{2 \times 1} - \frac{1}{2} e^{2 \times 0} \right)$$

$$= \pi \left(\frac{1}{2} e^2 - \frac{1}{2} \right)$$

$$= \frac{\pi}{2} (e^2 - 1) \text{ units}^3 \quad \text{///}$$

Question 8

a) $y = xe^{2x}$

i) $y = xe^{2x}$ (1) $u = x$ $v = e^{2x}$
 $\frac{dy}{dx} = vu' + uv'$ $u' = 1$ $v' = 2e^{2x}$
 $= e^{2x} \cdot 1 + x \cdot 2e^{2x}$ $\frac{d^2y}{dx^2} = e^{2x} + vu'' + uv''$
 $= e^{2x} + 2xe^{2x}$ $= e^{2x} + e^{2x} + 2xe^{2x}$
 $= e^{2x}(1+2x)$ (2) $= 2e^{2x} + 2xe^{2x}$
 $= e^{2x}(2+x)$ (3)

stationary points when $\frac{dy}{dx} = 0$

$0 = e^{2x}(1+x)$

$\therefore e^{2x} = 0$ or $(1+x) = 0$

$\log_e 0 = x$

undefined, \therefore

reject

when $\frac{d^2y}{dx^2} = 0$

$0 = e^{2x}(2+x)$

$\therefore 2+x = 0$
 $x = -2$

$x = -1$ sub into (1)

$y = -1 \times e^{-1}$

$= -\frac{1}{e}$

\therefore p.o.i $x = -2$

$y = 2e^{-2x}$

\therefore at $(-1, -\frac{1}{e})$

x	-1^-	-1	-1^+
$\frac{dy}{dx}$	$-ve$	0	$+ve$

x			
$\frac{d^2y}{dx^2}$			

\therefore local min t.p. at $(-1, -\frac{1}{e})$


i) $y = xe^{2x}$, as $x \rightarrow \infty$ $y \rightarrow \infty$

\therefore asymptote at $y = 0$

when $x = 0$, $y = 0 \times e^0$ \therefore y-intercept at $(0, 0)$

$= 0 \times 1$

iii)



b) $y = e^{2x}$, P $(\log_e 2, 4)$

$y = e^{2x}$
 $\frac{dy}{dx} = 2e^{2x}$

when $\frac{dy}{dx} = 0$, tangent at curve is

$0 = 2e^{2x}$

$0 = e^{2x}$

i) Normal cuts point P and x-axis at Q

$m = \frac{y_2 - y_1}{x_2 - x_1}$

$= \frac{0 - 4}{32 + \ln 2 - \ln 2}$

$= \frac{-4}{32}$

$= -\frac{1}{8}$

Equation of normal at P is

$y - y_1 = m(x - x_1)$

$y - 4 = -\frac{1}{8}(x - \ln 2)$

$8y - 32 = -x + \ln 2$

$x + 8y = 32 + \ln 2$

\therefore equation of normal is

$x + 8y = 32 + \ln 2$

i) Area of region bound by curve $y = e^{2x}$ and normal at P & co-ordinate axis is

$$A = A_1 + A_2$$

$$A = \int_0^{\ln 2} e^{2x} dx + \frac{1}{2}bh$$

$$= \left[\frac{1}{2} e^{2x} \right]_0^{\ln 2} + \left(\frac{1}{2} \times (32 + \ln 2 - \ln 2) \times 4 \right)$$

$$= \left[\frac{1}{2} e^{2 \ln 2} - \frac{1}{2} e^{2 \times 0} \right] + \frac{1}{2} \times 32 \times 4$$

$$= 2 \times 4 - 2 \times 1 + 64$$

$$= 8 - 2 + 64$$

$$= 70 \text{ units}^2$$

