

Question 1 (15 marks)

Marks

a) Evaluate :

i. $3 \lim_{x \rightarrow 0} \frac{\tan x}{x}$ 1

ii. $\lim_{x \rightarrow 0} \frac{\sin 5x}{4x}$ 2


b) Find, correct to the nearest degree, the acute angle between the lines $x + y - 4 = 0$ and $y = 2x + 1$ 4

c) Differentiate with respect to θ

i. $y = 4 \cos^3 \theta - 3 \cos \theta$ 2

ii. $y = \sec \theta - \cot \theta$ 2

d)



PS and QR are arcs of concentric circles with O as centre.
Calculate in terms of π , the perimeter of the shaded region PQR.

4

Question 2 (15 marks) (Start a new page)

Marks

a) Find the exact value of $\tan \frac{5\pi}{6}$, expressing your answer in surd form with a rational denominator. 2

b) i. Write out the expansion of $\sin(x+y)$ 1

ii. Hence show that $\sin(x + \frac{\pi}{2}) = \cos x$ 1

c) Find:

i. $\int_0^{\frac{\pi}{4}} \cos 3x dx$ 2

ii. $\int_0^{\frac{\pi}{2}} \sin^2 2x dx$ 2

d)

i. Express $\sqrt{3} \sin x + \cos x$ in the form $C \sin(x + \alpha)$ where $C > 0$ and $0 \leq \alpha \leq \frac{\pi}{2}$ 2

ii. Hence or otherwise, solve the equation $\sqrt{3} \sin x + \cos x = 1$ for $0 \leq x \leq 2\pi$ 2

e) Prove the identity $\frac{\cos 2\theta}{\sin \theta + \cos \theta} = \cos \theta - \sin \theta$ 3

Question 3 (15 marks) (Start a new page)

Marks

- a) (i) Express $\tan(\alpha - \beta)$ in terms of $\tan \alpha$ and $\tan \beta$. 1
(ii) Use the result from (i) to find the exact value of $\tan 15^\circ$. 3
- b) Let $t = \tan \frac{\theta}{2}$ and starting with $\tan \theta = \tan\left(\frac{\theta}{2} + \frac{\theta}{2}\right)$, show that $\tan \theta = \frac{2t}{1-t^2}$. 3
- c) Use the 't' results to solve $3\cos \theta + 4\sin \theta = -3$ for $0^\circ \leq \theta \leq 360^\circ$ giving your answer correct to the nearest minute. 4
- d) Find the exact value of the volume of solid of revolution formed when the region bounded by curve $y = \sin x$, the x-axis and the line $x = \frac{\pi}{4}$ is rotated about the x-axis. 4

Question 4 (15 marks) (Start a new page)

Marks

- a) Find the equation of the tangent to the curve $y = \frac{1}{2} \cos 4x$ at the point where $x = \frac{\pi}{8}$. 3
- b) A sector of a circle has an area of 24 cm^2 . Find the radius if the angle at the centre of the circle is 3 radians. 2
- c) Solve the equation $\tan^2 2x = 1$ for $0 \leq x \leq \pi$. 3
- d) Solve the equation $2 \cos^2 x - \sin x - 1 = 0$ for $0 \leq x \leq 2\pi$. 3
- e) (i) Draw a neat sketch of $y = \frac{1}{2} \sin 2x + 2$ for $0 \leq x \leq \pi$ showing all important features. 2
(ii) On the same graph sketch $y = 1$. 1
(iii) How many solutions of $\frac{1}{2} \sin 2x + 1 = 0$ are there for $0 \leq x \leq \pi$? 1

S.C.H.S. Ext (1) - Task (2) March 2008

A ✓
B ✓
C ✓
D ✓
E ✓
F ✓
G ✓
H ✓

Question 1 (15 marks)

a) i) $3 \lim_{x \rightarrow 0} \frac{\tan x}{x} = 3 \times 1 = 3$ (1)

(ii) $\lim_{x \rightarrow 0} \frac{\sin 5x}{4x} = \frac{5}{4} \lim_{x \rightarrow 0} \frac{\sin 5x}{5x} = \frac{5}{4}$ (2)

b) $x+y-4=0$ $y=2x+1$
 $y=-x+4$ $m_2=2$
 $m_1=-1$

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \left| \frac{-1 - 2}{1 + (-1 \times 2)} \right| = \left| \frac{-3}{1 - 2} \right|$$

$\tan \theta = 3$
 $\therefore \theta = 71^\circ 34'$
 $\approx 72^\circ$ (4)

c) i) $y = 4 \cos^3 \theta - 3 \cos \theta$

$$\frac{dy}{d\theta} = 4 \cdot 3 \cos^2 \theta \cdot (-\sin \theta) + 3 \sin \theta = 3 \sin \theta - 12 \sin \theta \cos^2 \theta$$
 (2)

(ii) $y = \sec \theta - \cot \theta = \frac{1}{\cos \theta} - \frac{\cos \theta}{\sin \theta}$

c) (ii) $y = (\cos \theta)^{-1} - (\tan \theta)^{-1}$

$$\frac{dy}{d\theta} = -1(\cos \theta)^{-2} \cdot (-\sin \theta) + 1(\tan \theta)^{-2} \times \sec^2 \theta$$

$$= \frac{\sin \theta}{\cos^2 \theta} + \frac{\sec^2 \theta}{\tan^2 \theta}$$

$$= \frac{\sin \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\sin^2 \theta \cdot \cos^2 \theta}$$

$$= \frac{\sin \theta}{\cos^2 \theta} + \frac{1}{\sin^2 \theta} = \tan \theta \cdot \sec \theta + \operatorname{cosec}^2 \theta$$
 (2)

d) $\theta = 60^\circ = \frac{\pi}{3}$

length of arc PS = $r\theta = 15 \times \frac{\pi}{3} = 5\pi$ cm
 length of arc OR = $r\theta = 25 \times \frac{\pi}{3} = \frac{25\pi}{3}$ cm

$PQ = SR = 25 - 15 = 10$ cm

\therefore Perimeter of Shaded Region = $5\pi + \frac{25\pi}{3} + 20 = \frac{40\pi}{3} + 20$ cm.

(4)

$$Q_2 \text{ a) } \tan\left(\frac{5\pi}{6}\right) = \tan\left(\pi - \frac{\pi}{6}\right)$$

$$= -\tan\frac{\pi}{6}$$

$$= -\frac{1}{\sqrt{3}} \quad (1\frac{1}{2} \text{ marks})$$

$$= -\frac{\sqrt{3}}{3} \quad (2 \text{ marks})$$

$$\text{b) i) } \sin(x+y) = \sin x \cos y + \cos x \sin y \quad (1 \text{ mark})$$

$$\begin{aligned} \text{ii) } \sin\left(x + \frac{\pi}{2}\right) &= \sin x \cos\frac{\pi}{2} + \cos x \sin\frac{\pi}{2} \\ &= \sin x \cdot 0 + \cos x \cdot 1 \\ &= \cos x \end{aligned} \quad (1 \text{ mark})$$

$$\text{c) i) } \int_0^{\frac{\pi}{4}} \cos 3x \, dx = \left[\frac{1}{3} \sin 3x \right]_0^{\frac{\pi}{4}}$$

$$= \frac{1}{3} \sin \frac{3\pi}{4} - 0$$

$$= \frac{1}{3} \times \frac{1}{\sqrt{2}}$$

$$= \frac{1}{3\sqrt{2}} \quad (1\frac{1}{2} \text{ marks})$$

$$= \frac{\sqrt{2}}{6} \quad (2 \text{ marks})$$

$$\begin{aligned} \text{ii) } \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \frac{1}{2} \int_0^{\frac{\pi}{2}} (1 - \cos 4x) \, dx \\ &= \frac{1}{2} \left[x - \frac{1}{4} \sin 4x \right]_0^{\frac{\pi}{2}} \quad (1 \text{ mark}) \end{aligned}$$

$$= \frac{1}{2} \left[\frac{\pi}{2} - 0 \right] - [0]$$

$$= \frac{\pi}{4} \quad (2 \text{ marks})$$

$$\text{d) I) } C \sin(x+\alpha) = C (\sin x \cos \alpha + \cos x \sin \alpha)$$

$$\sqrt{3} \sin x + \cos x = C \cos \alpha \sin x + C \sin \alpha \cos x$$

$$\therefore C \cos \alpha = \sqrt{3} \quad \text{--- (1)}$$

$$C \sin \alpha = 1 \quad \text{--- (2)}$$

$$C^2 (\sin^2 \alpha + \cos^2 \alpha) = 3 + 1$$

$$C^2 = 4$$

$$C = 2 \quad C > 0$$

$$\tan \alpha = \frac{1}{\sqrt{3}}, \quad \alpha = \frac{\pi}{6}$$

$$\sqrt{3} \sin x + \cos x = 2 \sin\left(x + \frac{\pi}{6}\right) \quad (2 \text{ marks})$$

$$\text{II) } 2 \sin\left(x + \frac{\pi}{6}\right) = 1$$

$$\sin\left(x + \frac{\pi}{6}\right) = \frac{1}{2} \quad (1 \text{ mark})$$

$$x + \frac{\pi}{6} = \frac{\pi}{6}, \quad \pi - \frac{\pi}{6}, \quad \frac{13\pi}{6}$$

$$x = 0, \quad \frac{2\pi}{3}, \quad 2\pi \quad (2 \text{ marks})$$

e)

$$\frac{\cos 2\theta}{\sin \theta + \cos \theta}$$

$$= \frac{\cos^2 \theta - \sin^2 \theta}{\sin \theta + \cos \theta} \quad \leftarrow (1 \text{ mark})$$

$$= \frac{(\cos \theta + \sin \theta)(\cos \theta - \sin \theta)}{(\sin \theta + \cos \theta)} \quad (2 \text{ marks})$$

$$= \cos \theta - \sin \theta \quad (3 \text{ marks})$$

Q3 a) i) $\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$

ii) $\tan 15^\circ = \tan(45^\circ - 30^\circ)$
 $= \frac{\tan 45^\circ - \tan 30^\circ}{1 + \tan 45^\circ \tan 30^\circ}$

Or:
 $\sqrt{3}(60^\circ - 45^\circ)$ etc.
 $= \frac{1 - \frac{1}{\sqrt{3}}}{1 + 1 \times \frac{1}{\sqrt{3}}} \times \frac{\sqrt{3}}{\sqrt{3}}$
 $= \frac{(\sqrt{3}-1) \times (\sqrt{3}-1)}{\sqrt{3}+1 (\sqrt{3}-1)}$
 $= \frac{4 - 2\sqrt{3}}{2}$
 $= 2 - \sqrt{3}$

b) $\tan \theta = \tan\left(\frac{\theta}{2} + \frac{\theta}{2}\right)$
 $= \frac{\tan \frac{\theta}{2} + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2} \cdot \tan \frac{\theta}{2}}$
 $= \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}}$
 $= \frac{2t}{1 - t^2}$

c) $3 \cos \theta + 4 \sin \theta = -3$ for $0^\circ \leq \theta \leq 360^\circ$

$\sin \theta = \frac{2t}{1+t^2}$, $\cos \theta = \frac{1-t^2}{1+t^2}$

$= \frac{3(1-t^2)}{1+t^2} + \frac{4(2t)}{1+t^2} = -3$

$3(1-t^2) + 8t = -3(1+t^2)$

$3 - 3t^2 + 8t = -3 - 3t^2$

$8t = -6$

$t = -\frac{3}{4}$

$\therefore \tan \frac{\theta}{2} = -\frac{3}{4}$ for $0^\circ \leq \theta \leq 180^\circ$

$\therefore \frac{\theta}{2}$ is an obtuse angle

$\frac{\theta}{2} = \tan^{-1}\left(-\frac{3}{4}\right)$

$= -36^\circ 52'$

$\therefore \theta = 117^\circ 04'$

but test $\theta = 180^\circ$ as 't' method does not allow for such a solution.

When $\theta = 180^\circ$

LHS = $3 \cos 180^\circ + 4 \sin 180^\circ$

$= 3 \times -1 + 0$

$= -3 \therefore$ (True)

$=$ RHS

\therefore solutions are $180^\circ, 256^\circ 16'$

d) $V = \pi \int_a^b y^2 dx$

where $y = \sin x$

$\therefore y^2 = \sin^2 x$

$\therefore V = \pi \int_0^{\frac{\pi}{4}} \sin^2 x dx$

but $\sin^2 x = \frac{1}{2}(\cos 2x + 1)$

$\therefore V = \pi \times \frac{1}{2} \int_0^{\frac{\pi}{4}} (\cos 2x + 1) dx$

$= \frac{\pi}{2} \left[\frac{1}{2} \sin 2x + x \right]_0^{\frac{\pi}{4}}$

$= \frac{\pi}{2} \times \left(\frac{1}{2} \sin \frac{\pi}{2} + \frac{\pi}{4} \right)$

$= \frac{\pi}{2} \left(\frac{1}{2} + \frac{\pi}{4} \right)$

$= \frac{\pi}{4} + \frac{\pi^2}{8}$ (u³)

$= \frac{\pi^2}{8} - \frac{\pi}{4}$

Question 4

a) $y = \frac{1}{2} \cos 4x$
 $\frac{dy}{dx} = -2 \sin 4x$ ✓
 at $x = \frac{\pi}{8}$, $m = -2 \sin \frac{\pi}{2} = -2$ ✓

$y = \frac{1}{2} \cos \frac{\pi}{2} = 0$ ✓
 Eqn of tangent
 $y - 0 = -2(x - \frac{\pi}{8})$
 $y = -2x + \frac{\pi}{4}$ #

[3]

b) $A = \frac{1}{2} r^2 \theta$ ✓, $A = 24$, $\theta = 3$
 $24 = \frac{3}{2} r^2$

$r^2 = 16$
 $r = 4$ cm ✓

[2]

c) $\tan^2 2x = 1$, $0 \leq x \leq \pi$
 $\tan 2x = \pm 1$ ✓, $0 \leq 2x \leq 2\pi$
 $2x = \frac{\pi}{4}, (\pi + \frac{\pi}{4}), (\pi - \frac{\pi}{4}), (2\pi - \frac{\pi}{4})$

$2x = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{3\pi}{4}, \frac{7\pi}{4}$ ✓
 $x = \frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}$ ✓ #


[3]

d) $2 \cos^2 x - \sin x - 1 = 0$, $0 \leq x \leq 2\pi$
 $2(1 - \sin^2 x) - \sin x - 1 = 0$ ✓
 $2 - 2 \sin^2 x - \sin x - 1 = 0$
 $-2 \sin^2 x - \sin x + 1 = 0$
 $2 \sin^2 x + \sin x - 1 = 0$
 $(2 \sin x - 1)(\sin x + 1) = 0$
 $\sin x = \frac{1}{2}$ or $\sin x = -1$ ✓

$x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$ ✓ #

[3]

e) i) $y = \frac{1}{2} \sin 2x + 2$
 amp = $\frac{1}{2}$, period = $\frac{2\pi}{2} = \pi$, Vertical shift +2 ✓



[2]

[1]

i) Award 1st mark for correct amp, period, vshift
 Award 1 mark for any two of above correct
 Award second mark for correct sketch with correct amp, period, v. shift. OR 1 mark for incorrect sketch (wrong period, pamp, etc of equivalent difficulty)

ii) as above

iii) $\frac{1}{2} \sin 2x + 1 = 0$

ie $\frac{1}{2} \sin 2x + 2 = 1$
 this is solved by finding the intersection of $y = \frac{1}{2} \sin 2x + 2$ and $y = 1$.

As the two graphs do not intersect there are NO solutions
 [No carry through, here as it changes the intent of the question] ✓

[1]

Total [15]