

SYDNEY GIRLS HIGH SCHOOL



2007 HSC Assessment Task 2

March 7, 2007

MATHEMATICS Extension 1

Year 12

Time allowed: 75 minutes

Topics: Trigonometry (I & II) First part of Polynomials

DIRECTIONS TO CANDIDATES:

- Attempt all questions
- Questions are not of equal value
- There are 5 questions with part marks shown
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work
- Board approved calculators may be used
- Write on one side of the paper only

QUESTION 1

Marks

a) Find $\frac{dy}{dx}$ if

5

(i) $y = \sin 4x$

(ii) $y = \cos\left(\frac{1}{x}\right)$

(iii) $y = \tan^6 x$

b) Find

6

(i) $\int 3 \sin 3x \, dx$

(ii) $\int \sec^2 4x \, dx$

(iii) $\int (\cos 2x + 2 \cos x) \, dx$

(iv) $\int \sin(4x - 3) \, dx$

c) Given that $\cos 36^\circ = \frac{\sqrt{5}+1}{4}$

4

find in simplest form, the exact value of $\cos 72^\circ$

QUESTION 2

- a) Given the lines $L_1 : 3y = x + 1$ and $L_2 : 3x - 4y = 12$,
Show that the acute angle formed by them is equal to
the acute angle formed by L_1 and the x-axis. 5

- b) If $\tan \frac{\theta}{2} = \frac{1}{2}$ find the exact values of 6
- (i) $\tan \theta$
- (ii) $\cos 2\theta$

- c) (i) Factorise $3x^3 + 3x^2 - x - 1$ 4

(ii) hence solve the equation

$$3 \tan^3 \theta + 3 \tan^2 \theta - \tan \theta - 1 = 0 \text{ for } 0 \leq \theta \leq \pi$$

QUESTION 3

- a) Sketch the polynomial $f(x) = x^2(x+1)(x+2)$ 2
(without using calculus)

- b) For the polynomial $P(x) = 2 + x - 5x^2 + 8x^4$ 3
state the
- (i) degree
- (ii) leading term
- (ii) constant term

- c) (i) Find the remainder when $ax^2 + bx + c$ is divided by $x - 1$ 10
- (ii) Under what conditions is $x - 1$ a factor of the quadratic?
- (iii) Show that $x - 1$ is a factor of $ax^3 + (b - a)x^2 + (c - b)x - c$
and find the other factor
- (iv) State necessary and sufficient conditions on a, b, c for the
cubic to have three real and distinct roots

QUESTION 4

- a) Find the volume of the solid of revolution obtained 3
 by revolving the area between $y = 2 \sec x$ and the x-axis
 between $x = 0$ and $x = \frac{\pi}{3}$ about the x-axis

- b) (i) Express $\sqrt{2} \sin x + \sqrt{2} \cos x$ in the form 5
 $R \sin(x + \alpha)$ where $R > 0$ and $0 \leq \alpha \leq \frac{\pi}{2}$

- (ii) Hence, sketch $y = \sqrt{2} \sin x + \sqrt{2} \cos x$ for $0 \leq x \leq 2\pi$
 (show intercepts and endpoints clearly)

- (iii) Hence, find the value(s) of k for which $\sqrt{2} \sin x + \sqrt{2} \cos x = k$
 has 3 solutions in the domain $0 \leq x \leq 2\pi$

- c) Express the solution of the equation $\sin 2\theta = \sin \theta$ 4
 in general form, if θ is in radians


- (d) Find $\int \cos^2 \theta \, d\theta$ 3

QUESTION 5

- a) Evaluate $\lim_{x \rightarrow 0} \frac{\sin 2x}{3x}$ 2

- b) In $\triangle AOB$, $OA = OB = a$, which is constant. 6
 $\angle AOB = x$ radians, where x is variable.

PQ is a circular arc, centre O and radius r . If the
 area of $\triangle AOB$ is twice that of the sector OPQ



- (i) Express r^2 in terms of a and x
 (ii) Find r in terms of a when $\angle AOB$ is a right angle
 (iii) Describe the behaviour of r as $x \rightarrow 0$

- c) A monic polynomial $P(x)$ of degree 4 is known to have 7
 zeros 2 and -2

- (i) Write down an equation for $P(x)$ to the extent
 specified so far.
 (ii) given further that $P(0) = 4$ and $P(1) = -3$, find $P(x)$

- (iii) Solve $P(x) = 0$ for real roots

Q (1) (b) (i) $y' = 4 \cos 4x$ (1)

(ii) $y = \cos(\frac{1}{x})$ let $u = \frac{1}{x}$

$\frac{dy}{du} = -\sin u$ $\frac{du}{dx} = -\frac{1}{x^2}$

$\frac{dy}{dx} = \frac{1}{x} \sin(\frac{1}{x})$ (2)

(iii) $y = (\tan x)^6$

$y' = 6(\tan x)^5 \times \sec^2 x$
 $= 6 \tan^5 x \sec^2 x$ (2)

(b) (i) $\int 3 \sin 3x dx$
 $= 3 \int \sin 3x dx$
 $= 3(-\frac{1}{3} \cos 3x) + C$
 $= -\cos 3x + C$ (2)

(ii) $\int \sec^2 4x dx = \frac{1}{4} \tan 4x + C$ (1)

(iii) $\int (\cos 2x + 2 \cos x) dx$
 $= \frac{1}{2} \sin 2x + 2 \sin x + C$ (2)

(iv) $\int \sin(4x-3) dx$
 $= -\frac{1}{4} \cos(4x-3) + C$ (1)

(c) $\cos 72^\circ = \cos(2 \times 36^\circ)$
 $= 2 \cos^2 36^\circ - 1$
 $= 2 \left(\frac{6+2\sqrt{5}}{16} \right) - 1$
 $= \frac{6+2\sqrt{5}}{8} - \frac{8}{8}$
 $= \frac{2\sqrt{5}-2}{8}$
 $= \frac{\sqrt{5}-1}{4}$ (4)

Q (2) (a) $L_1: y = \frac{1}{3}x + \frac{1}{3}$
 $L_2: y = \frac{3}{4}x - 3$
 $\therefore M_1 = \frac{1}{3}, M_2 = -\frac{3}{4}$


$\tan(\theta_2 - \theta_1) = \frac{M_2 - M_1}{1 + M_1 M_2}$
 $= \frac{\frac{3}{4} - \frac{1}{3}}{1 + \frac{3}{4} \times \frac{1}{3}}$
 $= \frac{\frac{9}{12} - \frac{4}{12}}{1 + \frac{1}{4}}$
 $= \frac{\frac{5}{12}}{\frac{5}{4}}$
 $= \frac{1}{3}$
 $= M_1$ (5)
 $= \tan \theta_1$

(b) (i) $\tan \theta = \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}}$
 $= \frac{2 \times \frac{1}{2}}{1 - \frac{1}{4}}$
 $= \frac{4}{3}$ (3)

Q (2) (b) $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$
 $= \frac{(1-t^2)}{(1+t^2)} - \frac{(2t)}{(1+t^2)}$
 $= \frac{(1-\frac{1}{4})}{(1+\frac{1}{4})} - \frac{(2 \times \frac{1}{2})}{(1+\frac{1}{4})}$
 $= \frac{9}{25} - \frac{16}{25}$
 $= -\frac{7}{25}$ (3)

(c) (i) $3x^2(x+1) - 1(x+1)$
 $= (x+1)(3x^2-1)$ (1)

(ii) let $x = \tan \theta$ $0 \leq \theta < \pi$
 $\therefore (\tan \theta + 1)(3 \tan^2 \theta - 1) = 0$
 $\tan \theta = -1$ or $\tan^2 \theta = \frac{1}{3}$
 $\theta = \frac{3\pi}{4}$ $\tan \theta = \pm \frac{1}{\sqrt{3}}$
 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}$
 $\therefore \theta = \frac{\pi}{6}, \frac{3\pi}{4}, \frac{5\pi}{6}$ (3)



- (b) (i) 4 (1)
- (ii) $8x^4$ (1)
- (iii) 0 (1)

(c) (i) $f(x) = a + b + c$ (1)

(ii) $a + b + c = 0$ (1)

(iii) $f(x) = a + (b-a)x + (c-b)x^2 - c$
 $= 0$
 $\therefore x-1$ is a factor (2)

$ax^2 + bx + c$
 $(x-1)(ax^2 + (b-a)x + (c-b)) - c$
 $ax^2 - ax^2$
 $bx^2 + (c-b)x$
 $bx^2 - bx$
 $cx - c$
 $cx - c$ (4)

\therefore other factor $ax^2 + bx + c$

(iv) $ax^2 + bx + c$ has 2 real distinct roots if $\Delta > 0$
 i.e. $b^2 - 4ac > 0$
 and also $a + b + c \neq 0$ (2)
 i.e. $x-1$ is not a factor of $ax^2 + bx + c$

$$4a) V = \pi \int_{\pi/3}^{\pi/2} y^2 dx$$

$$= \pi \int_{\pi/3}^{\pi/2} 4 \sec^2 x \cdot dx$$

$$= 4\pi \left[\tan x \right]_{\pi/3}^{\pi/2}$$

$$= 4\pi \cdot \sqrt{3} \quad \therefore \text{Volume} = 4\sqrt{3}\pi \text{ u}^3$$

$$b) i) \sqrt{2} \sin x + \sqrt{2} \cos x = R \sin(x+\alpha)$$


$$R \sin(x+\alpha) = R \sin x \cos \alpha + R \cos x \sin \alpha$$

$$\therefore R \cos \alpha = \sqrt{2}, R \sin \alpha = \sqrt{2}$$

$$\therefore \tan \alpha = 1, \alpha = \pi/4$$

$$\therefore R \cdot \frac{1}{\sqrt{2}} = \sqrt{2}, R = 2$$

$$\therefore y = 2 \sin(x + \frac{\pi}{4})$$



iii) 3 solutions if $k = \sqrt{2}$.

$$c) \sin 2\theta = \sin \theta$$

$$\therefore 2 \sin \theta \cos \theta = \sin \theta$$

$$\therefore \sin \theta (2 \cos \theta - 1) = 0$$

$$\therefore \sin \theta = 0, \cos \theta = 1/2$$

$$\therefore \theta = n\pi, 2n\pi \pm \pi/3$$

$$d) \int \cos^2 \theta \cdot d\theta = \frac{1}{2} \int (1 + \cos 2\theta) d\theta$$


$$= \frac{1}{2} (\theta + \frac{1}{2} \sin 2\theta) + C$$

$$= \theta/2 + \frac{1}{4} \sin 2\theta + C$$

$$Q5 a) \lim_{x \rightarrow 0} \frac{\sin 2x}{3x}$$

$$\text{as } x \rightarrow 0, \sin 2x \rightarrow 2x$$

$$\therefore \lim_{x \rightarrow 0} \frac{\sin 2x}{3x} \Rightarrow \frac{2x}{3x} = 2/3$$



$$i) \text{Area } \triangle AOB = \frac{1}{2} a^2 \sin \alpha$$

$$\text{Area } POQ = \frac{1}{2} r^2 \alpha$$

$$\therefore \frac{1}{2} a^2 \sin \alpha = r^2 \alpha$$

$$\therefore r^2 = \frac{a^2 \sin \alpha}{2\alpha}$$

$$ii) \angle AOB = \pi/2, r^2 = \frac{a^2 \cdot 2}{2 \cdot \pi} = \frac{a^2}{\pi}$$

$$\therefore r = \frac{a}{\sqrt{\pi}}$$

$$iii) \text{as } x \rightarrow 0, \frac{\sin x}{x} \rightarrow 1$$

$$\therefore x^2 \rightarrow a^2/2$$

$$r \rightarrow a/\sqrt{2}$$

$$c) i) P(x) = (x+2)(x-2)(x^2 + bx + c)$$

$$P(0) = 4 \quad \therefore 4 = -4 \cdot c \quad c = -1$$

$$P(1) = -3 \quad \therefore -3 = (3)(-1)(1 + b + c)$$

$$\therefore b + c + 1 = +1$$

$$\therefore b - 1 + 1 = +1, b = 1$$

$$\therefore P(x) = (x+2)(x-2)(x^2 + x - 1)$$

$$\therefore \text{Roots are } x = \pm 2, \frac{-1 \pm \sqrt{5}}{2}$$