

Sydney Girls High School



2003

MATHEMATICS EXTENSION 1

Year 11

Assessment Task 1

Time Allowed - 75 minutes

Instructions

- There are four questions.
- Attempt ALL questions.
- The questions are of equal value.
- All necessary working should be shown in every question.
- Marks may be deducted for careless or badly arranged work.
- Start each question on a new page.
- Write on one side of the paper only

Total 80 marks

Question 2. (20 marks)

Marks


1. Integrate the following

(i) $\int 6x^2 + \frac{4}{x^3} dx$ 2


(ii) $\int \frac{dx}{\sqrt{6x+1}}$ 2

2. Evaluate: (i) $\int_{-2}^{-1} (x - \frac{1}{x})^2 dx$ 2

(ii) $\int_0^4 \sqrt{t}(4-t)dt$ 2



4. A bowl is formed by rotating the part of the curve $y = \frac{x^2}{4}$ between $x = 0$ and $x = 2$ about the y axis.
Find the volume of the bowl (exact answer).



5.

4

x	1	1.5	2	2.5	3
$f(x)$	5	1	-2	3	7

Use Simpson's Rule with five function values to evaluate $\int_1^3 f(x)dx$

Question 4 (20 marks)

Marks

1. Find the values of k for which the equation


$$x^2 + (k+2)x + 4 = 0 \text{ has}$$

- (i) equal roots.
- (ii) real and distinct roots.


2
2

2. Find the volume of the solid of revolution formed by rotation of the curve

$$\frac{x^2}{2} + y^2 = 8 \text{ about the } x \text{ axis.}$$



3.



This diagram shows the cross section of a creek with the depths shown in metres at 4 metre intervals. The total width of the creek is 12 metres.

- (i) Use the trapezoidal rule to find an approximate value for the area of the cross section.


4

- (ii) Water flows through this section at a speed of 0.5 metres per second. Calculate the approximate volume of water that flows past this section in an hour.

2

4. (i) Find the coordinates of point A.
 (ii) Find the shaded area (to 1 decimal place).
 (iii) If this shaded area is rotated around the y axis, find the volume of revolution correct to 1 decimal place.

6



Question 1.

Maths Solutions

Year 2003 Exam Paper

$$1 \quad x^2 - 2x - 8 = 0$$

$$(x-4)(x+2) = 0$$

$$\therefore x = 4 \text{ and } x = -2$$

(2)

$$3 \quad 2x^2 + 7x - 3 = 0$$

$$i \quad \alpha + \beta = -b/a = -7/2 = -3\frac{1}{2}$$

$$ii \quad \alpha\beta = c/a = -3/2 = -1\frac{1}{2}$$

$$iii \quad \alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2(\alpha\beta)$$

$$= 12\frac{1}{4} + 3$$

$$= 15\frac{1}{4}$$

(3)

$$5 \quad -2x^2 + 3x + 14 > 0$$


$$\text{Put } -2x^2 + 3x + 14 = 0$$

$$\therefore 2x^2 - 3x - 14 = 0$$

$$(2x-7)(x+2) = 0$$

$$x = 3\frac{1}{2} \text{ and } -2$$

$$\text{Sketch } y = -2x^2 + 3x + 14$$



\therefore Solution to $-2x^2 + 3x + 14 > 0$

$$\text{is } -2 < x < 3\frac{1}{2} = \text{Ans}$$

$$7 \quad 4x^2 - 20x + C = 0$$

Let roots be $\alpha, (\alpha-2)$

$$\alpha + (\alpha-2) = -b/a = 20/4 = 5$$

$$2\alpha = 7 \quad \therefore \alpha = 3\frac{1}{2}$$

$$\alpha(\alpha-2) = 4\alpha = 14$$

$$\therefore C = 4\alpha(\alpha-2)$$

$$= 4 \times 7/2 \times 3/2$$


$$= 21$$

(2)

Question 2.

Year 2003 Exam Paper

$$2 \quad y = x^2 - 6$$



(3)

Question 2

$$i \quad \int 6x^2 + 4x^{-3} dx$$

$$= \frac{6x^3}{3} + \frac{4x^{-2}}{-2} + C$$

$$= 2x^3 - \frac{2}{x^2} + C \quad (2)$$

$$ii \quad \int (6x+1)^{-\frac{1}{2}} dx$$

$$= \frac{(6x+1)^{\frac{1}{2}}}{6 \times \frac{1}{2}} + C$$

$$= \frac{\sqrt{6x+1}}{3} + C \quad (2)$$

Total = 20

4 For positive definite
 $k > 0$ and $\Delta < 0$

$$\begin{aligned}\Delta &= b^2 - 4ac \\ &= 36 - 4 \times k \times 2 \\ &= 36 - 8k\end{aligned}$$

$$36 - 8k < 0$$

$$36 < 8k$$

$\therefore k > 4\frac{1}{2}$ for pos. definite

$$i \quad \int_{-2}^{-1} x^2 - 2 + x^{-2} dx$$

$$= \left[\frac{x^3}{3} - 2x + \frac{x^{-1}}{-1} \right]_{-2}^{-1}$$

$$= \left[\frac{x^3}{3} - 2x - \frac{1}{x} \right]_{-2}^{-1}$$

$$= \left(-\frac{1}{3} + 2 + 1 \right) - \left(-\frac{8}{3} + 4 + \frac{1}{2} \right)$$

$$= \frac{5}{6} \quad (1)$$

$$ii \quad \int_0^4 4t^{\frac{1}{2}} - t^{3/2} dt$$

$$= \left[\frac{2}{3} \cdot 4t^{3/2} - \frac{2}{5} t^{5/2} \right]_0^4$$

$$= \left[\frac{8}{3} t^{5/2} - \frac{2}{5} t^{5/2} \right]_0^4$$

$$= \left(\frac{8}{3} \times 4^{5/2} - \frac{2}{5} \times 16^{5/2} \right) - (0)$$

$$= 8\frac{8}{15} \quad (2)$$

$$5 \quad 3x^2 + 2x + k = 0$$


For no real roots $\Delta < 0$

$$b^2 - 4ac < 0$$

$$4 - 4 \times 3 \times k < 0$$

$$4 < 12k$$

$\therefore k > \frac{1}{3}$ for no real roots



$$i \quad 2x = 6x - x^2$$


$$x^2 - 4x = 0$$

$$x(x-4) = 0$$

$$\therefore x = 0 \text{ and } x = 4$$

$$y = 0 \quad y = 8$$

$$\text{Ans } (0,0) \text{ and } (4,8) \quad (2)$$



$$\begin{aligned}\text{Vol} &= \pi \int_0^1 x^2 dy \\ &= \pi \int_0^1 4y dy\end{aligned}$$

$$\begin{aligned}&= \pi [2y^2]_0^1 \\ &= 2\pi \text{ units}^3 \quad (4)\end{aligned}$$

$$8 \quad x^4 = 4(x^2 + 8)$$

$$\text{Let } a = x^2$$

$$a^2 = 4(a+8)$$

$$a^2 - 4a - 32 = 0$$

$$(a-8)(a+4) = 0$$

$$a = 8 \quad \text{and} \quad a = -4$$

$$x^2 = 8 \quad x^2 = -4$$

$$\therefore \text{No solution}$$

$$\therefore x = \pm \sqrt{8} \quad \text{or} \quad \pm 2\sqrt{2}$$

$$ii \quad \text{Area} = \int_0^4 6x - x^2 - 2x dx$$

$$= \int_0^4 4x - x^2 dx$$

$$= \frac{4x^2}{2} - \frac{x^3}{3}$$

$$= \left[2x^2 - \frac{x^3}{3} \right]_0^4$$

$$= (32 - \frac{64}{3}) - (0)$$

$$= 10\frac{2}{3} \text{ units}^2 \quad (2)$$

x	f(x)	w	wf(x)
1	5	1	5
1.5	1	4	4
2	-2	2	-4
2.5	3	4	12
3	7	1	7
$\sum w f(x)$			24


$$\int_1^3 f(x) dx = \frac{1}{3} \sum w f(x)$$


$$= \frac{0.5}{3} \times 24$$

(4)

Question 3

$$\begin{aligned} i & x^2 + 4x + 4 + y^2 + 8y + 16 = -11 + 20 \\ & (x+2)^2 + (y+4)^2 = 9 \\ & \text{Centre} = (-2, -4) \quad \text{Radius} = 3 \end{aligned}$$

- 2 i  $V = (0,0)$
 $y = x^2$ (6)
- ii Focal length $= 2$ ($= a$)
- iii Equation $x^2 = 4ay$
 $\therefore x^2 = 8y$



Let $P = (x, y)$ = point on locus

Condition: $PA = 2 \times PB$

$$\begin{aligned} \sqrt{(x-1)^2 + (y-5)^2} &= 2 \times \sqrt{(x-4)^2 + (y+1)^2} \\ x^2 - 2x + 1 + y^2 - 10y + 25 &= 4((x-4)^2 + (y+1)^2) \\ 0 &= 3x^2 + 3y^2 - 30x + 18y + 42 \\ \therefore x^2 + y^2 - 10x + 6y + 14 &= 0 \quad (\text{locus eqn}) \end{aligned}$$

- 3 i Let $P = (x, y)$ = point on locus
 Condition: $PA \perp PO$ ($m_1 \cdot m_2 = -1$)


$$\frac{y}{x} \cdot \frac{y}{x-4} = -1$$


$y^2 = -x^2 + 4x$
 $\therefore x^2 + y^2 - 4x = 0$ is locus which

$$\begin{aligned} ii & x^2 - 4x + y^2 = 0 \\ & x^2 - 4x + 4 + y^2 = 4 \\ & (x-2)^2 + y^2 = 4 \end{aligned}$$

$\therefore \text{Centre} = (2, 0)$ Radius = 2

Question 4

- i $x^2 + (k+2)x + 4 < 0$
 $\Delta = (k+2)^2 - 16$
 $= k^2 + 4k - 12$
 $\Delta = 0$ for equal roots.
 $(k+6)(k-2) = 0$
 $\therefore k = -6 \text{ or } 2 \quad \text{ans}$ (2)
- ii $\Delta > 0$ for real & distinct roots
 $(k+6)(k-2) > 0$ 
 $\therefore k < -6, k > 2 \quad \text{ans}$ (2)




$$\begin{aligned} \text{Vol} &= \pi \int_0^4 y^2 dx \\ &= \pi \int_0^4 8 - \frac{x^2}{2} dx \\ &= \pi \left[8x - \frac{x^3}{6} \right]_0^4 \\ &= \pi \left[(32 - \frac{64}{6}) - 0 \right] \\ &= \frac{64\pi}{3} \text{ units}^3 \end{aligned}$$

Area $\hat{=} \frac{1}{2} \times \sum wxy$

$$\begin{aligned} &= \frac{4}{2} \times 6 \\ &= 12 \text{ m}^2 \end{aligned}$$

ii Volume $= 0.5 \times 12 \times 60 \times 60$
 $= 21600 \text{ m}^3$ (2)

x	y	w	wxy
0	0	1	0
4	1.3	2	2.6
8	1.7	2	3.4
12	0	1	0
$\sum wxy = 6$			



$$\begin{aligned} i & x^2 = (x-2)^2 \\ & x^2 = x^2 - 4x + 4 \\ & 4x = 4 \\ & x = 1, y = 1 \\ \therefore \text{Point } A &= (1, 1) \end{aligned}$$

$$\begin{aligned} ii \text{ Area} &= \int_0^1 x^2 dx + \int_1^2 (x-2)^2 dx \\ &= \left[\frac{x^3}{3} \right]_0^1 + \left[\frac{(x-2)^3}{3} \right]_1^2 \\ &= \left(\frac{1}{3} \right) - (0) + (0) - \left(-\frac{1}{3} \right) \\ &= \frac{2}{3} \text{ units}^2 \end{aligned}$$

iii) $V = \pi \int x_1^v dy - \pi \int x_2^v dy$

$$\sqrt{y} = x - 2$$

From diagram, $x-2 = -\sqrt{y}$

$$x = 2 - \sqrt{y}$$

$$x^2 = 4 - 4\sqrt{y} + y$$

$$\begin{aligned} \therefore V &= \pi \int_0^1 (4 - 4\sqrt{y} + y - y) dy \\ &= \pi \left[4y - \frac{8}{3}y^{3/2} \right]_0^1 \\ &= 4\pi/3 \text{ units}^3 \end{aligned}$$