

# SYDNEY GIRLS HIGH SCHOOL



2006 HSC Assessment Task 3

June 13, 2006

MATHEMATICS Extension 1

Year 12

Time allowed: 80 minutes

**Topics: Parametrics, Circle Geometry, Inverse Functions, Integration by Substitution**

**DIRECTIONS TO CANDIDATES:**

- Attempt all questions
- Questions are of equal value
- There are 4 questions with part marks shown in brackets
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work
- Board approved calculators may be used
- Write on one side of the paper only

a) Find the inverse of the following functions and state the domain and range

(i)  $y = \log_e(x-3)$  2

(ii)  $y = x^2 - 4x + 5 \quad x \geq 2$  2

b) Differentiate

(i)  $y = \sin^{-1} 3x$  2

(ii)  $y = \cos^{-1} \frac{x}{4}$  2

c) Find the primitive function of

(i)  $\int \frac{1}{4+x^2} dx$  1

d)  $f(x) = x \sin^{-1} x$

(i) what is the domain of  $f(x)$  1

(ii) show that this is an even function 2

(iii) verify that when  $x = 0$ ,  $f(x)$  is stationary 2

(iv) sketch a graph of  $y = f(x)$  1

(e) (i) Show that  $\frac{d(x^2 \tan^{-1} x)}{dx} = 2x \tan^{-1} x + 1 - \frac{1}{1+x^2}$  2

(ii) Hence or otherwise find  $\int_0^{\sqrt{3}} x \tan^{-1} x \, dx$  3

QUESTION 2

a) Find the following indefinite integrals using the substitution given

MARKS

(i)  $\int x\sqrt{x^2 + 4} dx$                        $u = x^2 + 4$                       2

(ii)  $\int \frac{dx}{x(\log x_e)^3}$                        $u = \log x_e$                       2

(iii)  $\int \frac{e^x dx}{\sqrt{49 - e^x}}$                        $u = e^x$                       2

b) Evaluate the following definite integrals using the substitution given

(i)  $\int_{-5}^0 \frac{t dt}{\sqrt{4-t}}$                        $t = 4 - u^2$                       4

(ii)  $\int_0^{\frac{\pi}{2}} \frac{\sin \theta}{3 - 2 \cos \theta} d\theta$                        $y = 3 - 2 \cos \theta$                       4


c) The region R is bounded by the curve  $y = \frac{x}{x+1}$   
the x-axis and the vertical line  $x = 3$ .

Use the substitution  $u = x + 1$  to find

(i) the exact area R                      3

(ii) the exact volume generated when R is rotated about the x-axis                      3

- a)  $T(2at, at^2)$  is a variable point on the parabola  $x^2 = 4ay$  3
- (i) show that the gradient of the tangent at T is  $t$ .
- (ii) show that the equation of the tangent at T is  $y = tx - at^2$
- b) Write down the equation of the chord of contact of tangents drawn from a point  $(x_1, y_1)$  to the parabola  $x^2 = 4ay$  4
- (i) find the equation of the chord of contact from the point  $(3, -2)$  to the parabola  $x^2 = 8y$
- (ii) at what point does the line intersect the directrix
- c) If PM is a normal to the parabola  $x^2 = 4ay$  at a variable point  $P(2ap, ap^2)$  and FN is drawn through the focus F parallel to the tangent at P to cut the normal at N 4
- (i) prove that the locus of  $N(x, y)$  is  $x^2 = a(y - a)$



d) The tangents at  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  on the parabola  $x^2 = 4ay$  intersect at  $T(a(p+q), apq)$


(i) find  $M$  the midpoint of  $PQ$

Hence show that


(ii)  $TM$  is parallel to the axis of symmetry

(iii) if  $TM$  meets the parabola on  $R$ , then  $R$  bisects  $TM$

(iv) the tangent at  $R$  is parallel to the chord  $PQ$



a)




O is the centre of the circle 5  
 Prove that

- (i)  $x + y = 90$
- (ii)  $z - y = 2x$

b)


3



BA and BC are tangents to the circles  
 $DE = 5 \times BD$ . Prove  $BA = \sqrt{6} \times BD$

c)

5




PQ is a common tangent and  $PA \parallel QB$ . Prove that

- (i)  $PC \parallel QD$
- (ii) PQBC is a cyclic quadrilateral

d)

4




P and Q are the centres of the circles  
PAX and QAY are straight lines.

Prove that P, Q, X and Y are concyclic

e)

3



A and B are fixed points. P moves on the plane  
so that AB subtends an angle of  $30^\circ$  at P.

- (i) describe the locus of P
- (ii) describe what construction you would carry out  
to draw the locus of P

THE END

Yr 12 Ext 2, June 06

a) if  $y = \log(x-3)$


$f^{-1}: x = \log(y-3)$

$y-3 = e^x$

$y = e^x + 3$

Domain = the set of reals

Range =  $\{y: y > 3\}$



(2)

ii)  $y = x^2 - 4x + 5, x \geq 2$

$f^{-1}: x = y^2 - 4y + 5, y \geq 2$

$x-5 = y^2 - 4y$

$x-5+4 = y^2 - 4y + 4$

$x-1 = (y-2)^2, y \geq 2$

$y-2 = \sqrt{x-1}, y \geq 2$

$y = 2 + \sqrt{x-1}, y \geq 2$

Domain =  $\{x: x \geq 1\}$

Range =  $\{y: y \geq 2\}$  (2)

i)  $y = \sin^{-1} 3x$

$|y| = \frac{1}{\sqrt{1-(3x)^2}} \times 3$

$= \frac{3}{\sqrt{1-9x^2}}$  (2)

ii)  $y = \cos^{-1} \frac{x}{4}$

$|y| = \frac{-1}{\sqrt{16-x^2}}$  (2)

i)  $\int \frac{1}{4+x^2} dx$

$= \frac{1}{2} \tan^{-1} \frac{x}{2} + C$  (1)

d) i)  $f(x) = x \cdot \sin^{-1} x$

Domain =  $\{x: -1 \leq x \leq 1\}$  (1)

ii)  $f(a) = a \cdot \sin^{-1} a$

$f(-a) = -a \times \sin^{-1}(-a)$

$= -a \times -\sin^{-1}(a)$   
 $= a \times \sin^{-1}(a)$  (2)

$\therefore f(-a) = f(a)$

$\therefore f(x)$  is an even function

iii)  $f(x) = x \cdot \sin^{-1} x$

$f'(x) = \sqrt{\frac{du}{dx}} + u \cdot \frac{dv}{dx}$

$= \sin^{-1} x \times 1 + x \times \frac{1}{\sqrt{1-x^2}}$

$= \sin^{-1} x + \frac{x}{\sqrt{1-x^2}}$


$f'(0) = \sin^{-1} 0 + \frac{0}{\sqrt{1-0}}$

$= 0 + 0$

$\therefore f'(0) = 0$

$\therefore$  When  $x=0, (x, f(x))$  is stationary. (2)

iv)



(1)

e) i)  $\frac{d}{dx} (x^2 \cdot \tan^{-1} x) = \sqrt{\frac{du}{dx}} + u \frac{dv}{dx}$

$= \tan^{-1} x \times 2x + x^2 \times \frac{1}{1+x^2}$

$\therefore \frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + \frac{x^2}{x^2+1}$

$= 2x \cdot \tan^{-1} x + \frac{x^2+1-1}{x^2+1}$

$\therefore \frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + \frac{x^2+1}{x^2+1} - \frac{1}{x^2+1}$

$= \frac{d}{dx} (x^2 \cdot \tan^{-1} x) = 2x \cdot \tan^{-1} x + 1 - \frac{1}{x^2+1}$

(2)



$$ii) \frac{d}{dx} (x^2 \tan^{-1} x) = 2x \cdot \tan^{-1} x + 1 - \frac{1}{x^2+1}$$

$$x^2 \cdot \tan^{-1} x = \int 2x \cdot \tan^{-1} x dx + \int 1 dx - \int \frac{dx}{x^2+1}$$

$$x^2 \cdot \tan^{-1} x = \int 2x \cdot \tan^{-1} x dx + (x - \tan^{-1} x + c)$$

$$\frac{2}{\sqrt{3}} \cdot \tan^{-1} x - x + \tan^{-1} x + c = \int 2x \cdot \tan^{-1} x dx$$

$$\int_0^{\sqrt{3}} 2x \cdot \tan^{-1} x dx = \left[ x^2 \tan^{-1} x - x + \tan^{-1} x + c \right]_0^{\sqrt{3}}$$

$$\int_0^{\sqrt{3}} x \cdot \tan^{-1} x dx = \left[ \frac{x^2}{2} \tan^{-1} x - \frac{x}{2} + \frac{1}{2} \tan^{-1} x + c \right]_0^{\sqrt{3}}$$

$$= \left( \frac{3}{2} \cdot \tan^{-1} \sqrt{3} - \frac{\sqrt{3}}{2} + \frac{1}{2} \tan^{-1} \sqrt{3} + c \right) - (0 - 0 + 0 + c)$$

$$= \frac{3}{2} \times \frac{\pi}{3} - \frac{\sqrt{3}}{2} + \frac{1}{2} \times \frac{\pi}{3}$$

$$= \frac{4\pi}{6} - \frac{\sqrt{3}}{2}$$

$$= \frac{2\pi}{3} - \frac{\sqrt{3}}{2}$$

(3)

$$2/i) \int x \sqrt{x^2+4} dx, u = x^2+4$$

$$= \int \sqrt{x^2+4} \cdot x dx, du = 2x dx$$

$$= \int \sqrt{u} \cdot \frac{1}{2} du, du = x dx$$

$$= \int \frac{1}{2} u^{1/2} du$$

$$= \frac{1}{2} \times \frac{2}{3} u^{3/2} + c$$

$$= \frac{1}{3} (x^2+4)^{3/2} + c \quad (2)$$

$$ii) \int \frac{dx}{x(\log_e x)^3}, u = \log_e x, du = \frac{1}{x} dx$$

$$= \int \frac{du}{u^3}$$

$$= \int u^{-3} du$$

$$= -\frac{1}{2} u^{-2} + c$$

$$= -\frac{1}{2u^2} + c$$

$$= -\frac{1}{2(\log_e x)^2} + c \quad (2)$$

$$ii) \int \frac{e^x dx}{\sqrt{49-e^x}}, u = e^x, du = e^x dx$$

$$= \int \frac{du}{\sqrt{49-u}}$$

$$= \int (49-u)^{-1/2} du$$

$$= 2(49-u)^{1/2} + c \quad (2)$$

$$= -2\sqrt{49-e^x} + c \quad (2)$$

$$b) i) \int \frac{t dt}{-5\sqrt{4-t}}, t = 4-u, dt = -2u du$$

$$= \int \frac{(4-u)(-2u) du}{-5\sqrt{4-(4-u^2)}}, u=2$$

$$= \int \frac{-2u(4-u^2) du}{-5\sqrt{4-4+u^2}}, u=-3$$

$$= \int \frac{2u(4-u^2) du}{\sqrt{u^2}}, u^2=9$$

$$= \int_2^3 \frac{2u(4-u^2) du}{u}$$

$$= \int_2^3 2(4-u^2) du$$

$$= \int_2^3 (8-2u^2) du$$

$$= \left[ 8u - \frac{2}{3} u^3 + c \right]_2^3$$

$$= \left( 8 \times 3 - \frac{2}{3} \times 3^3 + c \right) - \left( 8 \times 2 - \frac{2}{3} \times 2^3 + c \right)$$

$$= (24 - 18) - (16 - 16/3)$$

$$= 6 - 16 + 16/3$$

$$= -10 + 5\sqrt{3}$$

$$= -4\sqrt{3}$$

(4)

$$v) \int_0^{\pi/2} \frac{\sin \theta}{3-2\cos \theta} d\theta, \quad y = 3-2\cos \theta$$

$$= \int_1^3 \frac{dy}{y}$$

$$= \int_1^3 \frac{dy}{2y}$$

$$= \left[ \frac{1}{2} \ln y + c \right]_1^3$$

$$= \left( \frac{1}{2} \ln 3 + c \right) - \left( \frac{1}{2} \ln 1 + c \right) \quad (4)$$

$$= \frac{1}{2} \ln 3$$

$$dy = 2 \sin \theta d\theta$$

$$\theta = \pi/2, \quad y = 3 - 2\cos \theta$$

$$= 3 - 0 = 3$$

$$\theta = 0, \quad y = 3 - 2\cos \theta$$

$$= 3 - 2 = 1$$

$$= \pi \int_1^4 \frac{u^2 - 2u + 1}{u^2} du$$

$$= \pi \int_1^4 \left( 1 - \frac{2}{u} + \frac{1}{u} \right) du$$

$$= \pi \int_1^4 \left( 1 - \frac{2}{u} + u^{-2} \right) du$$

$$= \pi \left[ u - 2 \ln u - u^{-1} + c \right]_1^4$$

$$= \pi \left[ \left( 4 - 2 \ln 4 - \frac{1}{4} \right) - \left( 1 - 2 \ln 1 - 1 \right) \right]$$


$$= \pi \left( 3\frac{3}{4} - 2 \ln 4 + 2 \ln 1 \right)$$

$$= \pi \left( \frac{15}{4} - 2 \ln 4 \right)$$

sq units

$$(3)$$

$$v) i) \quad y = \frac{x}{x+1}$$



$$A = \int_0^3 \frac{x}{x+1} dx, \quad u = x+1$$

$$du = 1 dx$$

$$= \int_1^4 \frac{u-1}{u} du$$

$$= \int_1^4 \left( 1 - \frac{1}{u} \right) du$$

$$= \left[ u - \ln u + c \right]_1^4$$

$$= (4 - \ln 4 + c) - (1 - \ln 1 + c)$$


$$= 4 - \ln 4 - 1 + \ln 1$$

$$= (3 - \ln 4) \text{ sq units} \quad (3)$$

$$v) \quad V = \pi \int_0^3 y^2 dx$$

$$= \pi \int_0^3 \frac{x^2}{(x+1)^2} dx$$

$$= \pi \int_1^4 \frac{(u-1)^2}{u^2} du$$



$$y = \frac{x^2}{4a}$$

$$\frac{dy}{dx} = \frac{2x}{4a} = \frac{x}{2a}$$


$$\text{at } x = 2at, \quad \frac{dy}{dx} = \frac{2at}{2a} = t$$

$$\therefore T \text{ is } y - y_1 = m(x - x_1)$$

$$\therefore y - at^2 = t(x - 2at)$$

$$y - at^2 = tx - 2at^2$$

$$y = tx - at^2$$



$$x_1 x_2 = 2a(y_1 + y_2)$$

$$x_1 = 3, y_1 = -2, a = 2$$

$$\therefore 3x = 4(y - 2)$$

$$3x = 4y - 8$$


$$3x - 4y + 8 = 0$$

$$\text{at } y = -2, \quad 3x + 8 + 8 = 0$$

$$3x = -16$$

$$x = -5/3$$

$$\therefore P \text{ is } (-5/3, -2)$$



$$x^2 = 4ay$$

$$2x = 4a \cdot \frac{dy}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{x}{2a}$$

$$\text{at } x = 2ap, \quad \frac{dy}{dx} = p$$

$$\therefore m_{\perp} = -\frac{1}{p}$$

$$\therefore \text{Eqn is } y - ap^2 = -\frac{1}{p}(x - 2ap)$$

$$py - ap^3 = -x + 2ap$$

$$\therefore x + py = 2ap + ap^3$$

Line through focus is

$$y - a = p(x - 0)$$

$$y = px + a$$

Sub into normal.

$$\therefore x + p(px + a) = 2ap + ap^3$$

$$x + p^2x = 2ap + ap^3$$

$$x(1 + p^2) = ap(1 + p^2)$$


$$\therefore x = ap \text{ \& } y = ap^2 + a$$

$$\therefore \frac{x}{a} = p \Rightarrow y = a\left(\frac{x^2}{a^2}\right) + a$$

$$y = \frac{x^2}{a} + a$$

$$ay = x^2 + a^2$$

$$\therefore x^2 = a(y - a)$$



$$i) M \text{ is } \frac{2ap + 2aq}{2}, \frac{ap^2 + aq^2}{2}$$

$$= a(p+q), \frac{a}{2}(p^2 + q^2)$$

ii)  $M$  &  $T$  have the same  $x$  value,  
 $\therefore$  line is vertical  
 (i.e. parallel to  $y$ -axis)

iii) Midpoint  $MT$  is

$$a(p+q), \frac{1}{2} \left\{ \frac{a}{2}(p^2 + q^2) + apq \right\}$$

$$= a(p+q), \frac{1}{2} \left\{ \frac{ap^2 + aq^2 + 2apq}{2} \right\}$$

$$= a(p+q), \frac{a}{4}(p+q)^2$$

Sub into  $x^2 = 4ay$

$$\text{LHS} = a^2(p+q)^2$$


$$\text{RHS} = 4a \cdot \frac{a}{4}(p+q)^2 = a^2(p+q)^2 = \text{LHS}$$

$\therefore R$  lies on  $x^2 = 4ay$ .

$$iv) \frac{dy}{dx} = \frac{x}{2a} \quad \text{At } R, \frac{dy}{dx} = \frac{1}{2} a(p+q) = \frac{p+q}{2}$$

$$m_{PQ} = \frac{ap^2 - aq^2}{2ap - 2aq} = \frac{a(p+q)(p-q)}{2a(p-q)} = \frac{p+q}{2} = \text{grad of } \dots$$

4a)



- i)  $\angle AOB = 2y^\circ$  (angle at centre)  
 $\angle OBA = x^\circ$  (isos  $\Delta$ ,  $OA = OB$ )

In  $\Delta AOB$

$$x + x + 2y = 180 \text{ (angle sum of } \Delta)$$

$$\therefore 2x + 2y = 180$$


$$x + y = 90$$

- ii)  $y + z = 180$  (opp  $\angle$ 's of cyclic quad)  
 $\& 2x + 2y = 180$  (from above)

$$\therefore y + z = 2x + 2y$$

$$\therefore z - y = 2x$$

4b)



$AB = BC$  (tangents to a circle)

$BC^2 = BD \cdot BE$  (tangent/intercept thm)

Let  $BD = x$ ,  $\therefore DE = 5x$ ,  $BE = 6x$ .


$$\therefore BC^2 = x \cdot 6x = 6x^2$$

But  $AB = BC$

$$\therefore AB^2 = 6x^2$$

$$AB = \sqrt{6} \cdot x = \sqrt{6} \cdot BD$$

4c)



i) Let  $\angle APE = \alpha$

$\therefore \angle BQE = \alpha$  (corres  $\angle$ 's  $AP \parallel BQ$ )


$\& \angle ACP = \angle BQD = \alpha$   
 (angle in alt. segment)

$\therefore PC \parallel DQ$  (corres  $\angle$ 's  $\angle C \& \angle D$  are equal)

ii)  $\angle ACP = \alpha$   
 $\angle PCB = 180 - \alpha$  (str. angle)

$\therefore PCBQ$  is a cyclic quad  
 (opp  $\angle$ 's are supplementary)

4d)



Let  $\angle PYA = \alpha$


$\therefore \angle YAP = \alpha$  (isos  $\Delta$ ,  $PY, PA$  radii)

$\therefore \angle XAQ = \alpha$  (vert opp  $\angle$ 's)

$\therefore \angle AXQ = \alpha$  (isos  $\Delta$ ,  $AQ, XQ$  radii)  
 $= \angle PYA$

$\therefore PYXQ$  is a cyclic quad because  $P \& Q$  are subtending equal angles.

e)



i)  $P$  is the major arc of a circle.

ii) If the angle at  $P$  on the circumference is  $30^\circ$ , the angle at the centre is  $60^\circ$ .

$\therefore$  Construct  $60^\circ$  angles at  $A \& B$  & the centre of the circle is where the construction lines meet

$\therefore$  With compass on point  $O$  & radius  $OA$ , draw the major arc of a circle.