

# Sydney Girls High School



## MATHEMATICS- Extension 1 HSC ASSESSMENT TASK 2

**March 2005**

Topics: Exponential and Logarithmic Functions, Trigonometric Functions, Trigonometric Functions 2.

**Time Allowed: 75 minutes + 5 minutes reading time.**

Instructions:

- There are four (4) questions of equal value.
- Attempt all questions.
- Start each question on a new page.
- Show all necessary working.
- Marks may be deducted for careless or poor setting out.
- Board approved calculators may be used.
- A table of standard integrals is provided at the back of this paper.

**Total = 80 marks**

**QUESTION 1 (20 marks)**

**Marks**

(a) Given  $\log_2 3 \doteq 1.6$  and  $\log_2 5 \doteq 2.3$  evaluate:

(i)  $\log_2 15$

2

(ii)  $\log_2 30$

2

(b) Solve  $e^{2x-3} = 17$  for  $x$ , correct to 1 decimal place.

2

(c) Draw a neat sketch of  $y = \log_e(x-4)$ , showing all relevant features.

2

(d) Differentiate each of the following with respect to  $x$ :

(i)  $e^{3x}$

1

(ii)  $7^x$

1

(iii)  $6xe^{4x^2}$

2

(iv)  $\log_e(5x^2 + 2x)$

2

(e) Find the equation of the tangent to the curve  $y = 4e^{3x+1}$  at the  $y$ -intercept.  $x=0$

3

(f) Find:

(i)  $\int e^{6x+1} dx$

1

(ii)  $\int \frac{x^2}{x^3-10} dx$

2


$$\int \frac{x^2}{x^3-10} dx$$
  
$$3x^2$$
  
*See*

**QUESTION 2 (20 marks)**

Marks

(a) A fan in the shape of a sector of a circle has its frame made of cane and the shaded area is material. The frame consists of two circular arcs,  $AB$  and  $DC$ , and the sides  $AO$  and  $OB$ .

- (i) How much cane is used in the frame, to the nearest cm? 2  
 (ii) What area of material is used in the shaded part of the fan, to the nearest  $\text{cm}^2$ ? 2



(b) Differentiate the following with respect to  $x$ :

- (i)  $y = \cos(2x + 5)$  1  
 (ii)  $y = \log_e(\sin x)$  2  
 (ii)  $y = \frac{x}{\tan 4x}$  2

(c) Evaluate:

(i)  $\lim_{x \rightarrow 0} \frac{\sin \frac{x}{5}}{x}$  2

(ii)  $\int_0^{\frac{\pi}{6}} \tan^2 x \, dx$  2

*h*  $\int_0^{\pi/6} 1 - \sec^2 x \, dx$

(d) Find the equation of the normal to the curve  $y = \sin x$  at the point where  $x = \frac{\pi}{4}$ . 3

(e) Sketch the following curves for  $0 \leq x \leq 2\pi$ . State the amplitude and period of each curve.

- (i)  $y = 3 \sin 2x$  2  
 (ii)  $y = \cos\left(2x + \frac{\pi}{2}\right)$  2

**QUESTION 3 (20 marks)**

**Marks**

(a) If  $\sin a = \frac{\sqrt{3}}{2}$ ,  $\frac{\pi}{2} < a < \pi$ , and  $\cos b = \frac{1}{\sqrt{2}}$ ,  $0 < b < \frac{\pi}{2}$ , evaluate  $\cos(a-b)$ .

2

(b) Prove the following identities

(i)  $\cos\left(\frac{\pi}{6} + x\right) + \sin\left(\frac{\pi}{3} + x\right) = \sqrt{3} \cos x$

(ii)  $\sin 4x = 4 \sin x \cos^3 x - 4 \cos x \sin^3 x$

(c)

(i) Show that  $\sin^4 x = \left[\frac{1}{2}(1 - \cos 2x)\right]^2$

(ii) Hence or otherwise, show that  $\int \sin^4 x \, dx = \frac{3}{8}x - \frac{1}{4}\sin 2x + \frac{1}{32}\sin 4x + C$

Handwritten notes and calculations:

$$\cos 2x = 1 - 2\sin^2 x$$

$$2\sin^2 x = 1 - \cos 2x$$

$$\sin^2 x = \frac{1}{2}(1 - \cos 2x)$$

Annotations:  $\sin^2 x$ ,  $(\sin^2 x)^2$ ,  $(1 - \cos 2x)^2$

3

(d)

(i) Find  $\frac{d}{dx}(\sin^3 3x)$

2

(ii) Hence or otherwise find  $\int \sin^2 3x \cos 3x \, dx$

2

(e) Find the acute angle, to the nearest minute, between the following lines:

3

$3x + y - 5 = 0$  and  $x - 2y + 2 = 0$

**QUESTION 4 (20 marks)**

**Marks**

(a) Show that  $y = xe^{-x}$  has a maximum point at  $\left(1, \frac{1}{e}\right)$

3

(b) Evaluate  $\int_1^{\log_e 2} \frac{e^x + 1}{e^x} dx$

2

(c) (i) Show that  $\frac{x+2}{x+5} = 1 - \frac{3}{x+5}$


1

(ii) Hence or otherwise evaluate  $\int_2^3 \frac{x+2}{x+5} dx$


2

(d) Find the exact area of the region bounded by the curve  $y = \frac{2}{x+3}$ , the  $y$ -axis, and the line  $y = 2$ .

3



(e) The diagram below shows the curves  $y = \cos 2x$  and  $y = 2\sin^2 x$  for  $-\pi \leq x \leq \pi$ .



(i) Find the coordinates of A, a point of intersection of the two curves.

3

(ii) Find the exact area enclosed between the curves  $y = \cos 2x$  and  $y = 2\sin^2 x$  from the origin to the point A.

3

(iii) Find the volume of the solid formed if the area in part (ii) is rotated about the  $x$ -axis. [Hint: Question 3(c) (ii) may be of some use]

3

Year 12 Extension 1 Mathematics  
Assessment Task 2, 2005.

QUESTION 1 (20 marks)

(a)  $\log_2 3 \doteq 1.6$  and  $\log_2 5 \doteq 2.3$


(i)  $\log_2 15 = \log_2 (3 \times 5)$   
 $= \log_2 3 + \log_2 5$   
 (2)  $= 3.9$

(ii)  $\log_2 30 = \log_2 (2 \times 3 \times 5)$   
 $= \log_2 2 + \log_2 3 + \log_2 5$   
 (2)  $= 4.9$

or  $\log_2 30 = \log_2 (2 \times 15) = 1 + 3.9 = 4.9$

(b)  $e^{2x-3} = 17$   
 $\log_e e^{2x-3} = \log_e 17$   
 $2x-3 = \log_e 17$   
 $x = \frac{1}{2} (\log_e 17 + 3)$   
 (2)  $\doteq 2.9$

(c)  $y = \log_e (x-4)$



(d) (i)  $y = (e^{3x})$

$\frac{dy}{dx} = 3e^{3x}$  (1)

(ii)  $y = 7^x$   
 $= \log_e 7 \cdot 7^x$  (1)

(iii)  $y = 6xe^{4x^2}$

$\frac{dy}{dx} = uv' + vu'$

$u = 6x$
$u' = 6$
$v = e^{4x^2}$
$v' = 8xe$

(2)  $= 6x(8xe^{4x^2}) + e^{4x^2} \cdot 6$   
 $= 6e^{4x^2}(8x^2 + 1)$

(iv)  $y = \log_e (5x^2 + 2x)$

$\frac{dy}{dx} = \frac{10x+2}{5x^2+2x}$  (2)

(e)  $y = 4e^{3x+1}$

$\frac{dy}{dx} = 12e^{3x+1}$

(3)

at  $x=0$

$m = 12e$

Note: at  $x=0$   
 $y = 4e$

$y - y_1 = m(x - x_1)$

$y - 4e = 12e(x - 0)$

$0 = 12ex - y + 4e$

(f) (i)  $\int e^{6x+1} dx$

$= \frac{1}{6} e^{6x+1} + C$  (1)

(ii)  $\int \frac{x^2}{x^3-10} dx$

$f(x) = x^3 - 10$

$= \frac{1}{3} \int \frac{3x^2}{x^3-10} dx$

$f'(x) = 3x^2$

$= \frac{1}{3} \log_e (x^3 - 10) + C$  (2)

QUESTION 2 (20 marks)

(a) (i)

Amount of cane used  
 $= 2(8+5) + \left(\frac{120}{180} \times \pi \times 8\right) + \left(\frac{120}{180} \times \pi \times 13\right)$   
 $= 26 + \frac{16\pi}{3} + \frac{26\pi}{3}$   
 $= 26 + 14\pi$   
 $\hat{=} 70 \text{ cm}$  (2)

(ii)  $A = \frac{1}{2}r^2\theta$

Area of material =  $\left(\frac{1}{2} \times 13^2 \times \frac{2\pi}{3}\right) - \left(\frac{1}{2} \times 8^2 \times \frac{2\pi}{3}\right)$   
 $\hat{=} 110 \text{ cm}^2$  (2)

(b) (i)  $y = \cos(2x+5)$

$\frac{dy}{dx} = -2 \sin(2x+5)$  (1)

(ii)  $y = \log_e(\sin x)$

$\frac{dy}{dx} = \frac{\cos x}{\sin x}$   
 $= \cot x$  (2)

(iii)  $y = \frac{x}{\tan 4x}$

let  $u=x$   $v = \tan 4x$   
 $u' = 1$   $v' = 4 \sec^2 4x$

$\frac{dy}{dx} = \frac{vu' - uv'}{v^2}$  (2)  
 $= \frac{\tan 4x(1) - x(4 \sec^2 4x)}{\tan^2 4x}$   
 $= \frac{\tan 4x - 4x \sec^2 4x}{\tan^2 4x}$

(c) (i)  $\lim_{x \rightarrow 0} \frac{\sin \frac{x}{5}}{x}$

$= \lim_{x \rightarrow 0} \frac{\sin \frac{x}{5}}{\frac{x}{5}} \times \frac{1}{5}$   
 $= \frac{1}{5} \lim_{x \rightarrow 0} \frac{\sin \frac{x}{5}}{\frac{x}{5}}$  (2)  
 $= \frac{1}{5} (1)$   
 $= \frac{1}{5}$

(ii)  $\int_0^{\frac{\pi}{6}} \tan^2 x \, dx$   
 $= \int_0^{\frac{\pi}{6}} (\sec^2 x - 1) \, dx$   
 $= [\tan x - x]_0^{\frac{\pi}{6}}$   
 $= \frac{1}{\sqrt{3}} - \frac{\pi}{6}$   
 $= \frac{6 - \sqrt{3}\pi}{6\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$  (2)  
 $= \frac{2\sqrt{3} - \pi}{6}$

(d)  $y = \sin x$

$\frac{dy}{dx} = \cos x$



at  $x = \frac{\pi}{4}$   $\frac{dy}{dx} = \cos \frac{\pi}{4}$   
 $m_1 = \frac{1}{\sqrt{2}}$

$m_2 = -\sqrt{2}$

at  $x = \frac{\pi}{4}$   $y = \frac{1}{\sqrt{2}}$  (3)

$y - y_1 = m_2(x - x_1)$   
 $y - \frac{1}{\sqrt{2}} = -\sqrt{2}\left(x - \frac{\pi}{4}\right)$   
 $y = -\sqrt{2}x + \frac{\pi\sqrt{2}}{4} + \frac{\sqrt{2}}{2}$


$\sqrt{2}x + y + \frac{\pi\sqrt{2}}{4} - \frac{\sqrt{2}}{2} = 0$



QUESTION 3 (20 marks)

(a)  $\sin a = \frac{\sqrt{3}}{2}$  ,  $\frac{\pi}{2} < a < \pi$

$\cos b = \frac{1}{\sqrt{2}}$  ,  $0 < b < \frac{\pi}{2}$



$$\begin{aligned} \cos(a-b) &= \cos a \cos b + \sin a \sin b \\ &= \left(\frac{1}{2} \times \frac{1}{\sqrt{2}}\right) + \left(\frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}}\right) \\ &= \frac{-1 + \sqrt{3}}{2\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \\ &= \frac{\sqrt{6} - \sqrt{2}}{4} \end{aligned} \quad (2)$$

(b)(i)

$$\begin{aligned} \text{LHS} &= \cos\left(\frac{\pi}{6} + x\right) + \sin\left(\frac{\pi}{3} + x\right) \\ &= \cos\frac{\pi}{6} \cos x - \sin\frac{\pi}{6} \sin x + \sin\frac{\pi}{3} \cos x + \cos\frac{\pi}{3} \sin x \\ &= \frac{\sqrt{3}}{2} \cos x - \frac{\sin x}{2} + \frac{\sqrt{3}}{2} \cos x + \frac{\sin x}{2} \\ &= \sqrt{3} \cos x \\ &= \text{RHS} \end{aligned} \quad (3)$$

(ii) LHS =  $\sin 4x$

$$\begin{aligned} &= 2 \sin 2x \cos 2x \\ &= 2 (2 \sin x \cos x) (\cos^2 x - \sin^2 x) \\ &= 4 \sin x \cos^3 x - 4 \cos x \sin^3 x \\ &= \text{RHS} \end{aligned} \quad (3)$$

(c)(i)

$$\begin{aligned} \text{LHS} &= (\sin x)^4 \\ &= (\sin^2 x)^2 \\ &= \left[\frac{1}{2}(1 - \cos 2x)\right]^2 \end{aligned} \quad (2)$$

since  $\cos 2x = 1 - 2\sin^2 x$

$$\begin{aligned} 2\sin^2 x &= 1 - \cos 2x \\ \sin^2 x &= \frac{1}{2}(1 - \cos 2x) \end{aligned}$$

(ii)  $\int \sin^4 x dx = \frac{1}{4} \int (1 - \cos 2x)^2 dx$

$$\begin{aligned} &= \frac{1}{4} \int 1 - 2\cos 2x + \cos^2 2x dx \\ &= \frac{1}{4} \left[ x - \frac{2}{2} \sin 2x + \frac{x}{2} + \frac{\sin 4x}{8} \right] + C \\ &= \frac{x}{4} - \frac{1}{4} \sin 2x + \frac{x}{8} + \frac{1}{32} \sin 4x + C \\ &= \frac{3x}{8} - \frac{1}{4} \sin 2x + \frac{1}{32} \sin 4x + C \end{aligned} \quad (3)$$

Note:  $\int \cos^2 ax dx$

$$\begin{aligned} &= \frac{1}{2} \int (1 + \cos 2ax) dx \\ &= \frac{x}{2} + \frac{1}{4a} \sin 2ax + C \end{aligned}$$

(d)(i)  $\frac{d}{dx} (\sin^3 3x)$

$$\begin{aligned} &= 3(\sin 3x)^2 \cdot 3\cos 3x \\ &= 9 \sin^2 3x \cos 3x \end{aligned} \quad (2)$$

(ii)  $\int 9 \sin^2 3x \cos 3x dx = \sin^3 3x + C$

$$\int \sin^2 3x \cos 3x dx = \frac{1}{9} \sin^3 3x + C$$

(e)  $m_1 = -\frac{3}{1} = -3$  ,  $m_2 = \frac{1}{2}$

$$\theta = 81^\circ 52' \quad (3)$$

let  $\theta$  be the acute angle

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \left| \frac{-3 - \frac{1}{2}}{1 - \frac{3}{2}} \right| = 7$$

$\theta = 81^\circ 52'$



QUESTION 4 (20 marks)

(a)  $y = xe^{-x}$   $u = x \quad u' = 1$   
 $v = e^{-x} \quad v' = -e^{-x}$

$$\frac{dy}{dx} = uv' + v u' = x(-e^{-x}) + e^{-x}(1) = e^{-x} - xe^{-x}$$

For stat pts  $\frac{dy}{dx} = 0$

$0 = e^{-x} - xe^{-x}$   
 $0 = e^{-x}(1-x)$   
 $1 = x$  At  $x=1 \quad y = \frac{1}{e}$

$\frac{d^2y}{dx^2} = -e^{-x} - [e^{-x} - xe^{-x}]$   
 $= -2e^{-x} + xe^{-x}$

at  $x=1$

$\frac{d^2y}{dx^2} = -2e^{-1} + e^{-1} = -\frac{3}{e}$

$< 0 \therefore$  max pt at  $(1, \frac{1}{e})$

(b)  $\int_1^{\ln 2} \frac{e^x + 1}{e^x} dx$

$= \int_1^{\ln 2} (1 + e^{-x}) dx$


$= [x - e^{-x}]_1^{\ln 2}$   
 $= [\ln 2 - e^{-\ln 2}] - [1 - e^{-1}]$

$= \ln 2 - e^{-\ln 2} - 1 + \frac{1}{e}$   
 $= \ln 2 - \frac{1}{2} - 1 + \frac{1}{e}$   
 $= \ln 2 + \frac{1}{e} - \frac{3}{2}$

(c) (i)  $\frac{x+2}{x+5} = \frac{x+5-3}{x+5}$   
 $= 1 - \frac{3}{x+5}$  (1)

(ii)  $\int_2^3 \frac{x+2}{x+5} dx$   
 $= \int_2^3 (1 - \frac{3}{x+5}) dx$

$= [x - 3 \log_e(x+5)]_2^3$  (2)  
 $= (3 - 3 \log_e 8) - (2 - 3 \log_e 7)$   
 $= 1 - 3(\log_e 8 - \log_e 7)$   
 $= 1 - 3 \log_e(\frac{8}{7})$



Area = Area of square -  $\int_{-2}^0 \frac{2}{x+3} dx$

$= 4 - 2 \int_{-2}^0 \frac{dx}{x+3}$  (3)  
 $= 4 - 2 [\ln(x+3)]_{-2}^0$   
 $= 4 - 2 [\ln 3 - \ln 1]$   
 $= 4 - 2 \ln 3 = 1.8 u^2$

or,  $A = \left| \int_{\frac{2}{3}}^2 \frac{2-3}{y} dy \right|$   
 $= |2 \ln 3 - 4|$   
 $= 4 - 2 \ln 3 = 1.8 u^2$

(e) (i) Point of intersection = A(x,y)

$2 \sin^2 x = \cos 2x$   
 $2 \sin^2 x - \cos 2x = 0$   
 $2 \sin^2 x - (1 - 2 \sin^2 x) = 0$   
 $4 \sin^2 x - 1 = 0$   
 $\sin^2 x = \frac{1}{4}$   
 $\sin x = \pm \frac{1}{2}$

But A lies between  $0 \leq x \leq \frac{\pi}{2}$   
 $\sin x = \frac{1}{2}$

$x = \frac{\pi}{6}$  (3)

if  $x = \frac{\pi}{6} \quad y = \cos 2(\frac{\pi}{6}) = \frac{1}{2}$

$\therefore A(x,y) = (\frac{\pi}{6}, \frac{1}{2})$

(ii)  $A = \int_0^{\frac{\pi}{6}} \cos 2x dx - 2 \int_0^{\frac{\pi}{6}} \sin^2 x dx$

$= [\frac{1}{2} \sin 2x]_0^{\frac{\pi}{6}} - 2 \cdot \frac{1}{2} \int_0^{\frac{\pi}{6}} (1 - \cos 2x) dx$

$= [\frac{1}{2} \sin 2x]_0^{\frac{\pi}{6}} - [x - \frac{1}{2} \sin 2x]_0^{\frac{\pi}{6}}$

$= [\frac{1}{2} \sin \frac{\pi}{3} - 0] - [\frac{\pi}{6} - \frac{1}{2} \sin \frac{\pi}{3} - 0]$

$= \frac{\sqrt{3}}{4} - \frac{\pi}{6} + \frac{\sqrt{3}}{4}$

$= \frac{\sqrt{3}}{2} - \frac{\pi}{6}$  (3)

$= \frac{3\sqrt{3} - \pi}{6} u^2$

(iii)  $V = \pi \left[ \int_0^{\frac{\pi}{6}} \cos^2 2x dx - \int_0^{\frac{\pi}{6}} 4 \sin^4 x dx \right]$

Now for

$\int_0^{\frac{\pi}{6}} \cos^2 2x dx = \int_0^{\frac{\pi}{6}} [x + \frac{1}{4} \sin 4x]$

$= \frac{1}{4} [\frac{\pi}{6} + \frac{1}{4} \sin \frac{2\pi}{3} - 0]$

$= \frac{\pi}{12} + \frac{1}{4} (\frac{\sqrt{3}}{2})$

$= \frac{\pi}{12} + \frac{\sqrt{3}}{16}$  (3)

Now for

$4 \int_0^{\frac{\pi}{6}} \sin^4 x dx = 4 \left[ \frac{3x}{8} - \frac{1}{4} \sin 2x + \frac{1}{32} \sin 4x \right]_0^{\frac{\pi}{6}}$

$= 4 \left[ \frac{3\pi}{48} - \frac{1}{4} \sin \frac{\pi}{3} + \frac{1}{32} \sin \frac{2\pi}{3} - 0 \right]$

$= \frac{\pi}{4} - \frac{\sqrt{3}}{3} + \frac{1}{8} \sin \frac{2\pi}{3}$

$= \frac{\pi}{4} - \frac{\sqrt{3}}{2} + \frac{1}{8} (\frac{\sqrt{3}}{2})$

$= \frac{\pi}{4} - \frac{7\sqrt{3}}{16}$

Vol. =  $\pi \left( \frac{\pi}{12} + \frac{\sqrt{3}}{16} - \frac{\pi}{4} + \frac{7\sqrt{3}}{16} \right) u^3$

$= \pi \left( \frac{8\sqrt{3}}{16} - \frac{\pi}{6} \right) u^3$

$= \left( \frac{\sqrt{3}\pi}{2} - \frac{\pi^2}{6} \right) u^3$

(3)