

SYDNEY GIRLS HIGH SCHOOL



2006 HSC Assessment Task 3

June 7, 2006

MATHEMATICS

Year 12

Time allowed: 90 minutes

Topics: Locus & Parabola, Quadratics, Integration

DIRECTIONS TO CANDIDATES:

- Attempt all questions
- Questions are of equal value
- Part marks shown in brackets
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work
- Board approved calculators may be used
- Write on one side of the paper only

SGHS Mathematics 2U – Assessment Task 3: June 2006

Question 1.

a) For the parabola $x^2 = 4(y-1)$, write down

- i) The vertex
- ii) The focus
- iii) The equation of the directrix
- iv) The equation of the axis of symmetry

[4]

b) A parabola has the point (2, -2) as its vertex and (2,0) as the focus. Write down the equation of the parabola. [2]

c) Find the locus of the set of points P(x,y) given A(-1,-1) and B(5,3), such that

- i) PA and PB are the same length
- ii) PA and PB are perpendicular.

[4]

d) For the parabola $x^2 = 16y$,

- i) Find the gradient at the point P(-8,4).
- ii) Hence find the equations of the tangent and normal at P.
- iii) If the tangent cuts the Y axis at M and the normal cuts the Y axis at N, find the area of triangle PMN. [6]

e) Find the locus of the set of points where P(x,y) is equidistant from A(2,3) and the line $y = 1$ [4]

Question 2.

a) Solve for x where: $x^2 - 7x - 18 = 0$ [2]

b) For what values of k is the expression $x^2 - 2(k-3)x + (k-1)$ positive definite?

[4]

c) If α and β are the roots of the equation $2x^2 - 7x + 2 = 0$, find the values of:

i) $\alpha + \beta$

ii) $\alpha\beta$

iii) $\frac{1}{\alpha} + \frac{1}{\beta}$

iv) $\alpha^2 + \beta^2$

v) $\frac{4}{\alpha^2} + \frac{4}{\beta^2}$

[7]

d) Find all the real numbers x that satisfy the equation: $x^4 = 4x^2 + 32$

[3]

e) Find the values of k for the function $f(x) = 2x^2 - (3k-1)x + (2k-5) = 0$ to have

i) Sum of roots to be 4

ii) The roots to be reciprocals

[4]

Question 3.

- a) i) Use the Trapezoidal Rule with 3 values to estimate $\int_0^1 4^x dx$ [3]

ii) Is the estimate an under estimate or over estimate. Justify your answer. [1]

- b) Find the following indefinite integrals:

i) $\int (4 - 3x)^5 dx$

ii) $\int \frac{x^4 - 1}{x^2} dx$

iii) $\int \left(x + \frac{1}{x} \right)^2 dx$

[6]

- c) The curve $y = f(x)$ has the gradient function $f'(x) = 3x^2 - 2x + 1$. If the curve passes through the point Q(2,3), find the equation of the function.

[3]

- d) Given the function $y = 16^x$

- i) Copy and complete the following table

x	0	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{3}{4}$	1
y					

- ii) Use two applications of Simpson's Rule to find the approximate area enclosed by the curve, the X axis and the line $x = 1$ [3]

e) Given $y = \sqrt{x^2 + 16}$,

- i) Find $\frac{dy}{dx}$

- ii) Hence or otherwise evaluate $\int_0^3 \frac{2x dx}{\sqrt{x^2 + 16}}$ [4]

Question 4.

a) A parabola has the equation $x^2 = 8(4 - y)$

- { i) Sketch the parabola and clearly indicate the directrix, focus and points of intersection with the co-ordinate axes.
- ii) Another parabola Q, with equation $x^2 = 8y$ intersects the parabola P at A and B. Find the co-ordinates of A and B
- iii) Calculate the area of the region bounded by the two parabolas P and Q: [8]

b) Given $2mx^2 - (4m+1)x + 2 = 0$, show that the equation has rational roots if m is rational.

[3]

c) i) Find the points of intersection of the curve $y = 4 - \sqrt{2x}$ with the X and Y axes.

ii) The area enclosed by the curve $y = 4 - \sqrt{2x}$ and the X and Y axes is rotated about the Y axis. Find the volume of the solid of revolution that is formed.

[5]

d) Consider the points A(-3,-1) and B (6,2). If a point P(x,y) moves so that PA is twice the distance PB, show that the locus of P is a circle and find its centre and radius.


[4]

-----end of exam-----



YR 12 2u 2006 Q. ①

- (a) (i) $(0,1)$ ①
 (ii) $(0,2)$ ①
 (iii) $y \geq 0$ ①
 (iv) $x = 0$ ①



(b) $x^2 = 4ay$
 $(x-h)^2 = 4a(y-k)$
 $(x-2)^2 = 8(y+2)$ ②

$\Delta = 2$.

$x^2 + 2x + 1 + y^2 + 2y + 1 = x^2 - 10x + 25 + y^2 - 6y + 9$
 $12x + 8y - 32 = 0$
 $3x + 2y - 8 = 0$



(ii) $M_{PA} \times M_{PB} = -1$
 $\frac{y+1}{x+1} \cdot \frac{y-1}{x-5} = -1$
 $y^2 - 2y - 1 = -1$ ②
 $x^2 - 4x - 5$
 $y^2 - 2y - 1 = -x^2 + 4x + 5$
 $x^2 - 4x + y^2 - 2y - 8 = 0$
 or $(x-2)^2 + (y-1)^2 = 13$

(a) (i) $y = \frac{x}{16}$ ①
 $y = \frac{x}{8}$
 at $x = -8$ $y = -1$

(ii) $y-4 = -1(x+8)$ ③
 $y-4 = -x-8$

$(y-4)(x+8)$ $x+y+4=0$ (Tangent)

$y-4 = 1(x+8)$
 $y-4 = x+8$
 $x-y+12=0$ (Normal)



Q2

a) $x^2 - 7x - 18 = 0$
 $(x-9)(x+2) = 0$
 $x-9=0 \quad \text{or} \quad x+2=0$
 $x=9$ ① $x=-2$ ②
 $= 49 - 2$
 $= \frac{47}{4}$

b) $\Delta = b^2 - 4ac$
 $= [-2(k-3)]^2 - 4(1)(k-1)$
 $= 4(k^2 - 6k + 9) - 4k + 4$
 $= 4k^2 - 24k + 36 - 4k + 4$
 $= 4k^2 - 28k + 40$
 $= 4(k^2 - 7k + 10)$ ②

v) $\frac{4}{\alpha^2} + \frac{4}{\beta^2} = \frac{4\beta^2 + 4\alpha^2}{\alpha^2 \beta^2}$
 $= \frac{4(\alpha^2 + \beta^2)}{(\alpha \beta)^2}$

for def when $\Delta < 0$ and $a > 0$ ①

$4(k^2 - 7k + 10) < 0$
 $k^2 - 7k + 10 < 0$
 $(k-5)(k-2) < 0$

$2 < k < 5$ ①

i) $x^2 = 4x^2 + 32$
 $x^2 - 4x^2 - 32 = 0$
 Let $m = x^2$
 $m^2 - 4m - 32 = 0$ ③
 $(m-8)(m+4) = 0$
 $m-8=0 \quad \text{or} \quad m+4=0$
 $m=8 \quad \text{or} \quad m=-4$
 $\frac{m^2}{2} = \frac{8}{2} \quad \frac{m^2}{2} = \frac{-4}{2}$
 $\frac{x^2}{2} = \pm \sqrt{4}$
 $\Rightarrow 2x^2 - (3k-1)x + (2k-5) = 0$

ii) $\alpha \beta = \frac{c}{a}$
 $= \frac{-2}{2}$
 $= 1$ ①

Let roots be $\alpha, \alpha - 4 - \alpha$.
 $\alpha + (\alpha - 4) = 3k - 1$
 $4 = \frac{3k-1}{2}$

iii) $\frac{1+1}{\alpha \beta} = \frac{\alpha + \beta}{\alpha \beta}$
 $\frac{2}{\alpha \beta} = \frac{1}{\alpha \beta}$
 $\frac{2}{\alpha \beta} = 1$
 $\alpha \beta = 2$ ①


iv) $\alpha \beta = 1$ (reciprocal roots)
 $\alpha \beta = 2k - 5$
 $1 = \frac{2k-5}{2}$
 $2k-5 = 2$
 $2k = 7$
 $k = \frac{7}{2}$ ②

Question 3.

$$\text{ii} \quad \int_0^1 4^x dx$$

x	$y = 4^x$	w	w.y
0	1	1	1
0.5	2	2	4
1	4	1	4
$\sum w.y = 9$			

$$\begin{aligned}\int_0^1 4^x dx &= \frac{1}{2} \times \sum w.y \\ &= \frac{1}{2} \times 9 \\ &= \frac{9}{4} \\ &= 2\frac{1}{4} \text{ (or } 2.25)\end{aligned}$$



(3)

From Graph = Overestimate.

$$\text{or } \int_0^1 4^x dx = \left[\frac{4^x}{\ln 4} \right]_0^1 = \frac{(4-1)}{\ln 4} = 2.16$$

$$y = 16^x$$

x	$y = 16^x$	w	w.y
0	1	1	1
$\frac{1}{4}$	2	4	8
$\frac{1}{2}$	4	2	8
$\frac{3}{4}$	8	4	32
1	16	1	16
$\sum w.y = 65$			

(1)

$$\begin{aligned}\text{i} \quad &\int (4-3x)^5 dx \quad (2) \\ &= \frac{(4-3x)^6}{-3 \times 6} = \frac{(4-3x)^6}{-18} + C\end{aligned}$$

$$\text{ii} \quad \int \frac{x^4 - 1}{x^2} dx \quad (2)$$

$$\begin{aligned}&= \int x^2 - x^{-2} dx \\ &= \frac{x^3}{3} - \frac{x^{-1}}{-1} = \frac{x^3}{3} + x^{-1} + C\end{aligned}$$

$$\text{iii} \quad \int (x + \frac{1}{x})^2 dx$$

$$\begin{aligned}&= \int x^2 + 2 + x^{-2} dx \quad (2) \\ &= \frac{x^3}{3} + 2x - x^{-1} + C \\ &\quad (\text{or } \frac{x^3}{3} + 2x - \frac{1}{x} + C)\end{aligned}$$

$$\therefore f'(x) = 3x^2 - 2x + 1$$

$$f(x) = x^3 - x^2 + x + C$$

$$\text{Substitute } (2, 3)$$

$$3 = 8 - 4 + 2 + C$$

$$\therefore C = -3 \quad (3)$$

$$\therefore f(x) = x^3 - x^2 + x - 3$$

$$\begin{aligned}\int_0^1 16^x dx &= \frac{1}{3} \times \sum w.y \\ &= \frac{1}{3} \times 65 \\ &= \frac{65}{12} \\ &= 5\frac{5}{12} \text{ units}^2 \quad (2)\end{aligned}$$

$$\therefore y = \sqrt{x^2 + 16} = (x^2 + 16)^{\frac{1}{2}}$$

$$\begin{aligned}\therefore \frac{dy}{dx} &= \frac{1}{2} \cdot 2x(x^2 + 16)^{-\frac{1}{2}} \\ &= \frac{x}{\sqrt{x^2 + 16}} \quad (2)\end{aligned}$$

$$\begin{aligned}\text{ii} \quad &\int_0^3 \frac{2x}{\sqrt{x^2 + 16}} dx \\ &= 2 \left[\sqrt{x^2 + 16} \right]_0^3 \\ &= 2 \left[\sqrt{25} - \sqrt{16} \right]\end{aligned}$$

Yr 12 2U 2006
Assessment Task 3

Q4.

a) i) $x^2 = -8(y-4)$


$\sqrt{(0, 4)}$

$4a = 8$

$a = 2$

$F(0, 2)$

directrix $y=6$



(2)

$y=0 \rightarrow x^2 = 32$
 $x = \pm 4\sqrt{2}$

ii) $x^2 = 8y$ — ①
 $x^2 = 8(4-y)$ — ②
① = ②

$8y = 32 - 8y$
 $16y = 32$

$\boxed{y=2}$

sub in ①

$x^2 = 16$
 $x = \pm 4$

A(4, 2)

B(-4, 2)

$y = \frac{x^2}{8}$
 $8y = 32 - x^2$

iii) $\text{Area} = 2 \int_0^4 \frac{32-x^2}{8} - \frac{x^2}{8} dx$
 $= 2 \int_0^4 4 - \frac{x^2}{4} dx$
 $= 2 \left[4x - \frac{x^3}{12} \right]_0^4$
 $= 2 \left[16 - \frac{64}{12} \right]$
 $= 21 \frac{1}{3} u^2$

(3)

b) For rational
 Δ is a perfect square

ii) cont

$$\Delta = b^2 - 4ac$$

$$= (-[4m+1])^2 - 4 \times 2 \times 2$$

$$= 16m^2 + 8m + 1 - 16m$$

$$= 16m^2 - 8m + 1$$

$$= (4m-1)^2 \quad (3)$$

If m is rational
then the roots are rational

c) i) $x=0 \quad y=4$
when $y=0$ ①


$4 = \sqrt{2}x$

$2x = 16$

$x = 8$

(1)

ii)



$$\text{Vol} = \pi \int_0^4 \left(\frac{(4-y)^2}{2} \right)^2 dy$$

$$= \pi \int_0^4 \frac{(4-y)^4}{4} dy$$

d) $PA = 2PB$

$(x+3)^2 + (y+1)^2 = 4(x-6)^2 + (y-4)^2$

$$-x^2 + 6x + 9 + y^2 + 2y + 1 =$$

$$4(x^2 - 12x + 36) + (y^2 - 8y + 16)$$

$$3x^2 - 54x + 150 + 3y^2 - 18y +$$

$$x^2 - 12x + y^2 - 8y = -50 +$$

$$(x-9)^2 + (y-3)^2 = 40$$

$C(9, 3) \quad r = 2\sqrt{10}$

(4)