




2010 Half-Yearly Examination

FORM V MATHEMATICS EXTENSION 1

Friday 7th May 2010


QUESTION ONE (12 marks) Start a new leaflet.

- (a) Write down the exact value of $\cos 150^\circ$.
- (b) Solve $|x + 1| = 3$.
- (c) Consider the straight line $\sqrt{3}x - y + 2 = 0$.
 - (i) Write down the gradient of the line.
 - (ii) Find the angle of inclination of the line.
- (d) Write down the value of $8^{-\frac{2}{3}}$.
- (e) Factorise $x^3 - 8$.
- (f)



The diagram above shows the curve $y = x(x - 3)$. Use the curve to solve the inequation $x(x - 3) \geq 0$.

(g)



The diagram above shows the graph of the relation $y = -\sqrt{9 - x^2}$.

- (i) Write down the domain of the relation.
 - (ii) Explain why the relation is a function.
- (h) Calculate the limiting sum of the geometric series $3 + 1 + \frac{1}{3} + \dots$

General Instructions

- Writing time — 2 hours
- Write using black or blue pen.
- Board-approved calculators may be used.
- All necessary working should be shown in every question.
- Start each question on a new leaflet.

Structure of the paper

- Total marks — 96
- All eight questions may be attempted.
- All eight questions are of equal value.

Collection

- Write your name, class and master clearly on each leaflet.
- Hand in the eight questions in a single well-ordered pile.
- Hand in a leaflet for each question, even if it has not been attempted.
- If you use a second leaflet for a question, place it inside the first.
- Write your name on the question paper and place it inside your leaflet for Question One.

5A: DNW	5B: DS	5C: TCW
5D: MLS	5E: RCF	5F: PKH
5G: KWM	5H: REP	5I: SJE

Checklist

- Writing leaflets: 8 per boy.
- Candidature — 149 boys

Examiner
KWM


QUESTION TWO (12 marks) Start a new leaflet.

- (a) The first term of an arithmetic progression is -1 and the third term term is 9 .
- Find the common difference.
 - Find the fiftieth term.
 - Find the sum of the first fifty terms.
- (b) By rationalising the denominator, express $\frac{3}{2 + \sqrt{5}}$ in the form $a + b\sqrt{5}$.
- (c) The function $f(x)$ is defined by $f(x) = x^2 - x$.
- Find $f(x + h)$.
 - Use the definition $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ to find $f'(x)$.
 - Find the equation of the tangent to the curve $y = x^2 - x$ at the point $P(1, 0)$.
- (d) Solve for x
- $$2 \log_a 2 + \log_a 3 = \log_a x.$$

Exam continues overleaf ...

QUESTION THREE (12 marks) Start a new leaflet.


(a)




In the diagram above $\triangle ABC$ has vertices $A(3, 2)$, $B(-2, 1)$ and $C(1, -3)$. AD is an altitude of the triangle.

- Find the length of the side BC .
 - Find the gradient of BC .
 - Show that the equation of BC is $4x + 3y + 5 = 0$.
 - Use the perpendicular distance formula to find the length of the altitude AD .
 - Hence or otherwise, calculate the area of $\triangle ABC$.
- (b) Find the exact value of the pronumeral in each diagram below.

(i)



(ii)



- (c) The sum of the first n terms of a sequence is given by $S_n = n^2 - n$.
- Find S_8 and S_9 .
 - Hence find the ninth term of the sequence.

Exam continues next page ...

QUESTION FOUR (12 marks) Start a new leaflet.

- (a) For each of the following functions find $\frac{dy}{dx}$.
- (i) $y = x^3 + 2x - 3$
 - (ii) $y = (3x + 2)^4$
 - (iii) $y = \sqrt{x}$
 - (iv) $y = \frac{3x - 2}{x}$
- (b) Use the quotient rule to differentiate $\frac{2x}{x^2 + 1}$.
- (c) Use the product rule to differentiate $(2x + 1)^2(x^2 + 1)$ and express the derivative in factored form.
- (d) (i) Fully factorise $x^3 - x^2 - 2x$.
 (ii) Sketch the curve $y = x^3 - x^2 - 2x$, indicating all intercepts with the axes.
 (iii) Hence or otherwise, solve the inequality $x^3 - x^2 - 2x < 0$.

QUESTION FIVE (12 marks) Start a new leaflet.

- (a) Given $\tan \theta = -\frac{5}{12}$ and $\cos \theta > 0$, find the exact value of $\sin \theta$.
- (b) Sketch the graph of the function $y = 1 - |x - 2|$.
- (c) Find the equation of the straight line that passes through the point of intersection of the lines $2x - y + 1 = 0$ and $x + y - 2 = 0$ and also passes through the point $(2, -1)$. Do not find the point of intersection of the two lines and leave your answer in general form.
- (d) A function $h(x)$ is defined by $h(x) = \frac{2}{x^2 + 1}$.
- (i) Evaluate $h(0)$.
 - (ii) Show that $h(x)$ is an even function.
 - (iii) What value does $h(x)$ approach as $x \rightarrow \infty$?
 - (iv) Sketch the function $y = h(x)$ and state its range.

QUESTION SIX (12 marks) Start a new leaflet.


- (a) (i) Solve $\sin x + \sqrt{3} \cos x = 0$, for $0^\circ \leq x \leq 360^\circ$.
 (ii) Solve $\sec^2 x + \tan x = 3$, for $0^\circ \leq x \leq 360^\circ$.
 (Give your solutions correct to the nearest minute.)
- (b) (i) Write down the radius and centre of the circle $(x + 2)^2 + (y - 3)^2 = 25$.
 (ii) Show that the line $4x - 3y - 8 = 0$ is a tangent to the circle $(x + 2)^2 + (y - 3)^2 = 25$.
- (c) Prove the identity $\frac{\cos \alpha}{1 + \sin \alpha} + \frac{1 + \sin \alpha}{\cos \alpha} = 2 \sec \alpha$.

QUESTION SEVEN (12 marks) Start a new leaflet.

- (a) The third and sixth terms of a geometric sequence are $\frac{1}{27}$ and $\frac{1}{729}$ respectively.
- (i) Find the first term and the common ratio.
 - (ii) Find the sum of the first seven terms. (Express your answer as a fraction in simplest terms.)
 - (iii) How many terms of the sequence exceed $\frac{1}{1\,000\,000}$?
- (b) (i) Show that $x = 60^\circ$ is a solution to the equation $\sin x = \frac{1}{2} \tan x$.
 (ii) On the same set of axes sketch the graphs of the functions $y = \sin x$ and $y = \frac{1}{2} \tan x$, for $-180^\circ \leq x \leq 180^\circ$.
 (iii) Use your graphs to solve $\sin x \leq \frac{1}{2} \tan x$, for $-90^\circ < x < 90^\circ$.

QUESTION EIGHT (12 marks) Start a new leaflet.

(a)



In the diagram above the tangent and normal to the curve $y = \sqrt{x}$ at a variable point $P(a^2, a)$, where $a > 0$, intersect the x -axis at the points Q and R respectively.

- (i) Find the equation of the tangent at the point $P(a^2, a)$.
 - (ii) Find the co-ordinates of Q and R .
 - (iii) Hence show that the difference in the distances of the points Q and R from the origin is constant.
- (b) The two curves $y = x^2 + ax + b$ and $y = cx - x^2$ share a common tangent at the point $(1, 0)$. Find the values of the constants a , b and c .
- (c) Find the equation of the line which bisects the acute angle between the lines $l_1 : 3x - 6y - 10 = 0$ and $l_2 : 2x - y - 4 = 0$.

END OF EXAMINATION

QUESTION 1

(a) $\cos 150^\circ = -\cos 30^\circ$
 $= -\frac{\sqrt{3}}{2}$

(b) $|x+1| = 3$
 $x+1=3$ or $x+1=-3$
 $x=2$ or $x=-4$

(c) $\sqrt{3}x - y + 2 = 0$
 $y = \sqrt{3}x + 2$

(i) gradient = $\sqrt{3}$

(ii) $\tan \alpha = \sqrt{3}$
 $\alpha = 60^\circ$

angle of inclination is 60°

(d) $8^{\frac{2}{3}} = (2)^{-2}$
 $= \frac{1}{4}$

(e) $x^3 - 8 = (x-2)(x^2 + 2x + 4)$

(f) $x(x-3) \geq 0$

$x \leq 0, x \geq 3$

(g) (i) Domain: $-3 \leq x \leq 3$

(ii) the graph satisfies the vertical line test.

(h) $3 + 1 + \frac{1}{3} + \dots$

$S_{\infty} = \frac{a}{1-r}$

$S_{\infty} = \frac{3}{1-\frac{1}{3}}$

$S_p = \frac{3}{\frac{2}{3}}$

$S_p = \frac{9}{2}$

$S_p = 4\frac{1}{2}$ (12)

QUESTION 2

(a) $a = -1$ } (1)
 $a + 2d = 9$ } (2)

(i) $2d = 10$
 $d = 5$

(ii) $t_n = a + (n-1)d$
 $t_{50} = -1 + 49 \times 5$
 $t_{50} = 244$

(iii) $S_n = \frac{n}{2}(a+d)$
 $S_{50} = 25(-1 + 244)$
 $S_{50} = 6075$

(b) $\frac{3}{2+\sqrt{5}} \times \frac{2-\sqrt{5}}{2-\sqrt{5}}$
 $= \frac{6-3\sqrt{5}}{4-5}$
 $= -6 + 3\sqrt{5}$

(c) $f(x) = x^2 - x$

(i) $f(x+h) = (x+h)^2 - (x+h)$
 $= x^2 + 2xh + h^2 - x - h$

(ii) $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x - h - x^2 + x}{h}$
 $= \lim_{h \rightarrow 0} \frac{2xh + h^2 - h}{h}$
 $= \lim_{h \rightarrow 0} \frac{h(2x + h - 1)}{h}$

$= \lim_{h \rightarrow 0} 2x - 1 + h$
 $= 2x - 1$

(iii) $f'(x) = 2x - 1$
 $f'(1) = 1$
 gradient = 1
 $y - y_1 = m(x - x_1)$
 $y = 1(x - 1) + 0$
 tangent: $y = x - 1$

(d) $2 \log_a 2 + \log_a 3 = \log_a x$
 $\log_a 4 + \log_a 3 = \log_a x$
 $\log_a 12 = \log_a x$

(12) $x = 12$

QUESTION 3

(a) $A(3,2), B(-2,1), C(1,-3)$

(i) $\overline{BC} = \sqrt{(-2-1)^2 + (1-(-3))^2}$

$$\overline{BC} = \sqrt{9+16}$$
$$= 5 \text{ units } \checkmark$$

(ii) gradient $BC = \frac{1-(-3)}{-2-1}$
$$= -\frac{4}{3} \checkmark$$

(iii) $m = -\frac{4}{3} \quad B(-2,1)$

$y - y_1 = m(x - x_1)$

$y - 1 = -\frac{4}{3}(x + 2)$

$3y - 3 = -4x - 8$

$4x + 3y + 5 = 0 \checkmark$

(iv) $d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$

$d = \frac{|12 + 6 + 5|}{\sqrt{16 + 9}} \checkmark$

$d = \frac{23}{5}$

$= 4\frac{3}{5} \text{ units } \checkmark$

(v) Area $\Delta ABC = \frac{1}{2}bh$
$$= \frac{1}{2} \overline{BC} \times AD$$

$$= \frac{1}{2} \times 5 \times \frac{23}{5}$$

$$= \frac{23}{2}$$

$$= 11\frac{1}{2} \text{ sq units. } \checkmark$$

(b)

(i) $\frac{x}{\sin 30^\circ} = \frac{10}{\sin 135^\circ} \checkmark$

$$\frac{x}{\frac{1}{2}} = \frac{10}{\frac{1}{\sqrt{2}}}$$

$$2x = 10\sqrt{2} \checkmark$$

$$x = 5\sqrt{2} \text{ units.}$$

(ii)

$y^2 = 4 + 9 - 2 \times 3 \times \cos 60^\circ \checkmark$

$y^2 = 4 + 9 - 6$

$y = \sqrt{7} \text{ units. } \checkmark$

(c)

$S_n = n^2 - n$

(i) $S_8 = 64 - 8 \quad S_9 = 81 - 9$
$$= 56 \checkmark \quad S_9 = 72$$

(ii) $t_9 = S_9 - S_8$
$$= 72 - 56$$

$t_9 = 16 \checkmark \quad (12)$

QUESTION 4

(a)

(i) $y = x^3 + 2x - 3$

$\frac{dy}{dx} = 3x^2 + 2 \checkmark$

(ii) $y = (3x+2)^4$

$\frac{dy}{dx} = 4(3x+2)^3 \times 3$
$$= 12(3x+2)^3 \checkmark$$

(ii) $y = \sqrt{x}$

$y = x^{\frac{1}{2}}$

$\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} \checkmark$
$$= \frac{1}{2\sqrt{x}}$$

(iv) $y = \frac{3x-2}{x}$

$y = 3 - 2x^{-1}$

$\frac{dy}{dx} = \frac{2}{x^2} \checkmark$

(b) $f(x) = \frac{2x}{x^2+1}$

$f'(x) = \frac{(x^2+1)2 - 2x \times 2x}{(x^2+1)^2} \checkmark$

$$= \frac{2x^2 + 2 - 4x^2}{(x^2+1)^2}$$

$$= \frac{2 - 2x^2}{(x^2+1)^2} \checkmark$$

$$= \frac{2(1-x)(1+x)}{(x^2+1)^2}$$

(c) $f(x) = (2x+1)^2/(x^2+1)$

$f'(x) = \frac{2(2x+1)(x^2+1)2 + (2x+1)^2 \times 2x}{(x^2+1)^2} \checkmark$

$$= 2(2x+1) \{ 2(x^2+1) + x(2x+1) \}$$


$$= 2(2x+1)(4x^2+x+2) \checkmark$$

(d) (i) $x^3 - x^2 - 2x$

$$= x(x^2 - x - 2)$$

$$= x(x+1)(x-2) \checkmark$$

(ii)



(b) $f(x) = \frac{2x}{x^2+1}$

$f'(x) = \frac{(x^2+1)2 - 2x \times 2x}{(x^2+1)^2} \checkmark$

$$= \frac{2x^2 + 2 - 4x^2}{(x^2+1)^2}$$

$$= \frac{2 - 2x^2}{(x^2+1)^2} \checkmark$$

$$= \frac{2(1-x)(1+x)}{(x^2+1)^2}$$


(iii) $x(x+1)(x-2) < 0$

$$x < -1 \text{ OR } 0 < x < 2 \checkmark$$

(12)


QUESTION 5

(a) $\tan \theta = -\frac{5}{12}$ ✓



$\sin \theta = -\frac{5}{13}$ ✓

(b) $y = 1 - |x-2|$



(c) $2x - y + 1 + k(x + y - 2) = 0$ ✓

(2, -1): $4 + 1 + 1 + k(2 - 1 - 2) = 0$
 $6 - k = 0$
 $k = 6$ ✓

$2x - y + 1 + 6(x + y - 2) = 0$
 $8x + 5y - 11 = 0$ ✓

(d) $h(x) = \frac{2}{x^2 + 1}$

(i) $h(0) = 2$ ✓


(ii) $h(-x) = \frac{2}{(-x)^2 + 1}$
 $= \frac{2}{x^2 + 1}$

$\therefore h(x)$ is an even function. ✓

5.

(iii) as $x \rightarrow \infty$, $h(x) \rightarrow 0$ ✓

(iv)



Range: $0 < y \leq 2$ ✓

(12)

QUESTION 6

(a) (i) $\sin x + \sqrt{3} \cos x = 0$

$\frac{\sin x}{\cos x} + \sqrt{3} = 0$
 $\tan x = -\sqrt{3}$ ✓

$x = 120^\circ$ or $x = 300^\circ$ ✓

(ii) $\sec^2 x + \tan x = 3$

$(1 + \tan^2 x) + \tan x = 3$

$\tan^2 x + \tan x - 2 = 0$ ✓

$(\tan x - 1)(\tan x + 2) = 0$ ✓

$\tan x = 1$ $\tan x = -2$

$x = 45^\circ, 225^\circ$ $x = 116^\circ 34', 296^\circ 34'$ ✓

(b) (i) $(x+2)^2 + (y-3)^2 = 25$

radius = 5 units ✓

centre (-2, 3) ✓

(ii) The distance from the line $4x - 3y - 8 = 0$ to the centre of the circle (-2, 3)

$d = \frac{|-8 - 9 - 8|}{\sqrt{16+9}}$ ✓

$d = \frac{25}{5}$

$d = 5$ units. ✓

Since the perpendicular distance from the line to the centre equals the radius of the circle, the line intersects the circle at one point only. Hence the line is a tangent. ✓

6.

(c)

LHS = $\frac{\cos \alpha}{1 + \sin \alpha} + \frac{1 + \sin \alpha}{\cos \alpha}$

$= \frac{\cos^2 \alpha + (1 + \sin \alpha)^2}{\cos \alpha (1 + \sin \alpha)}$ ✓

$= \frac{\cos^2 \alpha + \sin^2 \alpha + 2 \sin \alpha + 1}{\cos \alpha (1 + \sin \alpha)}$

$= \frac{2 + 2 \sin \alpha}{\cos \alpha (1 + \sin \alpha)}$ ✓

$= \frac{2(1 + \sin \alpha)}{\cos \alpha (1 + \sin \alpha)}$ ✓

$= \frac{2}{\cos \alpha}$

$= 2 \sec \alpha$

$= \text{RHS}$ ✓

(12)

QUESTION 7

i) $ar^2 = \frac{1}{27}$ (1)
 $ar^5 = \frac{1}{729}$ (2)

(2) $\frac{ar^5}{ar^2} = \frac{1}{729} \div \frac{1}{27}$
 $r^3 = \frac{27}{729}$
 $r = \frac{3}{9}$
 $r = \frac{1}{3}$ ✓ $a = \frac{1}{3}$


(ii) $S_n = \frac{a(1-r^n)}{1-r}$
 $S_7 = \frac{\frac{1}{3}(1-(\frac{1}{3})^7)}{1-\frac{1}{3}}$
 $S_7 = \frac{1}{2}(1 - \frac{1}{2187})$
 $= \frac{1}{2} \times \frac{2186}{2187}$
 $= \frac{1092}{2187}$ ✓

(iii) $a_n > 10^{-6}$
 $ar^{n-1} > 10^{-6}$
 $\frac{1}{3} (\frac{1}{3})^{n-1} > 10^{-6}$
 $\frac{1}{3^n} > \frac{1}{1000000}$
 $3^n < 1000000$
 $n \log 3 < \log 1000000$
 $n \log 3 < 6$
 $n < \frac{6}{\log 3}$
 $n < 12.57...$
 $n = 12$ ✓

12 terms of the sequence exceed 10^{-6} .

(b) (i) $\sin x = \frac{1}{2} \tan x$
 $x = 60^\circ$: LHS = $\sin 60^\circ = \frac{\sqrt{3}}{2}$

RHS = $\frac{1}{2} \tan 60^\circ$
 $= \frac{1}{2} \times \sqrt{3}$
 $= \frac{\sqrt{3}}{2}$ $\therefore 60^\circ$ is a solut. ✓



(ii) $\sin x \leq \frac{1}{2} \tan x$ $-90^\circ < x < 90^\circ$
 $60^\circ \leq x < 90^\circ$ or $-60^\circ \leq x \leq 0^\circ$ ✓

(12)

QUESTION 8

(a) (i) $y = \sqrt{x}$
 $\frac{dy}{dx} = \frac{1}{2\sqrt{x}}$
 at $P(a^2, a)$ gradient = $\frac{1}{2a}$ ✓

$y - y_1 = m(x - x_1)$
 $y - a = \frac{1}{2a}(x - a^2)$
 $2ay - 2a^2 = x - a^2$ ✓
 $x - 2ay + a^2 = 0$ tangent.

(ii) Q: find $y=0$.
 $x = -a^2$
 $Q(-a^2, 0)$
 equation of the normal.
 $y - a = -2a(x - a^2)$ ✓
 find $y=0$
 $-a = -2a(x - a^2)$
 $\frac{1}{2} = x - a^2$
 $\frac{x}{2} = a^2 + \frac{1}{2}$
 $R(a^2 + \frac{1}{2}, 0)$ ✓

(iii) OR $\vec{OQ} - \vec{OR}$
 $|\frac{a^2 + 1/2}{2}| - |-a^2|$
 $= a^2 + \frac{1}{4} - a^2$
 $= \frac{1}{4}$ ✓

(b) (i) $y = x^2 + ax + b$
 (ii) $y = cx - x^2$
 (1,0) lies on both curves
 (i) $1 + a + b = 0$
 (ii) $c - 1 = 0$
 $\therefore c = 1$ ✓

$\frac{dy}{dx} = 2x + a$
 $\frac{dy}{dx} = c - 2x$
 at $x=1$ the gradients are equal.
 $a + 2 = c - 2$ ✓
 $a + 2 = 1 - 2$
 $a + 2 = -1$
 $a = -3$
 (i) $1 - 3 + b = 0$
 $b = 2$ ✓

(c) let $P(x, y)$ be any point on the angle bisector, d_1 the distance from P to l_1 and d_2 the distance from P to l_2 .
 $\frac{|3x - 6y - 10|}{\sqrt{45}} = \frac{|2x - y - 4|}{\sqrt{5}}$ ✓
 $|3x - 6y - 10| = 3|2x - y - 4|$
 $3x - 6y - 10 = 6x - 3y - 12$
 $3x + 3y - 2 = 0$ OR
 $3x - 6y - 10 = -6x + 3y + 12$
 $9x - 9y - 22 = 0$

acute angle bisector must have a gradient that lies between $l_1: m_1 = \frac{1}{2}$ and $l_2: m_2 = 2$.
 $\therefore 9x - 9y - 22 = 0$ is ✓
 the acute angle bisector of l_1 and l_2 .

(12)