

3 UNIT MATHEMATICS FORM V

Time allowed: 3 hours

Exam date: 18th October 2001

Instructions:

- All questions may be attempted.
- All questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.

Collection:

Each question will be collected separately.

Start each question on a new Answer Booklet.


If you use a second booklet for a question, place it inside the first. Do not staple.

Write your name, class and master's initials on each answer booklet:

5A: WMP	5B: JNC	5C: DNW	5D: REN
5E: KWM	5F: BDD	5G: FMW	5H: TCW

QUESTION ONE (Start a new Answer Booklet)

- (a) Solve $\tan \theta = 1$, for $0^\circ \leq \theta \leq 360^\circ$.
- (b) Simplify $\frac{x^3 - 8}{x^2 - 4}$.
- (c) (i) Show that there are 21 terms in the arithmetic series $-2 + 1 + 4 + \dots + 58$.
(ii) Hence or otherwise find the sum.
- (d) Solve the inequation $|x + 1| < 2$.
- (e)




Find the exact value of x in the diagram above.

- (f) (i) Find the gradient of the chord joining the points $A(1, 0)$ and $B(2, 4)$ on the curve $y = 2x^2 - 2x$.
(ii) Find the gradient of the tangent to the curve $y = 2x^2 - 2x$ at the point $B(2, 4)$.
- (g) Consider the function $f(x) = x^2 + 3x$.
(i) Show that $f(x + h) - f(x) = 2xh + h^2 + 3h$.
(ii) Use the definition $f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$ to find $f'(x)$.

QUESTION TWO (Start a new Answer Booklet)

- (a) Find the perpendicular distance from the point $(2, 1)$ to the line $12x - 5y + 5 = 0$.
- (b) Find the values of a and b , given that $(3\sqrt{2} - 1)^2 = a - b\sqrt{2}$.
- (c) The limiting sum of the geometric series $a + \frac{a}{2} + \frac{a}{4} + \dots$ is 6. Find the value of a .
- (d) At any point on a curve, $\frac{dy}{dx} = 2x + 5$. If the curve passes through the point $P(1, -2)$, find the equation of the curve.

(e)



The diagram above shows a sketch of $y = f(x)$ with a maximum turning point at (c, b) , a minimum turning point at $(-c, -b)$, and a point of inflexion at the origin.

- (i) $y = f(x)$ is known to be an odd function. How does the graph illustrate this property?
- (ii) For what values of x is $f'(x) < 0$?
- (iii) Write down the co-ordinates of the point where the gradient of the curve is a maximum.
- (iv) Find the value of $\int_{-a}^a f(x) dx$.
- (f) Solve $\frac{1}{x+1} > 2$.

QUESTION THREE (Start a new Answer Booklet)

- (a) Write down the equation of the locus of the point $P(x, y)$ which moves so that its distance from the point $C(2, -1)$ is always 2 units.
- (b) Differentiate the following with respect to x :
- $y = 3x^4 + 2x^2 - 3$,
 - $y = \frac{3}{x}$,
 - $y = 6\sqrt{x}$,
 - $y = (4x^2 - 3x)^{10}$.
- (c) Consider the function $f(x) = x(x - 3)^2$.
- Find the intercepts with the x and y axes of the curve $y = f(x)$.
 - Show that $f'(x) = 3(x - 1)(x - 3)$, and find $f''(x)$.
 - Find the co-ordinates of any stationary points and determine their nature.
 - Find the co-ordinates of any points of inflexion.
 - Sketch the graph, indicating all the important features.

QUESTION FOUR (Start a new Answer Booklet)

- (a) Find each of the following indefinite integrals:

$$(i) \int (5x^2 + 2x - 3) dx,$$


$$(ii) \int \frac{x^3 - x}{x^3} dx.$$

- (b) (i) Copy and complete the table below for the function $y = 2^{-x}$.

x	-2	-1	0	1	2
y					

- (ii) Use Simpson's rule with the five function values from the table to find an approximation for $\int_{-2}^2 2^{-x} dx$.

(c)



The diagram above shows the region bounded by the parabola $y = 2x^2 - 2$, the x -axis and the line $x = 2$. Calculate the area of this region.

(d) Consider the parabola with the equation $x^2 - 6x + 4y + 5 = 0$.

- (i) Express the parabola in the form $(x - h)^2 = -4a(y - k)$, and hence find the co-ordinates of the vertex of the parabola.
- (ii) Find the co-ordinates of the focus of the parabola.

(e) (i) Differentiate $y = \sqrt{3x^2 - 2x}$.


(ii) Hence find $\int \frac{3x - 1}{\sqrt{3x^2 - 2x}} dx$.

QUESTION FIVE (Start a new Answer Booklet)

- (a) (i) Complete the square to find the minimum value of the quadratic function
 $y = x^2 + 4x - 6$.
- (ii) Find the values of k for which the quadratic equation $x^2 + 4x - (6 + k) = 0$ has real roots.
- (iii) Let the solutions of the quadratic equation $x^2 + 4x - 6 = 0$ be α and β . Find :
- (α) $\alpha + \beta$,
 - (β) $\alpha\beta$,
 - (γ) $\frac{1}{\alpha^2} + \frac{1}{\beta^2}$.

(b) Solve $2 \sin^2 \theta - 5 \sin \theta + 2 = 0$, for $-180^\circ \leq \theta \leq 180^\circ$.


(c)



The diagram above shows the curve $y = f(x)$ with points of inflection at $(2, 0)$ and $(4, 0)$. Sketch the graph of $y = f'(x)$.

QUESTION SIX (Start a new Answer Booklet)


(a)



The diagram above shows the parabola $x^2 = 4y$ and the chord of contact AB from the external point $P(1, -2)$.

- Write down the equation of the chord of contact.
 - Show that the endpoints of the chord are $A(-2, 1)$ and $B(4, 4)$.
 - Find the area of the shaded region.
- (b) (i) Sketch $y = |x - 4|$.
(ii) For what values of c does the equation $|x - 4| = \frac{1}{2}x + c$ have two distinct solutions?

(c)



The diagram above shows the parabola $x^2 = 8y$ with focus $S(0, 2)$. The tangent at the variable point $T(4t, 2t^2)$ meets the x -axis at A .

- Show that the equation of the tangent at T is $y = tx - 2t^2$.
- Find the co-ordinates of the mid-point M of TA .
- Find the Cartesian equation of the locus of M .


QUESTION SEVEN (Start a new Answer Booklet)

- (a) Use mathematical induction to prove that

$$\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{(n+1)(n+2)} = \frac{n}{2(n+2)}$$


for all positive integers n .

(b)



The diagram above shows the region bounded by the semi-circular curve $y = \sqrt{r^2 - x^2}$ and the line $y = r - h$, where $0 \leq h \leq r$. Show that the volume of the solid formed when this region is rotated about the y -axis is given by $V = \frac{\pi}{3}h^2(3r - h)$.

(c)



Water is pumped into a hemispherical dam of radius 4 metres at a rate of 3π cubic metres per hour. How fast is the water rising when the water depth h is 3 metres? (Hint: You will need to use the formula derived in part (b).)

- (d) Draw a possible sketch of a curve $y = f(x)$ for which:

- the domain is the set of all real numbers, and
- the curve is differentiable for all values of x except $x = 2$, and
- the x -axis is a horizontal asymptote in both directions, and
- the function has positive values for $x > 0$, and negative values for $x < 0$.

QUESTION EIGHT (Start a new Answer Booklet)


(a) Find where the curve $y = \frac{1-x^2}{1+x^2}$ is concave down.

(b) (i) Prove that $\frac{1}{(1+nx)(1+(n+1)x)} = \frac{1}{x} \left(\frac{1}{1+nx} - \frac{1}{1+(n+1)x} \right)$.

(ii) Hence find the sum of the first n terms of the series

$$\frac{1}{(1+x)(1+2x)} + \frac{1}{(1+2x)(1+3x)} + \frac{1}{(1+3x)(1+4x)} + \dots$$

(c)



The diagram above shows a container of fixed volume V with a square base of length x units. The eight edges AB , BC , CD , DA , AE , BF , CG and DH are constructed of metal piping for extra strength.

(i) Show that the total length L of these eight edges is $L = 4x + \frac{4V}{x^2}$.

(ii) Find the minimum value of L .

(d) (i) Factorise $a^5 - b^5$.


(ii) Show that $77^5 - 22^5 + 32^4 - 12^4$ is divisible by 5. You may not use your calculator.

KWM

QUESTION 1

(a) $\tan \theta = 1$

$$\theta = 45^\circ \text{ or } \theta = 225^\circ \checkmark$$



$$(b) \frac{n^3 - 8}{n^2 - 4} = \frac{(n-2)(n^2 + 2n + 4)}{(n-2)(n+2)}$$

$$= \frac{n^2 + 2n + 4}{n+2} \checkmark$$

(c) (i) Put $a + (n-1)d = 58 \checkmark$

$$-2 + (n-1)3 = 58$$

$$3n - 5 = 58$$

$$3n = 63$$

$$n = 21 \checkmark$$

58 is the 21st term.

(ii) $S_n = \frac{n}{2} (a + l)$

$$S_{21} = \frac{21}{2} \times 56$$

$$S_{21} = 588 \checkmark$$

(d) $|x+1| < 2$

$$x+1 < 2 \text{ and } x+1 > -2$$

$$x < 1 \text{ and } x > -3$$

$$-3 < x < 1 \checkmark$$

(e)

$$\frac{x}{\sin 30^\circ} = \frac{8}{\sin 45^\circ} \checkmark$$

$$x = \frac{8 \sin 30^\circ}{\sin 45^\circ}$$

$$x = 4 \times \frac{1}{2} \times \sqrt{2}$$

$$x = 4\sqrt{2} \text{ cm } \checkmark$$

(f) (i) gradient = $\frac{y_2 - y_1}{x_2 - x_1}$

$$= \frac{4-0}{2-1}$$

$$= 4 \checkmark$$

(ii) $y = 2x^2 - 2x$

$$\frac{dy}{dx} = 4x - 2 \checkmark \text{ A } + B(2,4),$$

$$\text{gradient} = 6. \checkmark$$

(g) (i) $f(n) = n^2 + 3n$

$$f(n+h) = (n+h)^2 + 3(n+h)$$

$$= n^2 + 2nh + h^2 + 3n + 3h$$

$$f(n+h) - f(n) = 2nh + h^2 + 3h \checkmark$$

(ii) $f'(n) = \lim_{h \rightarrow 0} \frac{f(n+h) - f(n)}{h}$

$$= \lim_{h \rightarrow 0} \frac{n^2 + 2nh + h^2 + 3n + 3h - n^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2nh + h^2 + 3h}{h} \checkmark$$

$$= \lim_{h \rightarrow 0} \frac{-h(2n + 3 + h)}{h}$$

$$= \lim_{h \rightarrow 0} (2n + 3 + h) \checkmark$$

$$= 2n + 3.$$

(15)

QUESTION 2

(a) $12x - 5y + 5 = 0 \quad (2,1)$

$d = \left| \frac{ax_0 + by_0 + c}{\sqrt{a^2 + b^2}} \right| \quad \checkmark$

$= \frac{24 - 5 + 5}{\sqrt{144 + 25}} \quad \checkmark$

$= \frac{24}{13} \text{ units} \quad \checkmark$

(b) $(3\sqrt{2} - 1)^2 = a - b\sqrt{2}$

$18 - 6\sqrt{2} + 1 = a - b\sqrt{2}$

$19 - 6\sqrt{2} = a - b\sqrt{2} \quad \checkmark$

$a = 19 \text{ and } b = 6 \quad \checkmark$

c) $a + \frac{a}{2} + \frac{a}{4} + \dots$, is a GPwith $r = \frac{1}{2}$.

Put $S_\infty = 6$

$\frac{a}{1-r} = 6 \quad \checkmark$

$a = 6 - 6r$

$a = 6 - 3$

$a = 3 \quad \checkmark$

d) $\frac{dy}{dx} = 2x + 5$

$y = x^2 + 5x + c \quad \checkmark$

since the curve passes through

$P(1, -2)$,

$-2 = 1 + 5 + c$

$c = -8$

So the equation of the curve
is $y = x^2 + 5x - 8 \quad \checkmark$ (e) i) The graph has point symmetry about the origin. (or rotational symmetry of order 2) \checkmark ii) When $f'(x) < 0$, the curve is decreasing. Hence $x < -c$ or $x > c$ iii) The gradient of the curve is a maximum at the point of inflection $(0, 0)$. \checkmark iv) $\int_{-a}^a f(x) dx = 0$, since the function is odd. \checkmark

(f) $\frac{1}{x+1} > 2 \quad \text{multiplying by}$


$(x+1)^2$,

$x+1 > 2(x+1)^2$

$x+1 > 2(x^2 + 2x + 1)$

$2x^2 + 3x + 1 < 0 \quad \checkmark$

$(2x+1)(x+1) < 0$



$-1 < x < -\frac{1}{2} \quad \checkmark$

15

QUESTION 3 The locus is a

(a) Circle, centre $(2, -1)$, radius = 2
 $(x-2)^2 + (y+1)^2 = 4 \checkmark$

(b) (i) $y = 3x^4 + 2x^2 - 3$
 $\frac{dy}{dx} = 12x^3 + 4x \checkmark$

(ii) $y = 3x^{-\frac{1}{2}}$
 $\frac{dy}{dx} = -3x^{-\frac{3}{2}}$
 $= -\frac{3}{x^{\frac{3}{2}}} \checkmark$

(iii) $y = 6x^{\frac{1}{2}}$
 $\frac{dy}{dx} = 3x^{-\frac{1}{2}}$
 $\frac{d^2y}{dx^2} = \frac{3}{\sqrt{x}} \checkmark$

(iv) $y = (4x^2 - 3x)^{10}$
 $\frac{dy}{dx} = 10(4x^2 - 3x)^9(8x - 3) \checkmark$

(i) $f(x) = x(x-3)^2$

(i). Y intercept: $(0, 0) \checkmark$

X intercepts: $(0, 0)$ and $(3, 0) \checkmark$

(ii) $f(x) = x(x-3)^2$

$$\begin{aligned} f'(x) &= (x-3)^2 + 2x(x-3) \\ &= (x-3)(x-3 + 2x) \checkmark \\ &= (x-3)(3x-3) \\ &= 3(x-3)(x-1) \end{aligned}$$

$$f'(x) = 3x^2 - 12x + 9$$

$$f''(x) = 6x - 12 \quad \checkmark$$

(iii) For stationary points,

$$\text{put } f'(x) = 0$$

$$3(x-3)(x-1) = 0$$

$x = 1$ or $x = 3$. Hence there are stationary

points at $(1, 4)$ and $(3, 0) \checkmark$

At $(1, 4)$, $f''(1) = -6$

$f''(1) < 0$, so there is a local maximum at $(1, 4) \checkmark$

At $(3, 0)$, $f''(3) = 6$

$f''(3) > 0$, so there is a local minimum at $(3, 0) \checkmark$

(iv)


$$\text{Put } f''(x) = 0$$

$$6x - 12 = 0$$

$x = 2$, so there is a possible point of inflection at $(2, 2) \checkmark$

x	1.9	2	2.1	A change
$f''(x)$	-	0	+	✓

in concavity occurs at $(2, 2)$, so there is a point of inflection at $(2, 2) \checkmark$



15

QUESTION 4

(a) (i) $\int (5x^2 + 2x - 3) dx$

$$= \frac{5}{3}x^3 + x^2 - 3x + C \quad \checkmark$$

(ii) $\int (1-x^{-2}) dx = x + \frac{1}{x} + C \quad \checkmark$

(b) (i) $y = 2^{-x}$

-2	-1	0	1	2
4	2	1	$\frac{1}{2}$	$\frac{1}{4}$

$$= \left[\left(\frac{2}{3} - 2 \right) - \left(-\frac{2}{3} + 2 \right) \right] + \left[\left(\frac{16}{3} - 4 \right) - \left(\frac{2}{3} - 2 \right) \right]$$

$$= \frac{8}{3} + \frac{8}{3}$$

$$= 5\frac{1}{3} \text{ square units.} \quad \checkmark$$

(d) (i) $x^2 - 6x + 4y + 5 = 0$

$$(x-3)^2 - 9 + 4y + 5 = 0$$

$$(x-3)^2 = -4y + 4$$

Hence the vertex is $(3, 1)$ and focal length = 1. \checkmark

(ii) Simpson's Rule.

(ii) the focus is $(3, 0)$. \checkmark

$$\int_{-2}^2 2^{-x} dx = \int_{-2}^0 2^{-x} dx + \int_0^2 2^{-x} dx$$

$$\therefore \frac{1}{3} \left(4 + 4 \times 2 + 1 \right) + \frac{1}{3} \left(1 + 4 \times \frac{1}{2} + \frac{1}{4} \right) \quad \checkmark$$

$$\therefore \frac{13}{3} + \frac{13}{12}$$

$$\therefore \frac{65}{12}$$

$$\therefore 5\frac{5}{12} \quad \checkmark$$

(e)

(i) $y = (3x^2 - 2x)^{\frac{1}{2}}$

$$\frac{dy}{dx} = \frac{1}{2} (3x^2 - 2x)^{-\frac{1}{2}} (6x - 2) \quad \checkmark$$

$$= \frac{3x-1}{\sqrt{3x^2-2x}} \quad \checkmark$$

(ii) From part (i),

$$\int \frac{3x-1}{\sqrt{3x^2-2x}} dx = \sqrt{3x^2-2x} + C \quad \checkmark$$

(c)

$$\text{Area} = - \int_{-1}^1 (2x^2 - 2) dx + \int_1^2 (2x^2 - 2) dx \quad \checkmark$$

$$= - \left[\frac{2x^3}{3} - 2x \right]_{-1}^1 + \left[\frac{2x^3}{3} - 2x \right]_1^2 \quad \checkmark$$

QUESTION 5

(a) (i) $y = x^2 + 4x - 6$
 $y = (x+2)^2 - 4 - 6$
 $y = (x+2)^2 - 10 \quad \checkmark$
minimum value = $-10. \quad \checkmark$

(ii) $x^2 + 4x - (6+k) = 0$

for real roots $\Delta \geq 0.$

$$\begin{aligned}\Delta &= b^2 - 4ac \\ &= 16 + 4(6+k) \\ &= 40 + 4k. \quad \checkmark \\ 40 + 4k &\geq 0 \\ 4k &\geq -40\end{aligned}$$

for real roots $k \geq -10. \quad \checkmark$

(iii) $x^2 + 4x - 6 = 0$


(a) $\alpha + \beta = -\frac{b}{a}$
 $= -4 \quad \checkmark$

(b) $\alpha\beta = \frac{c}{a}$
 $= -6 \quad \checkmark$

$$\begin{aligned}(c) \quad \frac{1}{\alpha^2} + \frac{1}{\beta^2} &= \frac{\beta^2 + \alpha^2}{\alpha^2 \beta^2} \\ &= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{(\alpha\beta)^2} \quad \checkmark \\ &= \frac{16 + 12}{36} \\ &= \frac{7}{9} \quad \checkmark\end{aligned}$$

(d) $2\sin^2\theta - 5\sin\theta + 2 = 0$
 $(2\sin\theta - 1)(\sin\theta - 2) = 0 \quad \checkmark$
 $2\sin\theta = 1 \quad \text{or} \quad \sin\theta = 2$
 $\sin\theta = \frac{1}{2} \quad \checkmark \quad \text{no solutions.}$
 $\theta = 30^\circ \quad \text{or} \quad \theta = 150^\circ \quad \checkmark$

(e)



(15)

QUESTION 6

(a) (i) The chord of contact from $P(1, -2)$ is

$$x_0 x = 2a(y + y_0). \text{ When } a=1,$$

$$x = 2(y - 2)$$

$$x = 2y - 4$$

$$x - 2y + 4 = 0$$

$$(ii) \text{ Solve } x^2 = 4ay \dots \textcircled{1}$$

$$x = 2y - 4 \dots \textcircled{2}$$

Simultaneously,

$$\textcircled{2} \quad (2y - 4)^2 = 4ay$$

$$4y^2 - 16y + 16 = 4ay$$

$$4y^2 - 20y + 16 = 0$$



$$y^2 - 5y + 4 = 0 \quad \checkmark$$

$$(y-1)(y-4) = 0$$

When $y=1$, $x=-2$. When $y=4$, $x=4$. So the end points are $A(-2, 1)$ and $B(4, 4)$.

$$\begin{aligned} (\text{iii}) \quad \text{Area} &= \int_{-2}^4 \left(\frac{1}{2}x + 2 - \frac{1}{4}x^2 \right) dx \quad \checkmark \\ &= \left[\frac{1}{4}x^2 + 2x - \frac{1}{12}x^3 \right]_{-2}^4 \quad \checkmark \\ &= \left(4 + 8 - \frac{64}{12} \right) - \left(1 - 4 + \frac{8}{12} \right) \\ &= 15 - \frac{72}{12} \\ &= 9 \text{ square units.} \quad \checkmark \end{aligned}$$

$$(b) (i) \quad y = |x - 4|$$



$|x-4| = \frac{1}{2}x + c$ will have two distinct solutions when $c > -2$. \checkmark

(c) (i)

$$y = \frac{1}{8}x^2$$

$$\frac{dy}{dx} = \frac{1}{4}x$$

The gradient is t . $\frac{dy}{dx} = t$ So the tangent

$$T \text{ is } y - y_1 = m(x - x_1)$$

$$y - 2t^2 = t(x - 4t) \quad \checkmark$$

$$y - 2t^2 = tx - 4t^2$$

$$y = tx - 2t^2$$

$$(\text{ii}) \quad \text{Put } y = 0$$

$$tx = 2t^2$$

$$x = 2t. \text{ Hence } A = (2t, 0)$$

So the mid-point M is $(3t, t^2)$. \checkmark

$$(\text{iii}) \quad x = 3t \dots \textcircled{1}$$

$$y = t^2 \dots \textcircled{2}$$

From $\textcircled{1}$ $t = \frac{x}{3}$ substituting into $\textcircled{2}$ \checkmark

$$y = \frac{x^2}{9}$$

$$x^2 = 9y \quad \checkmark$$

This is a parabola with vertex $(0, 0)$ and focal length $\frac{9}{4}$. $\textcircled{15}$

QUESTION 7

a) To prove

$$\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{(n+1)(n+2)} = \frac{n}{2(n+2)}$$

A₁, when $n=1$, LHS = $\frac{1}{2 \times 3}$
 $= \frac{1}{6}$

LHS = $\frac{1}{2 \times 3}$
 $= \frac{1}{6}$ The statement is true
 for $n=1$.

B₁, Suppose k is a positive integer
 for which the statement is true.

That is $\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{(k+1)(k+2)} = \frac{k}{2(k+2)}$

We prove the statement true for
 $n=k+1$. That is we prove

$$\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{(k+1)(k+2)} + \frac{1}{(k+2)(k+3)} = \frac{k+1}{2(k+3)}$$

$$\begin{aligned} \text{LHS} &= \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{(k+1)(k+2)} + \frac{1}{(k+2)(k+3)} \\ &= \frac{k}{2(k+2)} + \frac{1}{(k+2)(k+3)} \quad (\text{by the induction hypothesis}) \end{aligned}$$

$$= \frac{k(k+3) + 2}{2(k+2)(k+3)}$$

$$= \frac{k^2 + 3k + 2}{2(k+2)(k+3)}$$

$$= \frac{(k+1)(k+2)}{2(k+2)(k+3)}$$

$$= \frac{k+1}{2(k+3)}$$

$$= \text{RHS.}$$

C₁, It follows from parts A and B by mathematical induction that the

statement is true for all positive integers. n.

(b)

$$V = \pi \int_{r-h}^r r^2 - x^2 \cdot dy \quad \checkmark$$

$$= \pi \left[r^2 x - \frac{x^3}{3} \right]_{r-h}^r \quad \checkmark$$

$$= \pi \left[\left(r^3 - \frac{r^3}{3} \right) - \left((r-h)^2(r-h) - \frac{(r-h)^3}{3} \right) \right]$$

$$= \pi \left[\frac{2r^3}{3} - \left(r^3 - r^2 h - \frac{1}{3}(r^3 - 3r^2 h + 3r h^2) \right) \right]$$

$$= \pi \left[\frac{2r^3}{3} - \left(\frac{2}{3}r^3 - r^2 h - \frac{1}{3}h^3 \right) \right]$$

$$= \pi \left[r^2 h - \frac{1}{3}h^3 \right]$$

$$V = \frac{\pi}{3} h^2 (3r - h) \quad \checkmark$$

$$(c) V = \frac{\pi}{3} h^2 (3r - h)$$

$$= 4\pi h^2 - \frac{\pi}{3} h^3, \text{ since } r=4.$$

$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt} \quad (\text{chain rule})$$


$$\frac{dV}{dt} = (8\pi h - \pi h^2) \frac{dh}{dt} \quad \checkmark$$

Substitute $\frac{dV}{dt} = 3\pi$ and $h=3$.

$$3\pi = (24\pi - 9\pi) \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{1}{5} \text{ m/hr.} \quad \checkmark$$

(d)



asymptote $y=0$.

15

QUESTION 8

$$(a) \quad y = \frac{1-x^2}{1+x^2}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{(1+x^2)(-2x) - (1-x^2)2x}{(1+x^2)^2} \\ &= \frac{-2x(1+x^2 + 1-x^2)}{(1+x^2)^2} \end{aligned}$$

$$\frac{dy}{dx} = \frac{-4x}{(1+x^2)^2} \quad \checkmark$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{(1+x^2)^2(-4) - (-4x)2(1+x^2)2x}{(1+x^2)^4} \\ &= \frac{-4(1+x^2)\{1+x^2 - 4x^2\}}{(1+x^2)^4} \\ &= \frac{4(3x^2-1)}{(1+x^2)^3} \quad \checkmark \end{aligned}$$

. So y'' has zeros

at $x = \frac{1}{\sqrt{3}}$ and $x = -\frac{1}{\sqrt{3}}$ and no discontinuities.

x	-1	$-\frac{1}{\sqrt{3}}$	0	$\frac{1}{\sqrt{3}}$	1
$\frac{d^2y}{dx^2}$	1	0	-4	0	1

The curve is concave down for
 $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$. \checkmark

$$\begin{aligned} (b) (i) \quad RHS &= \frac{1}{x} \left(\frac{1}{1+nx} - \frac{1}{1+(n+1)x} \right) \\ &= \frac{1}{x} \left(\frac{1+(n+1)x - (1+nx)}{(1+nx)(1+(n+1)x)} \right) \\ &\checkmark \quad = \frac{1}{x} \left(\frac{1+nx+x-1-nx}{(1+nx)(1+(n+1)x)} \right) \\ &= \frac{1}{x} \left(\frac{nx}{(1+nx)(1+(n+1)x)} \right) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{(1+nx)(1+(n+1)x)} \\ &= LHS. \end{aligned}$$

(ii)

$$\begin{aligned} &\frac{1}{(1+x)(1+2x)} + \frac{1}{(1+2x)(1+3x)} + \frac{1}{(1+3x)(1+4x)} + \dots \\ &+ \frac{1}{(1+nx)(1+(n+1)x)}. \quad \text{Using the result from part (i)} \end{aligned}$$

$$\begin{aligned} &= \frac{1}{x} \left(\frac{1}{1+x} - \frac{1}{1+2x} + \frac{1}{1+2x} - \frac{1}{1+3x} + \dots \right. \\ &\quad \left. + \frac{1}{1+nx} - \frac{1}{1+(n+1)x} \right) \quad \checkmark \\ &= \frac{1}{x} \left(\frac{1}{1+x} - \frac{1}{1+(n+1)x} \right) \quad \checkmark \\ &= \frac{1}{x} \left(\frac{1+(n+1)x - (1+nx)}{(1+nx)(1+(n+1)x)} \right) \\ &= \frac{1}{x} \times \frac{nx}{(1+nx)(1+(n+1)x)} \\ &= \frac{n}{(1+nx)(1+(n+1)x)} \quad \checkmark \end{aligned}$$

(c) (i) Let $HD = y$.

Since $V = x^2y$,
 $y = \frac{V}{x^2}$.

$$L = 4x + 4y$$

$$L = 4x + \frac{4V}{x^2} \quad \checkmark$$

$$\begin{aligned} (ii) \quad \frac{dL}{dx} &= 4 - \frac{8V}{x^3} \\ 4 - \frac{8V}{x^3} &= 0 \\ 4x^3 - 8V &= 0 \\ 4x^3 &= 8V \\ x &= (2V)^{\frac{1}{3}} \quad \checkmark \end{aligned}$$

so $\frac{dL}{dx}$ has a zero at $x = (2V)^{\frac{1}{3}}$.

$$\frac{dL}{dx} = 24Vx^{-4}$$

so $\frac{d^2L}{dx^2} > 0$ for all x , so

L has a minimum at $x = (2V)^{\frac{1}{3}}$.

ii) When $x = (2V)^{\frac{1}{3}}$,

$$L = 4x + \frac{4V}{x^2}$$

$$L = 4 \left\{ (2V)^{\frac{1}{3}} + \frac{V}{(2V)^{\frac{2}{3}}} \right\} \checkmark$$

$$= 4 \left\{ (2V)^{\frac{1}{3}} + 2^{-\frac{2}{3}} \cdot V^{\frac{1}{3}} \right\}$$

$$= 2^2 \cdot 2^{\frac{1}{3}} V^{\frac{1}{3}} + 2^2 \cdot 2^{-\frac{2}{3}} V^{\frac{1}{3}}$$

$$= 4(2V)^{\frac{1}{3}} + 2^{\frac{4}{3}} \cdot V^{\frac{1}{3}}$$

$$= 4(2V)^{\frac{1}{3}} + 2 \cdot 2^{\frac{1}{3}} V^{\frac{1}{3}}$$

$$= 6(2V)^{\frac{1}{3}} \checkmark$$

which is the minimum value of L .

i)

$$(i) a^5 - b^5 = (a-b)(a^4 + a^3b + a^2b^2 + ab^3 + b^4) \checkmark$$

$$(ii) 77^5 - 22^5 + 32^4 - 12^4$$

$$= (77-22)m + (32-12)n \checkmark$$

$$= 55m + 20n \text{ (where } m \text{ and } n \text{ are positive integers.)}$$

$$= 5(11m + 4n) \checkmark$$

the number is divisible by 5.