



# FORM VI

## MATHEMATICS EXTENSION 1

Examination date  
 Tuesday 10th August 2004

Time allowed  
 2 hours (plus 5 minutes reading time)

### Instructions

- All seven questions may be attempted.
- All seven questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

### Collection

- Write your candidate number clearly on each booklet.
- Hand in the seven questions in a single well-ordered pile.
- Hand in a booklet for each question, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Keep the printed examination paper and bring it to your next Mathematics lesson.

### Checklist

- SGS booklets: 7 per boy. A total of 1000 booklets should be sufficient.
- Candidature: 121 boys.

### Examiner

MLS

**QUESTION ONE** (12 marks) Use a separate writing booklet.

Marks


- (a) Solve the inequality  $\frac{4}{5-x} \leq 1$ . 3
- (b) For what value of  $p$  is the expression  $4x^3 - x + p$  divisible by  $x + 3$ ? 2
- (c) Expand  $(a + \frac{1}{2})^5$ , expressing each term in its simplest form. 2
- (d) Given the points  $A(1, 4)$  and  $B(5, 2)$ , find the co-ordinates of the point that divides the interval  $AB$  externally in the ratio 1 : 3. 2
- (e) Find  $\int x(1 - x^2)^5 dx$ , using the substitution  $u = 1 - x^2$ , or otherwise. 3

**QUESTION TWO** (12 marks) Use a separate writing booklet.

Marks

- (a) Consider the parabola  $x = 4t, y = 2t^2$ .
- (i) Find the gradient of the parabola at the point where  $t = 4$ . 1
- (ii) Find the equation of the tangent to the parabola at  $t = 4$ . 2

(b)



In the diagram above, two circles touch one another externally at the point  $A$ . A straight line through  $A$  meets one of the circles at  $T$  and the other at  $S$ . The tangents at  $T$  and  $S$  meet the common tangent at  $A$  at  $X$  and  $Y$  respectively.


Let  $\theta = \angle XTA$ .

- (i) Explain why  $\angle XAT$  is  $\theta$ . 1
- (ii) Prove that  $TX \parallel YS$ . 2
- (c) (i) Write down the first three terms in the expansion of  $(1 + mx)^n$ . 1
- (ii) If  $(1 + mx)^n \equiv 1 - 4x + 7x^2 - \dots$ , find the values of  $m$  and  $n$ . 3
- (d) Evaluate  $\lim_{x \rightarrow 0} \frac{5x \cos 2x}{\sin x}$ , showing your reasoning. 2

**QUESTION THREE** (12 marks) Use a separate writing booklet.

Marks

(a)



The diagram above shows a container in the shape of a right circular cone. The semi-vertical angle  $\theta = \tan^{-1} \frac{1}{2}$ .

Water is poured in at the constant rate of  $10 \text{ cm}^3$  per minute.

Let the height of the water at time  $t$  seconds be  $h$  cm, let the radius of the water surface be  $r$  cm, and let the volume of water be  $V \text{ cm}^3$ .

- (i) Show that  $r = \frac{1}{2}h$ . 1
- (ii) Show that  $V = \frac{1}{12}\pi h^3$ . 1
- (iii) Find the exact rate at which  $h$  is increasing when the height of the water in the cone is 50 cm. 2

(b) Show that there is no term independent of  $x$  in the expansion of  $\left(2x^2 - \frac{1}{4x}\right)^{11}$ . 3

(c) Evaluate  $\int_{-1}^0 x\sqrt{1+x} dx$ , using the substitution  $u = 1+x$ . 4

(d) Find  $\int \sin x \cos^3 x dx$ . 1

**QUESTION FOUR** (12 marks) Use a separate writing booklet.

Marks


(a) If  $y = \frac{1}{200}te^{-t}$ , show that  $\frac{dy}{dt} = \frac{1}{200}(1-t)e^{-t}$ . 1

(b) Fred has recently consumed three standard alcoholic drinks. Immediately after he has finished his last drink, his blood alcohol level is measured over a four-hour period. Let his blood alcohol level at any time  $t$  be  $A$ , where  $t$  is the time in hours after his last drink.

It is found that the rate of change  $\frac{dA}{dt}$  of his blood alcohol content is given by

$$\frac{dA}{dt} = \frac{1}{200}(1-t)e^{-t}, \text{ where } 0 \leq t \leq 4.$$

- (i) Show that his blood alcohol content increases during the first hour and decreases after the first hour. 2
- (ii) Initially his blood alcohol content was 0.0005. Find  $A$  as a function of  $t$ . You will need to use part (a). 2
- (iii) Determine his maximum alcohol content during the four-hour period. Give your answer correct to four decimal places. 1



The graph of the curve  $y = \frac{\pi}{2} - 2 \tan^{-1} x$  is drawn above. It cuts the  $y$ -axis at  $(0, \frac{\pi}{2})$ .

- (i) Write down the domain of the inverse function of  $y = \frac{\pi}{2} - 2 \tan^{-1} x$ . 1
- (ii) Find the equation of the inverse function of  $y = \frac{\pi}{2} - 2 \tan^{-1} x$ . 1
- (iii) Find the volume generated when the shaded region is rotated about the  $y$ -axis. 4

**QUESTION FIVE** (12 marks) Use a separate writing booklet.

Marks

(a) Evaluate  $\int_0^4 \frac{1}{3+\sqrt{x}} dx$ , using the substitution  $x = (u-3)^2$ . 3

(b) (i) Write down the expansion of  $(1+x)^n$  in ascending powers of  $x$ . Then differentiate both sides of your identity. 1

(ii) Make an appropriate substitution for  $x$  to show that 1

$$\binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + 4\binom{n}{4} + \dots + n\binom{n}{n} = n(2^{n-1}).$$

(iii) Hence find an expression for 1

$$2\binom{n}{1} + 3\binom{n}{2} + 4\binom{n}{3} + 5\binom{n}{4} + \dots + (n+1)\binom{n}{n}.$$

(c) Find values for  $R$  and  $\alpha$  if  $\sqrt{3}\sin\theta - \cos\theta = R\cos(\theta + \alpha)$ , where  $R$  and  $\alpha$  are positive constants and  $0 < \alpha < 2\pi$ . 2


(d) Use the method of mathematical induction to prove that 4

$$\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{n}{(n+1)!} = \frac{(n+1)! - 1}{(n+1)!}, \text{ for all positive integers } n.$$

**QUESTION SIX** (12 marks) Use a separate writing booklet.

Marks

(a)



The sketch above shows the curve  $y = 2\sin x$  and the line  $x - 3y = 0$ . The graphs meet at the point  $A$  in the first quadrant.

(i) Write down an equation whose solution gives the  $x$ -coordinate of  $A$ . 1

(ii) An approximate value for the  $x$ -coordinate of  $A$  is  $x = 3$ . Apply Newton's method once to find a closer approximation for this value. Give your answer correct to one decimal place. 2

(b) Newton's law of cooling states that a body cools according to the equation

$$\frac{dT}{dt} = -k(T - S),$$

where  $T$  is the temperature of the body at time  $t$ ,  $S$  is the temperature of the surroundings and  $k$  is a constant.


(i) Show that  $T = S + Ae^{-kt}$  satisfies the equation, where  $A$  is a constant. 1

(ii) A metal rod has an initial temperature of  $470^\circ\text{C}$  and cools to  $250^\circ\text{C}$  in 10 minutes. The surrounding temperature is  $30^\circ\text{C}$ .

(a) Find the value of  $A$  and show that  $k = \frac{1}{10} \log_e 2$ . 2

(b) Find how much longer it will take the rod to cool to  $70^\circ\text{C}$ , giving your answer correct to the nearest minute. 2

(c)



In the diagram above, the straight line  $ACD$  is a tangent at  $A$  to the circle with centre  $O$ . The interval  $AOB$  is a diameter of the circle. The intervals  $BC$  and  $BD$  meet the circle at  $E$  and  $F$  respectively.

Let  $\angle BAF = \beta$ .

Copy or trace this diagram into your answer booklet.

(i) Explain why  $\angle ABF = \frac{\pi}{2} - \beta$ . 1

(ii) Prove that the quadrilateral  $CDFE$  is cyclic. 3

**QUESTION SEVEN** (12 marks) Use a separate writing booklet. Marks

(a) Car A and car B are travelling along a straight level road at constant speeds  $V_A$  and  $V_B$  respectively. Car A is behind car B, but is travelling faster.

When car A is exactly  $D$  metres behind car B, car A applies its brakes, producing a constant deceleration of  $k \text{ m/s}^2$ .

- (i) Using calculus, find the speed of car A after it has travelled a distance  $x$  metres under braking. 2
  - (ii) Prove that the cars will collide if  $V_A - V_B > \sqrt{2kD}$ . 4
- (b) A particle is moving in simple harmonic motion of period  $T$  about a centre  $O$ . Its displacement at any time  $t$  is given by  $x = a \sin nt$ , where  $a$  is the amplitude.
- (i) Draw a neat sketch of one period of this displacement-time equation, showing all intercepts. 1
  - (ii) Show that  $\dot{x} = \frac{2\pi a}{T} \cos \frac{2\pi t}{T}$ . 1
- (c) The point  $P$  lies  $D$  units on the positive side of  $O$ . Let  $V$  be the velocity of the particle when it first passes through  $P$ . 4
- Show that the time between the first two occasions when the particle passes through  $P$  is  $\frac{T}{\pi} \tan^{-1} \frac{VT}{2\pi D}$ .

END OF EXAMINATION

Q1. a

(i)  $\frac{4}{5-x} \leq 1$

$4(5-x) \leq (5-x)^2$  ✓

$20 - 4x \leq 25 - 10x + x^2$

$x^2 - 6x + 5 \geq 0$  ✓



$1 \leq x \leq 5$  ✓

(ii)  $p(x) = 4x^3 - x + p$

$p(-3) = 0$  ✓

$4(-27) - 3 + p = 0$  ✓

$p = 105$  ✓

(iii)  $(a + \frac{1}{2})^5$   
 $= \binom{5}{0} a^5 + \binom{5}{1} a^4 \cdot \frac{1}{2} + \binom{5}{2} a^3 \cdot \frac{1}{4} + \binom{5}{3} a^2 \cdot \frac{1}{8} + \binom{5}{4} a \cdot \frac{1}{16} + \binom{5}{5} \frac{1}{32}$  ✓

(iv) A(1, 4), B(5, 2)  $\vec{AB}$  external

$x = \frac{m(1) - n(5)}{m - n}$   $y = \frac{m(4) - n(2)}{m - n}$

$\frac{5 - 3}{-3}$   $\frac{2 - 12}{-3}$  ✓

$= -1$   $= 5$  ✓

$(-1, 5)$

(12)

(v)  $\int k(1-x^2)^5 dx$

Let  $u = 1 - x^2$

$\frac{du}{dx} = -2x$  ✓

$I = -\frac{1}{2} \int u^5 \cdot \frac{du}{x}$  ✓

$= -\frac{1}{2} \int u^5 du$  ✓

$= -\frac{1}{2} \left[ \frac{u^6}{6} \right] + C$  ✓

$= -\frac{(1-x^2)^6}{12} + C$

(a)  $x = 4t, y = 2t^2$   
 $t = \frac{x}{4}$   
 $y = 2 \left(\frac{x}{4}\right)^2$   
 $y = \frac{2x^2}{16}$   
 $y = \frac{x^2}{8}$


(b)  $\frac{dy}{dx} = \frac{x}{4}$

when  $t = 4, x = 16, y = 32$   
 $m = 4$

(c) when  $t = 4, x = 16, y = 32$

$4 - 32 = 4(x - 16)$   
 $4 - 32 = 4x - 64$   
 $4 = 4x - 32$  ✓

Quicker to use "chain rule"  
 i.e.  $x = 4t \Rightarrow \frac{dx}{dt} = 4$   
 $y = 2t^2 \Rightarrow \frac{dy}{dt} = 4t$   
 $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$   
 $= 4t \times \frac{1}{4} = C$  At  $t = 4, m = 4$ .



(i) Tangents from an external point X are equal to points on the circle (i.e. XA & XT)

$$\therefore XA = XT \quad (\text{Sine rule } AX = XT)$$

(ii)  $\angle AS = \theta$  (vert. opposite)

$AY = YS$  (tangents from external point)

$$\therefore AY = YS$$

$\therefore XT \parallel YS$  (Sine alternate angles are equal)

(iii)  $(1+mx)^n = \binom{n}{0} 1 + \binom{n}{1} mx + \binom{n}{2} m^2 x^2 + \dots$

(i)  $\binom{n}{1} mx = -4x \implies \frac{n!}{(n-1)!} m = -4 \implies nm = -4$

(ii)  $\binom{n}{2} m^2 x^2 = 7x^2 \implies \frac{n!}{(n-2)!} m^2 = 7 \implies n(n-1)m^2 = 7$

$\frac{nm}{n-1} = -4$   
 $nm = -4(n-1)$   
 $m = \frac{-4(n-1)}{n}$

$\frac{n!}{(n-2)!} m^2 = 7$   
 $n(n-1)m^2 = 7$   
 $n(n-1) \left(\frac{-4(n-1)}{n}\right)^2 = 7$   
 $n(n-1) \cdot 16(n-1)^2 = 7n^2$   
 $16(n-1)^2 = 7n$   
 $16n^2 - 32n + 16 = 7n$   
 $16n^2 - 39n + 16 = 0$   
 $n = \frac{39 \pm \sqrt{39^2 - 4 \cdot 16 \cdot 16}}{2 \cdot 16}$   
 $n = \frac{39 \pm \sqrt{1521 - 1024}}{32}$   
 $n = \frac{39 \pm \sqrt{497}}{32}$

$\frac{16}{n} (n-1) = 14$

$16n - 16 = 14n$   
 $2n = 16$   
 $n = 8$

$m = -\frac{4}{3}$

(iv)  $\lim_{x \rightarrow 0} \frac{5x \cos 2x}{\sin x}$


$\cos 2x = (1 - \sin^2 x) - \sin^2 x = 1 - 2\sin^2 x$

$= \lim_{x \rightarrow 0} \frac{5x \cdot (1 - 2\sin^2 x)}{\sin x}$

Note:  $5 \lim_{x \rightarrow 0} \frac{x}{\sin x} \times \lim_{x \rightarrow 0} \frac{\cos 2x}{1}$

$= \lim_{x \rightarrow 0} \frac{5x}{\sin x} = 5 \lim_{x \rightarrow 0} \frac{x}{\sin x} = 5 \times 1 = 5$

$= \lim_{x \rightarrow 0} 5 \times \frac{x}{\sin x} = \lim_{x \rightarrow 0} 10x \sin x = 5$



By Similar Triangles (ii)  $V = \frac{1}{3} \pi r^2 h$   
 $= \frac{1}{3} \pi \left(\frac{r}{h}\right)^2 h$   
 $= \frac{1}{3} \pi \frac{r^2}{h}$   
 $r = \frac{1}{2} h$

(ii)  $\frac{dV}{dt} = 10$

$\frac{dV}{dh} = \frac{\pi h^2}{4}$

$\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}$   
 $= 10 \times \frac{4}{\pi h^2}$

at  $h = 50$

$\frac{dh}{dt} = \frac{4}{2500\pi} = \frac{4}{250\pi} = \frac{2}{125\pi}$

(b)  $(2x^2 - \frac{1}{4x})^n$

$T_{r+1} = \binom{n}{r} (2x^2)^{n-r} \left(-\frac{1}{4x}\right)^r$   
 $= \binom{n}{r} 2^{n-r} 4^{-r} x^{2n-2r} \cdot x^{-r}$   
 $= \binom{n}{r} 2^{n-r} 4^r x^{2n-3r}$

For independent of x,  $2n-3r=0$

$2n-3r=0$

$2n=3r$

$r = \frac{2n}{3}$

$\therefore$  No Independent Term since r must be an integer

(c)  $\int_{-1}^0 x \sqrt{x+1} dx$

Let  $u = 1+x \implies x = u-1$  At  $x=0, u=1$   
 $du = dx$  At  $x=-1, u=0$

$\int_0^1 (u-1)u^{1/2} du$   
 $= \int_0^1 u^{3/2} - u^{1/2} du$   
 $= \left[ \frac{2u^{5/2}}{5} - \frac{2u^{3/2}}{3} \right]_0^1$   
 $= \frac{2}{5} - \frac{2}{3}$   
 $= \frac{4}{15}$

(d)  $\int \sin x \cdot \cos^3 x dx$

$= -\frac{\cos^4 x}{4} + c$

Question 4.  
 (a)  $y = \frac{1}{200} t e^{-t}$   $u = t$   $v = e^{-t}$   
 $u' = 1$   $v' = -e^{-t}$

$\frac{dy}{dt} = \frac{1}{200} [u'v + uv']$   
 $= \frac{1}{200} [e^{-t} + t(-e^{-t})]$   
 $= \frac{1}{200} e^{-t} [1 - t]$

(b) At  $t=0, y=0$   
 $t=1, y = \frac{1}{200e} = 0.0018$   
 $t=2, y = \frac{2}{200e^2} = 0.0015$   
 $\therefore$  Increases at  $t=1$ , then decreases at  $t=2$ .

(ii)  $A = \frac{1}{200} t e^{-t}$   
 $t=0, y = 0.0005$   
 $0.0005 = \frac{1}{200} e$   
 $A = 0.0005 + \frac{t e^{-t}}{200}$  ✓

(iii)  $t=4$   
 $A = 0.0005 + \frac{4}{200 e^4}$   
 $= 0.0009$  (4 dp).  
 Max. alcohol content is at  $t=1$   
 $\therefore A = \frac{1}{200} \cdot 1 \cdot e^{-1}$   
 $= 0.0018$  (to 4 dp).

(c) (i)  $y = \pi/2 - 2 \tan^{-1} x$   
 D: All real  $x$  R:  $-\pi/2 < y < \pi/2$   
 For its inverse function  $D: -\pi/2 < x < \pi/2$

(ii)  $y = \pi/2 - 2 \tan^{-1} x$   
 $\Rightarrow \tan^{-1} x = \frac{\pi/2 - y}{2}$   
 $x = \tan(\pi/4 - y/2)$   
 $x^2 = \tan^2(\pi/4 - y/2)$  ✓

(iii)  $y = \pi/2 - 2 \tan^{-1} x$   
 $x = \tan(\pi/4 - y/2)$   
 $V = \pi \int_0^{\pi/2} \tan^2(\pi/4 - y/2) dy$   
 $= \pi \int_0^{\pi/2} \sec^2(\pi/4 - y/2) - 1 dy$  ✓  
 $= \pi \left[ \frac{1}{\pi/4} \tan(\pi/4 - y/2) - y \right]_0^{\pi/2}$   
 $= -\pi \left[ \frac{1}{\pi/4} \tan(\pi/4 - \pi/4) - \frac{\pi}{2} \right] - \left[ \frac{1}{\pi/4} \tan \frac{\pi}{4} - 0 \right]$   
 $= -\pi \left\{ \frac{\pi}{4} - \frac{\pi}{2} \right\}$   
 $= \pi \left\{ \frac{\pi}{2} - \frac{\pi}{4} \right\} = \frac{\pi}{2} \{4 - \pi\}$

Question 5.  
 (a)  $\int_0^4 \frac{1}{6.3 + \sqrt{u}} du$   
 Let  $u = (u-3)^2$   
 $\frac{du}{du} = 2(u-3)$   
 $dx = 4, u = 15$   
 $x=0, u=3$

$I = \int_3^{15} \frac{2(u-3) du}{3 + u - 3}$  ✓  
 $= \frac{2}{2} \int_3^{15} \frac{(u-3) du}{u}$   
 $= \frac{2}{2} \int_3^{15} \left( 1 - \frac{3}{u} \right) du$   
 $= \frac{2}{2} \left[ u - 3 \ln u \right]_3^{15}$   
 $= \frac{2}{2} \left[ 15 - 3 \ln 15 - 3 + 3 \ln 3 \right]$   
 $= 2 \left[ 12 - 3 \ln \left( \frac{15}{3} \right) \right]$   
 $= 2 \left[ 12 - 3 \ln 5 \right]$

(a) (i)  $(1+k)^n = 1 + (n)k + (n/2)k^2 + (n/6)k^3 + \dots + (n/2)k^k + \dots + (n)k^n$   
 $\frac{d}{dk} n(1+k)^{n-1} = (n) + 2k(n/2) + 3k^2(n/6) + \dots + k^{k-1}(n/2) + \dots + n(1+k)^{n-1}$   
 (ii)  $LHS = n(1+k)^{n-1} = (n) + 2k(n/2) + 3k^2(n/6) + \dots + n(1+k)^{n-1}$

(iii) Add  $(n) + (n) + (n) + \dots + (n)$   
 $LHS = (n) + (n) + \dots + (n) + n(1+k)^{n-1}$   
 $= 2^n - 1 + n(1+k)^{n-1}$   
 $= 2^n - 1 + n(1+k)^{n-1}$  ✓

(b)  $\sqrt{5} \sin \theta - \cos \theta$   
 $= 2 \cos(\theta + \pi/6)$   
 $n=1, \theta = \pi/6$   
 $= 2 \cos(\pi/6 + \pi/6)$  ✓

(c)  $\frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{(n+1)!} = \frac{(n+1)! - 1}{(n+1)!}$   
 Assume true for  $n=k$   
 $LHS = \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{(k+1)!} = \frac{(k+1)! - 1}{(k+1)!}$   
 Assume true for  $n=k+1$   
 $LHS = \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{(k+2)!} = \frac{(k+2)! - 1}{(k+2)!}$

$\frac{(k+2)! - 1}{(k+2)!} = \frac{(k+2)! - 1}{(k+2)!} + \frac{k+1}{(k+2)!}$   
 $= \frac{(k+2)[(k+1)! - 1] + k+1}{(k+2)!}$   
 $= \frac{(k+2)! - k+2 + k+1}{(k+2)!}$   
 $= \frac{(k+2)! - 1}{(k+2)!}$  ✓  
 If true for  $n=k$ , and  $n=k+1$ , therefore by the principles of Mathematical Induction, true for all positive integers.


Question 6.  
 (a) (i)  $x - 3y = 0, y = 2 \sin x$   
 $xy = x$   
 $y = \pi/3$   
 $2 \sin x = \pi/3$   
 $\sin x = \pi/6$  ✓  
 $f(x) = 6 \cos x - 1$

(ii)  $x_1 = 3$   
 $x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$   
 $= 3 - \frac{(-2 + 153)}{-6.439}$   
 $= 2.8$  (1 dp)


(b) (i)  $T = S + Ae^{-kt}$   
 $T - S = Ae^{-kt}$   
 $\frac{dT}{dt} = -Ake^{-kt}$   
 $= -k[T - S]$

(ii) (a)  $t = 0, T = 450$   
 $450 = 30 + Ae^0$   
 $A = 420$   
 $t = 10, T = 250$   
 $250 = 30 + 420e^{-10k}$   
 $\frac{220}{420} = e^{-10k}$   
 $\ln\left(\frac{11}{21}\right) = -10k$   
 $-\ln 2 = -10k$   
 $k = \frac{\ln 2}{10}$


(b)  $70 = 30 + 420e^{-\frac{\ln 2}{10}t}$   
 $\frac{40}{420} = 2^{-\frac{t}{10}}$   
 $\ln\left(\frac{2}{21}\right) = -\frac{t \ln 2}{10}$   
 $t = -\frac{10 \ln\left(\frac{2}{21}\right)}{\ln 2}$   
 $= 26.44 \text{ hrs}$   
 $= 26 \text{ hrs } 26 \text{ mins}$



(i)  $\angle AEB = \angle AFB = \theta$  (Angle in the same arc)  
 $\angle AEB + \angle AFB = \pi - \theta$  (Angle sum of the  $\Delta$ )  
 $\angle AEB = \pi - \theta$  (Radius to the tangent)  
 $\angle ADB = \pi - (\pi - \theta) = \theta$  (Angle sum)  
 $\angle BEF = \theta$  (Angle subtended by the same arc)  
 $\angle EAF = \pi - \theta$  (Supplementary angle)  
 $\angle CEF + \angle CDF = \pi$   
 $\therefore \angle CEF$  is cyclic (Opposite angles =  $\pi$ )




(i) At  $x = a, \dot{x} = -k \text{ m/s}$   
 $\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -k$   
 $\frac{1}{2}v^2 = -kx + C$   
 $v^2 = -2kx + C$   
 At  $x = a, v = V_A$   
 $(V_A)^2 + 2ka = C$



(ii)  $x = 0$   
 $0 = a \sin(\omega t)$   
 $\sin^{-1}\left(\frac{0}{a}\right) = \omega t$   
 $t = \frac{\sin^{-1}\left(\frac{0}{a}\right)}{\omega}$

$T = \frac{2\pi}{\omega}$   
 $\omega = \frac{2\pi}{T}$

7(a)(ii)  $\dot{x} = -k$   
 $\frac{dv}{dt} = -k$   
 $v = \int_0^t -k dt$   
 $[v]_0^t = [-kt]_0^t$   
 $v - v_0 = -kt$   
 $v_B - v_A = -kt$   
 $t = -\frac{v_B - v_A}{k}$



$\dot{x} = -k$   
 $\frac{d}{dx}\left(\frac{1}{2}v^2\right) = -k$   
 $\frac{1}{2}v^2 = -kx + C$   
 At  $x = 0, v = V_A$   
 $\therefore C = \frac{1}{2}V_A^2$   
 $\therefore \frac{1}{2}v^2 = -kx + \frac{1}{2}V_A^2$   
 $\therefore v^2 = -2kx + V_A^2$

(ii) When  $x = D$  and  $v = V_B$ , the cars will collide  
 $V_B^2 = -2kD + V_A^2$   
 $\therefore V_A^2 - V_B^2 = 2kD$   
 But  $(V_A - V_B)^2 > V_A^2 - V_B^2 = 2kD$   
 $\therefore V_A - V_B > \sqrt{2kD}$  as reqd.

B. Speed = Distance B  
 Time

$V_B t = D_B = \dots$   
 $t = \frac{D_B + D}{V_B}$   
 $\therefore -\frac{V_B + V_A}{k} = \frac{D_B + D}{V_B}$   
 $V_B [V_A - V_B] = D_B k + D$

$V_A - V_B$

P.T.O.