

## 2 UNIT MATHEMATICS FORM VI

Time allowed: 3 hours (plus 5 minutes reading)

Exam date: 8th August 2001

## Instructions:

- All questions may be attempted.
- All questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

## Collection:

- Each question will be collected separately.
- Start each question in a new 4-leaf answer booklet.
- If you use a second booklet for a question, place it inside the first. Don't staple.
- Write your candidate number on each answer booklet.

$$b) (i) \frac{1}{(x-1)x} = \frac{Ax + B(x-1)}{(x-1)x}$$

$$\therefore 1 = Ax + Bx - B$$

$$B = -1 \text{ and } A + B = 0$$

$$\therefore A = 1$$

✓✓

$$\therefore \frac{1}{(x-1)x} = \frac{1}{x-1} - \frac{1}{x}$$

$$(ii) S_n = \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{(n-1) \times n}$$

$$= \left(\frac{1}{1} - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \dots + \left(\frac{1}{n-1} - \frac{1}{n}\right)$$

$$= \frac{1}{1} + \left(-\frac{1}{2} + \frac{1}{2}\right) + \left(-\frac{1}{3} + \frac{1}{3}\right) + \dots + \left(\frac{-1}{n-1} + \frac{1}{n-1}\right) - \frac{1}{n}$$

$$= 1 - \frac{1}{n}, \text{ as required.}$$

✓  
✓

$$(iii) \sum_{n=2}^{\infty} \frac{1}{(n-1)n} = \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots$$

$$= \lim_{n \rightarrow \infty} S_n \text{ (from (ii))}$$

$$= \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)$$

$$= 1.$$

✓

✓

**QUESTION ONE** (Start a new answer booklet)

Marks


- 1 (a) Convert  $\frac{4\pi}{5}$  to degrees.
- 1 (b) Write down a primitive of  $\sec^2 5x$ .
- 2 (c) The line  $5x - ky = 7$  passes through the point  $(1, 1)$ . Find the value of  $k$ .
- 2 (d) Differentiate  $y = 5x^3 - 2x + 9$  with respect to  $x$ .
- 2 (e) Express  $\frac{4}{\sqrt{3}-1}$  with a rational denominator in simplest form.
- 2 (f) Find the exact value of  $\tan \frac{\pi}{3} + \tan \frac{\pi}{4}$ .
- 2 (g) Solve  $|x - 1| = 11$ .

**QUESTION TWO** (Start a new answer booklet)

Marks

- 2 (a) Differentiate the following with respect to  $x$ :
  - (i)  $x^2 e^x$ ,
  - (ii)  $\ln(3x - 2)$ ,
  - (iii)  $\sin^2 x$ .
- 2 (b) Find a primitive function of  $(3x - 4)^6$ .
- (c) Evaluate the following definite integrals:
  - 2 (i)  $\int_1^2 6x^2 dx$ ,
  - 2 (ii)  $\int_0^{\frac{\pi}{2}} \sin 2x dx$ .

**QUESTION 10**



(i)

$$\cos \theta = \frac{AP}{l}$$

$$\therefore AP = l \cos \theta$$

$$S = \frac{1}{2} \cdot l \cos \theta \cdot l \cos \theta \cdot \sin \theta$$

$$S = \frac{l^2}{2} \cdot \cos^2 \theta \cdot \sin \theta, \text{ as required.}$$

$$= \frac{l^2}{2} \cdot (1 - \sin^2 \theta) (\sin \theta)$$

$$= \frac{l^2}{2} (\sin \theta - \sin^3 \theta)$$

(ii)  $\frac{dS}{d\theta} = \frac{l^2}{2} (\cos \theta - 3\sin^2 \theta \cos \theta) = \frac{l^2}{2} \cos \theta (1 - 3\sin^2 \theta)$

$\frac{dS}{d\theta} = 0$  when  $\cos \theta = 0$  or  $\sin \theta = \pm \frac{1}{\sqrt{3}}$

Since  $0 < \theta < 90^\circ$ ,  $\sin \theta = \frac{1}{\sqrt{3}}$  is only possible solution.

$\sin \theta$	$\frac{1}{2}$	$\frac{1}{\sqrt{3}}$	$\frac{3}{4}$
$\frac{dS}{d\theta}$	+	0	-

$\therefore$  When  $\sin \theta = \frac{1}{\sqrt{3}}$ ,  $S$  is a maximum.

$$S = \frac{l^2}{2} \cdot \left( \frac{1}{\sqrt{3}} - \frac{1}{3\sqrt{3}} \right)$$

$$= \frac{l^2}{2} \cdot \frac{(3-1)}{3\sqrt{3}}$$

$$= \frac{l^2}{3\sqrt{3}}$$

$$= \frac{\sqrt{3}l^2}{9} \text{ is the maximum area.}$$

QUESTION 9

a) (i)  $V = Ce^{-kt}$   
 When  $t = 0$ ,  $V = 65000$   
 $\therefore C = 65000$  ✓  
 When  $t = 1$ ,  $V = 55000$   
 $\therefore 55000 = 65000 e^{-k}$   
 $e^{-k} = \frac{11}{13}$

$-k = \ln\left(\frac{11}{13}\right)$

$\therefore k = -\ln\left(\frac{11}{13}\right)$  } ✓ (either)  
 $\approx 0.16705...$

(ii) When  $t = 5$ ,  
 $V = 65000 e^{-5k}$   
 $= \$28194$  ✓

(iii) We need  $t$  such that  
 $V < \frac{65000}{2}$

i.e)  $65000 e^{-kt} < \frac{65000}{2}$

$e^{-kt} < \frac{1}{2}$  } ✓ (any)  
 $-kt < \ln \frac{1}{2}$   
 $t > 4.149...$

$\therefore$  Car falls below half its cost price in 2005 ✓

b) (i)  $R = 9t^2 - t^4$   
 When  $t = 2$ ;  $R = 9 \cdot 4 - 2^4$   
 $= 20 \text{ m}^3/\text{h}$  ✓

(ii) When  $t = 3$ ,  $R = 0$  and when  $t > 3$ ,  $R < 0$  which suggests that concrete is going back into the truck! Since  $t \geq 0$  we have  $0 \leq t \leq 3$

(iii)  $\frac{dR}{dt} = 18t - 4t^3$  and  $\frac{d^2R}{dt^2} = 18 - 12t^2$   
 $= 2t(9 - 2t^2)$  ✓

$\frac{dR}{dt} = 0$  when  $t = 0$  or  $\frac{3}{\sqrt{2}}$  or  $-\frac{3}{\sqrt{2}}$

Since  $t \geq 0$ , ignore  $t = -\frac{3}{\sqrt{2}}$ . When  $t = \frac{3}{\sqrt{2}}$ ,  $\frac{d^2R}{dt^2} > 0$  and when  $t = \frac{3}{\sqrt{2}}$ ,  $\frac{d^2R}{dt^2} < 0$ .

So maximum flow-rate occurs when  $t = \frac{3}{\sqrt{2}}$ .  
 $\bar{R} = 9 \cdot \frac{9}{2} - \frac{81}{4}$  ✓


(iv)  $A = \int 9t^2 - t^4 dt = \frac{81}{4} \text{ m}^3/\text{h}$  ✓  
 $A = 3t^3 - \frac{t^5}{5} + c$  ✓

When  $t = 0$ ,  $A = 1000 \therefore c = 1000$ .

$\therefore A = 3t^3 - \frac{t^5}{5} + 1000$  ✓

(Be generous in part (ii).)

QUESTION THREE (Start a new answer booklet)




Marks

- 2 (i) Find the length of  $BC$  correct to the nearest centimetre.
- 2 (ii) Find the area of  $\triangle ABC$  correct to the nearest square metre.

(b) Consider the geometric series  $1 - \frac{1}{3} + \frac{1}{9} - \dots$

- 1 (i) Explain why the series has a limiting sum.
- 1 (ii) Find the limiting sum.
- 3 (c) Find the equation of the tangent to the curve  $y = \ln x$  at the point  $(e, 1)$ .
- (d) If  $\alpha$  and  $\beta$  are roots of the equation  $x^2 + 8x + 11 = 0$ , find:
  - 1 (i)  $\alpha + \beta$ ,
  - 1 (ii)  $\alpha\beta$ ,
  - 1 (iii)  $\alpha^2 + \beta^2$ .

QUESTION FOUR (Start a new answer booklet)




In the diagram above, the graph of  $y = f(x)$  is drawn.

Marks

- 1 (i) Sketch the graph of  $y = f(x) + 2$ .
- 1 (ii) Given that  $\int_{-1}^3 f(x) dx = \frac{15}{2}$ , evaluate  $\int_{-1}^3 (f(x) + 2) dx$ .

(b) A particle moves in a straight line so that its displacement  $x$  metres at time  $t$  seconds is given by  $x = 2t^3 - t^2$ .


- 2 (i) At what times is the particle at rest?
- 2 (ii) How far does the particle travel between these times?



In the diagram above,  $\triangle ABC$  is isosceles with  $\angle B = \angle C = 40^\circ$ , and  $AD = AE$ . Let  $\angle DAC = \theta$ .

- 1 (i) Explain why  $\angle ADB = 40^\circ + \theta$ .
- 2 (ii) Find an expression for  $\angle DAE$  in terms of  $\theta$ .
- 3 (iii) Show that  $\angle EDB = \frac{1}{2}\theta$ .

**QUESTION FIVE** (Start a new answer booklet)



In the diagram above,  $O$  is the centre of a circle of radius 4.8 centimetres, and  $\angle AOB = 75^\circ$ .

- Marks
- 2 (i) Find the exact length of arc  $AB$ .
  - 2 (ii) Find the exact area of the sector  $AOB$ .
  - 2 (iii) Find the area of the minor segment that has been shaded. Give your answer correct to three decimal places.

**QUESTION 8**

- a) (i)  $A_1 = 1500 (1.0075) = \$1511.25$  ✓
- (ii)  $A = 1500 (1.0075)^{120} = \$3677.04$  ✓
- (iii) Total Amount =  $1500 [1.0075^{120} + 1.0075^{119} + 1.0075^{118} + \dots + 1.0075]$  ✓
- $= 1500 \left[ 1.0075 \frac{(1.0075^{120} - 1)}{(1.0075 - 1)} \right]$  ✓
- $= 1500 \times 194.9656342$  ✓
- $= \$292448$  ✓
- (iv) New Total =  $1600 \times 194.9656342 = \$311945$  ✓
- $\therefore$  Difference =  $\$19497$  ✓
- b) (i) Area ABCD =  $\frac{1}{2} \cdot 2 \left(x + \frac{1}{e}\right)$  ✓
- $= \frac{e^2 + 1}{e} (u^2)$  ✓
- (ii) Area under the curve =  $\int_1^e e^x dx$  ✓
- $= [e^x]_1^e$
- $= e - e^{-1}$
- $= e - \frac{1}{e}$
- $= \frac{e^2 - 1}{e} (u^2)$  ✓
- $\therefore$  Area between the curves =  $\frac{e^2 + 1}{e} - \frac{e^2 - 1}{e}$
- $= \frac{2}{e} (u^2)$  ✓

QUESTION 7

a)  $\angle BWA = 80^\circ$  (external  $\angle$  of  $\triangle AWD$ ) ✓  
 $\angle BXC = 56^\circ$  (external  $\angle$  of  $\triangle CXE$ ) ✓  
 $x = 180 - (80 + 56)$  ( $\angle$  sum of  $\triangle$ ) ✓  
 $\therefore x = 44$  ✓

b)  $\angle DPA = x$  (alternate  $\angle$  on  $\parallel$  lines) ✓  
 $\therefore \triangle ADP$  is isosceles  $\Rightarrow AD = AP$  — 1. ✓  
 $\angle BPC = y$  (alternate  $\angle$  on  $\parallel$  lines) ✓  
 $\therefore \triangle BCP$  is isosceles  $\Rightarrow BC = BP$  — 2. ✓

Since ABCD is a parallelogram,  $AD = BC$  — 3.  
 By 1, 2 & 3;  $AD = AP = BC = BP$ .

$\therefore AB = AP + BP$   
 $= AD + AD$   
 $= 2AD$ , as required.

✓✓ (There are other acceptable methods. Allocate marks similarly if possible)


c) (i)  $\Delta = 2 - 4.5k$  ✓  
 $= 4 - 20k$  ✓  
 (ii) For real roots,  $\Delta \geq 0$  } ✓ (either)  
 $\therefore 4 - 20k \geq 0$  }  
 $\therefore k \leq \frac{1}{5}$  ✓

d)  $\ln 16 = 2 \ln x$  ✓  
 $\therefore 2 \ln 4 = 2 \ln x$  ✓  
 $\therefore x = 4$  ✓

[or  $\ln 16 = \ln x^2$  ✓  
 $\therefore x^2 = 16$  ✓  
 $x = \pm 4$   
 but  $x > 0 \therefore x = 4$  only] ✓

- 2 (b) (i) Solve  $\tan x = -3$  for  $0 \leq x \leq 2\pi$ . Give your answer in radians correct to three decimal places.  
 2 (ii) On the same diagram, sketch graphs of  $y = \tan x$  and  $y = -3$  for  $0 \leq x \leq 2\pi$ .  
 2 (iii) How many solutions are there to the equation  $\tan x = -3$  in the domain  $-2\pi \leq x \leq 2\pi$ ?

QUESTION SIX (Start a new answer booklet)




In the diagram above, the graph of  $y = 3x^2 + 2x - 1$  and the tangent to the curve at the point  $A(-1, 0)$  are drawn.


- Marks  
 2 (i) Show that the equation of the tangent is  $y + 4x + 4 = 0$ .  
 1 (ii) Show that the tangent meets the y-axis at  $B(0, -4)$ .  
 2 (iii) Find the equation of the line that passes through B and which is perpendicular to the tangent.  
 1 (iv) Show that this line meets the x-axis at the point  $C(16, 0)$ .  
 2 (v) Find the area of  $\triangle ABC$ .  
 4 (b) The region bounded by the curve  $y = \tan x$  and the x-axis from  $x = 0$  to  $x = \frac{\pi}{4}$  is rotated about the x-axis. The volume of the solid formed is given by  $V = \pi \int_0^{\frac{\pi}{4}} \tan^2 x dx$ . Use Simpson's rule with the three function values  $x = 0, \frac{\pi}{8}$  and  $\frac{\pi}{4}$  to approximate the volume. Give your answer correct to three decimal places.

**QUESTION SEVEN** (Start a new answer booklet)

Marks  
3 (a)



Find the value of  $x$ . (Give reasons.)



In the diagram above,  $ABCD$  is a parallelogram. The point  $P$  lies on  $AB$  and it is known that  $\angle ADP = \angle CDP = x$  and  $\angle BCP = \angle DCP = y$ .  
Prove that  $2AD = AB$ . (Give reasons.)

- 1 (c) (i) Write down the discriminant of  $5x^2 - 2x + k$ .
- 2 (c) (ii) For what values of  $k$  does  $5x^2 - 2x + k = 0$  have real roots?
- 2 (d) Solve  $\log_e 16 = 2 \log_e x$ .

**QUESTION 6**

a) (i)  $y = 3x + 2x - 1$   
 $\frac{dy}{dx} = 6x + 2$

At  $x = -1$ ,  $\frac{dy}{dx} = -4$  ✓

Eqn is  $y - 0 = -4(x + 1)$   
 i.e)  $y + 4x + 4 = 0$  ✓

(ii) When  $x = 0$ ,  $y + 4 = 0$   
 $\therefore y = -4$  ✓

So  $B = (0, -4)$  ✓

(iii)  $m = \frac{1}{4}$  ✓

$y + 4 = \frac{1}{4}(x - 0)$  ✓

i.e)  $4y + 16 = x$  ✓

(iv) When  $y = 0$ ,  $x = 16$   
 $\therefore C = (16, 0)$  ✓

(v) Area  $\Delta ABC = \frac{1}{2} \times 4 \times 16$  ✓

$= 32$  units<sup>2</sup> ✓

(b)

$x$	0	$\frac{\pi}{8}$	$\frac{\pi}{4}$
$y$	0	0.17157	1

$V = \pi \int_0^{\frac{\pi}{4}} x \tan^2 x \, dx$

$= \pi \left[ \frac{\pi}{24} (0 + 4 \times 0.17157 + 1) \right]$  ✓✓

$= 0.693$  (u<sup>3</sup>) ✓✓ (subtract 1 mark for incorrect rounding or if they have not attempted to round off)

QUESTION 5

a) (i)  $75^\circ = 75 \times \frac{\pi}{180}$

$= \frac{5\pi}{12}$  ✓

$\therefore l = \frac{5\pi}{12} \times 4.8$

$= 2\pi$  cm ✓

(subtract 1 mark if either answer has been approximated: i.e. 6.283... or 15.0796...)

(ii) Area of sector =  $\frac{1}{2} \cdot (4.8)^2 \cdot \frac{5\pi}{12}$  ✓

$= \frac{24\pi}{5}$  (or  $4.8\pi$ ) cm<sup>2</sup> ✓

(iii) Area of segment = Area of sector - Area of  $\Delta$   
 $= \frac{24\pi}{5} - \frac{1}{2} \cdot (4.8)^2 \cdot \sin \frac{5\pi}{12}$  ✓

$= 3.952$  cm<sup>2</sup> ✓

(b) (i)  $\tan x = -3$


Related angle = 1.249 ✓

$\therefore x = \pi - 1.249$  or  $2\pi - 1.249$

$= 1.893$  or  $5.034$  ✓

(need both for this mark. No marks for degree equivalent)

(ii) Since there are 2 solutions from  $0 \leq x \leq 2\pi$ , there are 4 solutions in the range  $-2\pi \leq x \leq 2\pi$  ✓



✓✓ (Do not penalise if they draw graphs outside domain)

QUESTION EIGHT (Start a new answer booklet)

(a) Kerry deposits \$1500 into a superannuation fund on January 1st 2001. He makes further deposits of \$1500 on the first of each month up to and including December 1st 2010. The fund pays compound interest at a monthly rate of 0.75%. In each of the following questions give your answer to the nearest dollar.

Marks

1

(i) How much is in the fund on January 31st 2001?

1

(ii) How much is the first \$1500 deposit worth on December 31st 2010?


3

(iii) Form a geometric series and hence determine the total amount in the fund on December 31st 2010.

2

(iv) If each deposit was increased to \$1600, what difference does it make to the total amount in the fund on December 31st 2010?

(b)



In the diagram above, a straight line has been drawn which intersects with  $y = e^x$  at the points  $C(1, e)$  and  $D(-1, \frac{1}{e})$ . The point  $A$  has coordinates  $(-1, 0)$  and  $B$  has coordinates  $(1, 0)$ . The area between the curves has been shaded.

2

(i) Show that the area of the trapezium  $ABCD$  is given by  $\frac{e^2 + 1}{e}$ .

3

(ii) Hence, or otherwise, find the exact area between the curves.

**QUESTION NINE** (Start a new answer booklet)

Marks

2

- (a) The value \$V\$ of a car is given by the formula  $V = Ce^{-kt}$ , where  $C$  and  $k$  are constants and  $t$  is the time measured in years. Michael bought a car on June 30th 2001 which cost \$65 000 and which was worth \$55 000 after one year.

1

- (i) Evaluate the constants  $C$  and  $k$ .
- (ii) Find the value of the car after 5 years. Give your answer correct to the nearest dollar.

2

- (iii) In which year will the value of the car fall below half its cost price for the first time?

- (b) Concrete is pumped from a truck into a building foundation. The rate  $R$  m<sup>3</sup>/hour at which the concrete is flowing is given by the expression  $R = 9t^2 - t^4$  for  $0 \leq t \leq 3$ , where  $t$  is the time measured in hours after the concrete begins to flow.

1

- (i) Find the rate of flow at time  $t = 2$ .

1

- (ii) Explain why  $t$  is restricted to  $0 \leq t \leq 3$ .

3


- (iii) Find the maximum flow rate of concrete.

2

- (iv) When the concrete begins to flow, the foundation has 1000 m<sup>3</sup> already in place. Find an expression for the amount of concrete in the foundation at time  $t$ .

**QUESTION 10 IS ON THE NEXT PAGE.**

**QUESTION 4**



(i)  $\int_{-1}^3 f(x) + 2 \, dx = \frac{15}{2} + 8$   
 $= \frac{31}{2}$

b)  $x = 2t^3 - t^2$

(i)  $\dot{x} = 6t^2 - 2t$

At rest when  $\dot{x} = 0$

$\therefore 2t(3t - 1) = 0$

$\therefore t = 0$  or  $\frac{1}{3}$

(ii) At  $t = 0$ ,  $x = 0$

$t = \frac{1}{3}$ ,  $x = 2 \cdot \frac{1}{27} - \frac{1}{9}$

$= -\frac{1}{27}$

$\therefore$  Distance travelled is  $\frac{1}{27}$  m.

(c) (i)  $\angle ADB = 40 + \theta$  (exterior angle of  $\Delta$ )

(ii)  $\angle DBE = 40$  (isosceles  $\Delta$ )

$\angle DAE + 40 + \theta + 40 = 180$  (angle sum)

$\therefore \angle DAE = 100 - \theta$

(iii)  $\angle ADE = \frac{1}{2} (180 - (100 - \theta))$  (base  $\angle$ s, isosceles  $\Delta ADE$ )

$= 40 + \frac{\theta}{2}$

$\therefore \angle EDB = (40 + \theta) - (40 + \frac{\theta}{2})$

$= \frac{\theta}{2}$



QUESTION 3

(a) (i)  $BC^2 = 9.81^2 + 15.24^2 - 2 \times 9.81 \times 15.24 \times \cos 31^\circ 20'$  ✓  
 $= 73.089... \text{ m}$   
 $\therefore BC \doteq 8.55 \text{ m}$  (nearest cm) ✓ (Penalise incorrect rounding but ignore no rounding to 2 d.p.)

(ii) Area  $\Delta ABC = \frac{1}{2} ab \sin C$   
 $= \frac{1}{2} \cdot 9.81 \cdot 15.24 \cdot \sin 31^\circ 20'$  ✓  
 $= 38.87... \text{ m}^2$   
 $\doteq 39 \text{ m}^2$  ✓

(b) (i)  $r = -\frac{1}{3}$ . Since  $|r| < 1$ ,  $S_\infty$  exists ✓  
 (ii)  $S_\infty = \frac{a}{1-r}$   
 $= \frac{1}{\frac{4}{3}}$  ✓  
 $= \frac{3}{4}$  ✓


(c)  $\frac{dy}{dx} = \frac{1}{x}$   
 At  $x = e$ ,  $\frac{dy}{dx} = \frac{1}{e}$  } ✓ (for either or both)

Equation of tangent is:  
 $y - 1 = \frac{1}{e}(x - e)$  ✓  
 $y - 1 = \frac{x}{e} - 1$

$\therefore x = ey$  ✓  
 (d) (i)  $\alpha + \beta = -8$  ✓  
 (ii)  $\alpha\beta = 11$  ✓  
 (iii)  $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$   
 $= (-8)^2 - 2 \times 11$  ✓  
 $= 42$  ✓

QUESTION TEN (Start a new answer booklet)

(a)



In the diagram above,  $P$  is a point on the circle with diameter  $AB = l$ . The point  $Q$  is on the diameter such that  $AP = AQ$ . Let  $\angle PAQ = \theta$  and let  $S$  be the area of  $\Delta PAQ$ .

Marks

- 3 (i) Show that  $S = \frac{l^2}{2} \cos^2 \theta \sin \theta$ .
- 3 (ii) Find the maximum area of  $\Delta APQ$  as  $P$  moves along the circumference of the circle.
- 2 (b) (i) Find  $A$  and  $B$  such that  $\frac{1}{(x-1)x} = \frac{A}{x-1} + \frac{B}{x}$ .
- 2 (ii) Let  $S_n = \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{(n-1)n}$ . Show that  $S_n = 1 - \frac{1}{n}$ .
- 2 (iii) Hence or otherwise evaluate  $\sum_{n=2}^{\infty} \frac{1}{(n-1)n}$ .

QUESTION 1

a)  $144^\circ$  ✓

b)  $\frac{1}{5} \tan 5x + c$  ✓

c)  $5 - k = 7$  ✓  
 $\therefore k = -2$  ✓

d)  $\frac{dy}{dx} = 15x^2 - 2$  ✓✓ (1 each)

e)  $\frac{4}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} = \frac{4(\sqrt{3}+1)}{2}$  ✓  
 $= 2(\sqrt{3}+1)$  ✓

f)  $\sqrt{3} + 1$  ✓✓ (1 each)

g)  $|x-1| = 11$

$x-1 = 11$  or  $x-1 = -11$

$\therefore x = 12$  or  $-10$  ✓✓

Grammar 2001  
SOLUTIONS

QUESTION 2

a) (i)  $\frac{dy}{dx} = 2xe^x + x^2e^x$  ✓✓  
 $= xe^x(2+x)$  (not necessary)

(ii)  $\frac{dy}{dx} = \frac{3}{3x-2}$  ✓✓

(iii)  $\frac{dy}{dx} = 2 \sin x \cdot \cos x$  ✓✓

b)  $\frac{(3x-4)^7}{21}$  ✓✓

c) (i)  $\int_1^2 6x^2 dx = \left[ 2x^3 \right]_1^2$  ✓  
 $= 2 \times 8 - 2 \times 1$  ✓  
 $= 14$

(ii)  $\int_0^{\frac{\pi}{2}} \sin 2x dx = -\frac{1}{2} \left[ \cos 2x \right]_0^{\frac{\pi}{2}}$  ✓  
 $= -\frac{1}{2} [\cos \pi - \cos 0]$   
 $= -\frac{1}{2} [-1 - 1]$  ✓  
 $= 1$  ✓