



St Catherine's
School
Waverley, Sydney

Student Number: _____

Extension I Mathematics
Assessment Task 3
6 June-2007

Time allowed: 55
minutes

General Instructions


- Attempt ALL questions
- Write your NAME or Student NUMBER at the top of this page and on any extra writing paper used
- Answer the questions in the spaces provided in this paper

Questions

Marks

Question 1.

The diagram shows a parabolic drinking cup of height 12 cm and radius 8 cm. The cup is being filled with water at a constant rate of 20 cm^3 per second. The height of the water at time t seconds is h cm and the radius is r cm.



- (i) Using similar triangles, show that $r = \frac{2}{3}h$ (1m)
- (ii) Find the rate at which the height is increasing, when the height of the water level is 10 cm. Leave your answer in terms of π .
(Volume is given by $V = \frac{1}{3}\pi r^2 h$) (3m)

Question 2.

The rate at which a body cools in air is assumed to be proportional to the difference between its temperature T and the constant temperature S of the surrounding air. This can be expressed by the differential equation

$$\frac{dT}{dt} = k(T - S),$$

where t is the time in hours and k is a constant.

- (i) Show that $T = S + Ae^{kt}$, where A is a constant is a solution of the differential equation. (1m)

A heated body cools from 90°C to 60°C in 2 hours. The surrounding air temperature S the body is 25°C .

- (ii) Show that $T = 25 + 65e^{-0.3095t}$ (3m)
- (iii) Find the time taken for the body to have a temperature of 30°C (2m)
- (iv) What is the limiting value of the temperature. (1m)
- (v) Draw a sketch of the equation relating T in terms of t . (2m)

Question 3.

The velocity of a particle travelling in a straight line is given by $v = 5x$, where x is the displacement from the origin.

- (i) Given that initially the particle has a velocity of 5 metres per second, find an expression for x , the displacement of the particle in terms of t , the time. (4m)
- (ii) Find an expression of acceleration a of the particle in terms of t , the time. (1m)

Question 4.

A ladder 5 metres in length is leaning against a wall. It is slipping down the wall at the rate is 0.2 metres per second. Find the rate at which its foot is slipping on the floor when its foot is 3 metres away from the wall.

(4m)

Question 5

The acceleration of a particle in motion is given by


$$\frac{d^2x}{dt^2} = 4 - 2x$$

The particle has a velocity of 4 metres per second at the point $x=1$.

- (i) Show that the velocity of the particle is given by $v^2 = 2(9 - (x - 2)^2)$ (3m)
- (ii) Find the centre, the end points and the amplitude of the motion (3m)
- (iii) Find the period of motion. (1m)

Question 6.

The diagram shows the path of a projectile launched with a velocity of 40 metres per second at an angle of elevation θ to the horizontal from the top of a building 50 metres high. The acceleration due to gravity is 10 metres per second per second.



- (i) Show that

$$x = (40 \cos \theta) t \quad \text{and} \quad y = -5t^2 + (40 \sin \theta) t + 50 \quad (2\text{m})$$

- (ii) The maximum height reached is 90 metres from the ground.
Prove that the angle of projection is 45° (3m)

- (iii) Find only the magnitude of the velocity of the particle when $t = \sqrt{2}$ (2m)

End of Paper.

Question 2

$$\frac{dT}{dt} = k(T-S)$$

(i) $T = S + Ae^{kt}$

$$\frac{dT}{dt} = A \cdot k \cdot e^{kt}$$

$$\frac{dT}{dt} = kAe^{kt}$$

$$= k(T-S)$$

(ii) $t=0, T=90 \quad t=2, T=60$

$$90 = S + Ae^0$$

$$= 25 + A$$

$$\therefore A = 65$$

(1)

$$60 = 25 + 65e^{2k}$$

$$35 = 65e^{2k}$$

$$2k = \frac{\ln \frac{35}{65}}$$

$$k = \frac{1}{2} \ln \frac{35}{65}$$

$$= -0.3095$$

(2)

$$\therefore T = 25 + 65e^{-0.3095t}$$

(3)

(iii) $30 = 25 + 65e^{-0.3095t}$

$$\frac{5}{65} = e^{-0.3095t}$$

$$t = \frac{-1}{0.3095} \ln \frac{5}{65}$$

$$= 8.29 \text{ hrs}$$


(4)

$$k \rightarrow \infty; e^{-0.3095t} \rightarrow 0$$

$$\therefore T \rightarrow 25$$

(1)

(v)



(2)

Q4:

$$\frac{8}{12} = \frac{r}{h}$$

$$\therefore r = \frac{8h}{12}$$

$$= \frac{2}{3}h$$

$$V = \frac{1}{3} \cdot \pi \cdot \left(\frac{2}{3}h\right)^2 \cdot h$$

$$= \frac{1}{3} \pi \cdot \frac{4}{9} h^3$$

$$= \frac{4\pi}{27} h^3$$

(1)

$$\frac{dv}{dt} = \frac{4\pi}{27} \cdot 3h^2 \cdot \frac{dh}{dt}$$

(2)

$$20 = \frac{4\pi}{27} \cdot 3 \times 10^2 \cdot \frac{dh}{dt}$$

(3)

$$\frac{dx}{dt} = \frac{20 \times 27^8}{8 \times 10^2 \times 4 \pi}$$

$$= \frac{9}{20 \pi}$$

$\frac{1}{2}$

Q.3

$$u = 5x$$

$$\frac{dr}{dx} = \frac{1}{5x}$$

$$r = \frac{1}{5} \ln x + C$$

$$t=0, u=5^0 \therefore x=1$$

$$r = \frac{1}{5} \ln 1 + C$$

$$\therefore C = 0$$

$$\therefore r = \frac{1}{5} \ln x$$

$$\ln x = 5r$$

$$x = e^{5r}$$


$$\frac{dx}{dr} = 5e^{5r}$$

$$\frac{d^2x}{dr^2} = 25e^{5r}$$

$$\frac{d^2x}{dr^2}$$

$\frac{1}{2}$
 $\frac{1}{2}$

Q.4



$$\frac{dy}{dt} = -0.2$$

$$\frac{dx}{dt} = 0.2$$

$$x = 3$$

$$s^2 = x^2 + y^2$$

$$y = \sqrt{s^2 - 3^2}$$

$$0 = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$0 = 3 \times \frac{dx}{dt} + 4 \times 0.2$$

$$\frac{dx}{dt} = \frac{0.8}{3}$$

$$= 0.26 = \frac{4}{15}$$

[minus sign - deduct $\frac{1}{2}$]

Q.5

$$\frac{d^2x}{dt^2} = 4 - 2x$$

$$= -2(x - 2)$$

$$\frac{d}{dx} \left(\frac{1}{2} v^2 \right) = 4 - 2x$$

$$\frac{1}{2} v^2 = 4x - x^2 + C$$

$$\frac{1}{2} \times 4^2 = 4 - 1^2 + C$$

$$C = 5$$

$$\frac{1}{2}v^2 = 4x - x^2 + 5$$

$$v^2 = 8x - 2x^2 + 10$$

$$= 2(5 - x^2 + 4x)$$

$$= 2(9 - (x^2 - 4x + 4)) \quad (1)$$

$$= 2(9 - (x-2)^2)$$

(1) Centre is at $x=2$; for $\frac{d^2x}{dt^2} = -2(x-2)$


at end pts ; $v=0$

$$9 - (x-2)^2 = 0$$

$$x-2 = \pm 3$$

$$x = 5, -1$$

amplitude is 3



(14) Period = $\frac{2\pi}{\sqrt{2}} = \sqrt{2}\pi$ per sec. (1)

$$T = \frac{2\pi}{\omega} \left(\frac{1}{2}\pi\right)$$

Question 6

$$x'' = 0 \quad ; \quad y = -10$$

$a = \text{const}$

$$y = -10t + C$$

$t=0$
 $y=40$

$$y = -10t + 40 \sin \theta$$

$$y = -5t^2 + 40 \sin \theta t + C$$

$t=0$
 $y=50$

$\therefore C=0$

$$y = -5t^2 + 40 \sin \theta t + m$$

$t=0$

$$x = 40 \cos \theta$$

$$x = 40 \cos \theta$$

$$x = 40 \cos \theta + C$$

$x=0$ what $t=0$ $\therefore C=0$

$$\therefore x = (40 \cos \theta) t$$

At max v ; $y' = 0$ (1/2)

$$-10t + 40 \sin \theta = 0$$

$$t = 4 \sin \theta \quad (2)$$

Sub in y ;

$$90 = -5(4 \sin \theta)^2 + (40 \sin \theta)(4 \sin \theta) + 50 \quad (1)$$

$$40 = -80 \sin^2 \theta + 160 \sin^2 \theta$$

$$= 80 \sin^2 \theta$$

$$\sin^2 \theta = \frac{1}{2}$$

$$\sin \theta = \pm \frac{1}{\sqrt{2}}$$

$$\theta = 45^\circ \quad (\text{positive angle or acute angle})$$

(11)

$$x = (40 \cos \theta) t$$

$$= \frac{40}{\sqrt{2}} = \frac{20\sqrt{2}}{1}$$

$$y = -10t^2 + 40 \sin \theta$$

$$= -10(\sqrt{2})^2 + \frac{40}{\sqrt{2}} = 20\sqrt{2} - 10\sqrt{2}$$

$$= 20\sqrt{2} - 10\sqrt{2}$$

$$u^2 = x^2 + y^2 = 800 + 200$$

$$u = \sqrt{(20\sqrt{2})^2 + (10\sqrt{2})^2} = 30\sqrt{2} = 10\sqrt{18}$$