

St. Catherine's School
Waverley

February 2008
HSC ASSESSMENT TASK
EXAMINATION

Extension I Mathematics

Time allowed: 55 minutes

INSTRUCTIONS

- There are 2 sections of value 17, 21
- Marks for each part of a question are indicated
- All questions should be attempted on the separate paper provided
- All necessary working should be shown
- Start each question on a new page
- Approved scientific calculators and drawing templates may be used badly arranged work.

Section A

17 Marks

- Find the remainder when the polynomial $P(x) = x^3 - 4x$ is divided by $(x + 3)$ [2]
- For the polynomial equation $x^3 - 2x^2 + x + 5 = 0$
 - Show that one root of this equation lies in the interval $-2 < x < -1$. [1]
 - Use Newton's method once to find an approximation of this root starting with $x = -1.5$ as a first approximation. [2]
- When the polynomial $P(x) = x^4 - 3x^3 + ax^2 + bx - 6$ is divided by $(x + 1)$ the remainder is 8. [3]
If $(x - 3)$ is a factor of $P(x)$, find the values of a and b .
- For the polynomial $P(x) = x^4 - 3x^2 - 2x$
 - Show that $P(x)$ has a double zero at $x = -1$. [2]
 - Write $P(x)$ as the product of its linear factors. [2]
- Show that $(x - 1)(x - 2)$ is a factor of [2]
 $P(x) = x^n(2^m - 1) + x^m(1 - 2^n) + (2^n - 2^m)$
where m and n are positive integers
- The polynomial $P(x) = 8x^3 - 20x^2 + 6x + 9$ has two equal roots. [3]
Find all the roots of $P(x) = 0$

Section B – Start a new page

21 Marks

1. Solve the equation $2\sin^2\theta = \sin 2\theta$ for $0 \leq \theta \leq 360^\circ$ [3]

2. (i) Write $\cos x - \sqrt{3}\sin x$ in the form $R\cos(x + \alpha)$ [2]

(ii) Hence or otherwise solve the trigonometric equation [2]


$$\cos x - \sqrt{3}\sin x = \sqrt{3} \quad \text{for } 0^\circ \leq x \leq 360^\circ$$

3. If $\tan \frac{\theta}{2} = t$, prove $\frac{1 + \sin\theta - \cos\theta}{1 + \sin\theta + \cos\theta} = t$ [3]

4. A surveyor who is y metres south of a tower sees the top of it with an angle of elevation 8° . A second surveyor is x metres east of the tower. From his position the angle of elevation is 10° to the top of the tower. The two surveyors are 940m apart.

(i) Show that $y = h \cot 8^\circ$ [1]

(ii) Find the height of the tower to the nearest metre. [2]



$$\tan 8^\circ = \frac{y}{h}$$

$$y = \tan 8^\circ h$$

$$x = \tan 10^\circ h$$

$$\tan^2 8^\circ h^2 + \tan^2 10^\circ h^2 = 940^2$$

$$h^2 ($$

5. Prove that $\frac{2 \tan A}{1 + \tan^2 A} = \sin 2A$ [2]

6. Find the general solution of $2\cos^2 x = \sqrt{3}$ [3]

7. Prove that $\frac{\tan\theta + 1}{\sec\theta} - \frac{\cot\theta + 1}{\operatorname{cosec}\theta}$ is independent of θ [3]

End of Task

Course:

Marking Scheme for Task:

Academic Year: 2007-8

Solutions	Marks	Comments
<p>Q3 $P(x) = x^4 - 3x^3 + ax^2 + bx + 6$ $P(1) = 1 + 3 + a - b - 6 = 8$ $\therefore a - b = 10$ — (1) $P(3) = 81 - 81 + 9a + 3b - 6 = 0$ $\therefore 9a + 3b = 6$ — (2) Now (1) \times 3 + (2) $12a = 36$ $\therefore a = 3$ and $b = -7$</p>	1 1 1	
<p>Q4 (i) $P(x) = x^4 - 3x^2 - 2x$ $P(1) = 1 - 3 + 2 = 0$ $P'(x) = 4x^3 - 6x - 2$ $P'(1) = -4 + 6 - 2 = 0$ $\therefore x = -1$ is a double root of $P(x)$</p>	1 1	
<p>(ii) $P(x) = (x+1)^2 Q(x)$ $\therefore x^4 - 3x^2 - 2x = (x^2 + 2x + 1)(x^2 - 2x)$ $= x(x-2)(x+1)^2$</p>	2	
<p>OR clearly $P(x) = x(x^3 - 3x - 2)$ $= x(x+1)^2(x-2)$ clearly $P(x) = x(x+1)^2(x-2)$ Since product of roots of $x^3 - 3x - 2$ is 2.</p>	1 1	

Course:

Marking Scheme for Task:

Academic Year: 2007-8


Solutions	Marks	Comments
<p>Q5 $P(x) = 1^n(2^m - 1) + 1^m(1 - 2^n) + 2^n - 2^m$ $= 2^m - 1 + 1 - 2^n + 2^n - 2^m$ $= 0$ $\therefore (x-1)$ is a factor. $P(2) = 2^n(2^m - 1) + 2^m(1 - 2^n) + (2^n - 2^m)$ $= 2^{n+m} - 2^n + 2^m - 2^{m+n} + 2^n - 2^m$ $= 0$ $\therefore (x-2)$ is a factor $\therefore (x-1)(x-2)$ is a factor of $P(x)$.</p>	1 1	
<p>Q6 $P(x) = 8x^3 - 20x^2 + 6x + 9$ let the roots be α, α, β \therefore sum of roots $= 2\alpha + \beta = \frac{5}{2}$ — (1) sum of roots $2\alpha = \alpha^2 + 2\alpha\beta = \frac{3}{4}$ — (2) product of roots $= \alpha^2\beta = -\frac{9}{8}$ — (3) from (1) $\beta = \frac{5}{2} - 2\alpha$ — (4) Sub in (2) $\alpha^2 + 2\alpha(\frac{5}{2} - 2\alpha) = \frac{3}{4}$ $\alpha^2 + 5\alpha - 4\alpha^2 = \frac{3}{4}$ $-3\alpha^2 + 5\alpha = \frac{3}{4}$ $-12\alpha^2 + 20\alpha - 3 = 0$ $12\alpha^2 - 20\alpha + 3 = 0$ $(6\alpha - 1)(2\alpha - 3) = 0$ $\therefore \alpha = \frac{1}{6}, \frac{3}{2}$ now $P(\frac{3}{2}) = 0 \therefore \frac{3}{2}$ is the double zero. Sub in (4) $\beta = \frac{5}{2} - 3 = -\frac{1}{2}$ \therefore roots are $x = \frac{3}{2}, \frac{3}{2}, -\frac{1}{2}$</p>	1 1 1 1 1 1 1 1	

Course:

Page no. of

Academic Year: 2007-8

Marking Scheme for Task:

Solutions	Marks	Comments
<u>Section B</u>		
<p>Q1 $2 \sin^2 \theta = \sin 2\theta$ $2 \sin^2 \theta - \sin 2\theta = 0$ $2 \sin^2 \theta - 2 \sin \theta \cos \theta = 0$ $2 \sin \theta (\sin \theta - \cos \theta) = 0$ $\therefore \sin \theta = 0$ or $\sin \theta - \cos \theta = 0$ $\theta = 0^\circ, 180^\circ, 360^\circ$ or $0, \pi, 2\pi$ $\sin \theta = \cos \theta$ $\tan \theta = 1$ ($\cos \theta \neq 0$) $\theta = 45^\circ, 225^\circ$ or $\frac{\pi}{4}, \frac{5\pi}{4}$ $\therefore \theta = 0^\circ, 45^\circ, 180^\circ, 225^\circ, 360^\circ$</p>	1	
<p>Q2 (i) $\cos x - \sqrt{3} \sin x$ $(\cos x \cos \alpha - \sin x \sin \alpha)$ $\therefore \cos x - \sqrt{3} \sin x = 2 \cos(x + 60^\circ)$</p>  <p>$\tan \alpha = \sqrt{3}$ $\alpha = 60^\circ$</p>	2	r=2 (mark) d=60 (mark)
<p>(ii) $\cos x - \sqrt{3} \sin x = \sqrt{3}$ $\therefore 2 \cos(x + 60^\circ) = \sqrt{3}$ $\cos(x + 60^\circ) = \frac{\sqrt{3}}{2}$ $x + 60^\circ = \cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$ $x + 60^\circ = 30^\circ, 330^\circ, 390^\circ, \dots$ $\therefore x = -30^\circ, 270^\circ, 330^\circ, \dots$ $\therefore x = 270^\circ, 330^\circ$</p>	1	

Course:

Page no. of

Academic Year: 2007-8

Marking Scheme for Task:

Solutions	Marks	Comments
<p>Q3 $\frac{1 + \sin \theta - \cos \theta}{1 + \sin \theta + \cos \theta} = \frac{1 + \frac{2t}{1+t^2} - \frac{1-t^2}{1+t^2}}{1 + \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2}}$ $= \frac{1+t^2+2t-1+t^2}{1+t^2+2t+1-t^2}$ $= \frac{2t^2+2t}{2+2t}$ $= \frac{2t(t+1)}{2(t+1)}$ $= t$</p>	1	
<p>Q4 (i) $\tan 8^\circ = \frac{h}{y}$ $\therefore y = \frac{h}{\tan 8^\circ}$ $= h \cdot \frac{1}{\tan 8^\circ}$ $= h \cot 8^\circ$</p> <p>(ii) now similarly $x = h \cot 10^\circ$ and by Pythagoras' theorem $(h \cot 8^\circ)^2 + (h \cot 10^\circ)^2 = 940^2$ $h^2 \cot^2 8^\circ + h^2 \cot^2 10^\circ = 940^2$ $\therefore h^2 = \frac{940^2}{\cot^2 8^\circ + \cot^2 10^\circ}$ $h^2 = 10672.60126 \dots$ $\therefore h = 103 \text{ m (nearest m)}$</p>	1	

