

# St Catherine's School

Year: 12  
Subject: Extension II Mathematics  
Time Allowed: 55 minutes  
Assessment task 3  
16/6/2004  
Task weight 15%

Exam number: 1416 9652

## Directions to candidates:

- All questions are to be attempted..
- All necessary **working** must be shown in every question.
- Full marks may not be awarded for careless or badly arranged work.
- Approved calculators may be used.
- Hand in your work in **1 bundle**:

Attach the question paper to your answer sheets

21 + 15 = 36  
excellent

100%

Q.1 Integrate the following:

(a)  $\int \frac{2x+5}{x+3} dx$  (2m)

(b)  $\int \frac{2x+3}{x^2+2x+5} dx$  (4m)


(c)  $\int_0^2 \frac{x^2}{\sqrt{4-x^2}} dx$  (4m)

(d)  $\int \sin^4 x \cos^3 x dx$  (3m)

(e) Use integration by parts to find  $\int e^{2x} \cos x dx$  (4m)

(f) If  $u_n = \int \sin^n x dx$ , show that  $u_n = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} u_{n-2}$  (4m)

Q.2 The shaded area shows the area bounded by the curve  $y = x(x-2)$  and the lines  $x=1$  and  $y=8$ . This area is rotated about the line  $x=1$ .



The coordinates of the point of intersection in the first quadrant between the line  $y=8$  and the given curve is  $(4,8)$ .

Use the method of cylindrical shells to find the volume generated. (5m)

Q.3 The area bounded by the curve  $y = x(4-x)$  and the x axis is rotated around the y axis. Take a slice perpendicular to the axis of rotation and use the slice method to find the volume generated.

(5m)

Please turn over for Q.4.

Question 1.

(a)  $\int \frac{2x+5}{x+3} dx.$

$$= \int 2 - \frac{1}{x+3} dx.$$

$$= 2x - \ln|x+3| + c.$$

$$\begin{array}{r}
 2 \\
 x+3 \overline{) 2x+5} \\
 \underline{2x+6} \\
 -1
 \end{array}$$

(b)  $\int \frac{2x+3}{x^2+2x+5} dx.$

$$= \int \frac{2x+3}{(x+1)^2+4} dx.$$

$$= \int \frac{2x}{(x+1)^2+4} dx.$$

$$= \int \frac{2x+2+1}{x^2+2x+5} dx.$$

$$= \int \frac{2x+2}{x^2+2x+5} + \frac{1}{x^2+2x+5} dx.$$

$$= \ln|x^2+2x+5| + \int \frac{1}{(x+1)^2+4} dx.$$

$$= \ln|x^2+2x+5| + \frac{1}{2} \tan^{-1}\left(\frac{x+1}{2}\right) + c.$$

(c)  $\int_0^2 \frac{x^2}{\sqrt{4-x^2}} dx.$

$$= \int_0^{\frac{\pi}{2}} \frac{4\sin^2\theta}{\sqrt{4-4\sin^2\theta}} \cdot 2\cos\theta d\theta.$$

$$= \int_0^{\frac{\pi}{2}} \frac{4\sin^2\theta}{\sqrt{4(1-\sin^2\theta)}} \cdot 2\cos\theta d\theta.$$

$$= \int_0^{\frac{\pi}{2}} \frac{4\sin^2\theta}{2\cos\theta} \cdot 2\cos\theta d\theta.$$

$$= 4 \int_0^{\frac{\pi}{2}} \sin^2\theta d\theta.$$

$$= 2 \int_0^{\frac{\pi}{2}} 1 - \cos 2\theta d\theta.$$

$$= 2 \left[ \theta - \frac{1}{2} \sin 2\theta \right]_0^{\frac{\pi}{2}}.$$

$$= 2 \left( \frac{\pi}{2} \right)$$

$$= \pi$$

let  $x = 2\sin\theta.$

$$dx = 2\cos\theta d\theta.$$

$$x = 2, \quad \theta = \frac{\pi}{2}.$$

$$x = 0, \quad \theta = 0.$$

$$(d) \int \sin^4 \theta \cos^3 \theta d\theta.$$

$$= \int \sin^4 \theta (1 - \sin^2 \theta) \cos \theta d\theta.$$

$$= \int u^4 (1 - u^2) du.$$

$$u = \sin \theta$$

$$du = \cos \theta d\theta.$$

~~to find~~

$$= \int u^4 - u^6 du.$$

$$= \frac{1}{5} u^5 - \frac{1}{7} u^7 + C.$$

$$= \frac{1}{5} \sin^5 \theta - \frac{1}{7} \sin^7 \theta + C.$$

$$(e) \int e^{2x} \cos x dx = e^{2x} \sin x - \int 2e^{2x} \sin x.$$

$$= e^{2x} \sin x - 2 \int e^{2x} \sin x.$$

$$= e^{2x} \sin x - 2 \left[ -e^{2x} \cos x + \int 2e^{2x} \cos x dx \right]$$

$$= e^{2x} \sin x + 2e^{2x} \cos x - 4 \int e^{2x} \cos x dx.$$

$$5 \int e^{2x} \cos x dx = e^{2x} \sin x + 2e^{2x} \cos x.$$

$$\int e^{2x} \cos x dx = \frac{1}{5} (e^{2x} \sin x + 2e^{2x} \cos x) + C.$$

det :

$$u = e^{2x}$$

$$v' = \cos x.$$

$$u' = 2e^{2x}$$

$$v = \sin x.$$

$$u = e^{2x}$$

$$v' = \sin x$$

$$u' = 2e^{2x}$$

$$v = -\cos x.$$

$$(f) U_n = \int \sin^n x dx.$$

$$= \int \sin^{n-1} x \sin x dx.$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos^2 x dx.$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x (1 - \sin^2 x) dx.$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x - \sin^n x dx.$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x dx - (n-1) \int \sin^n x dx.$$

$$= -\sin^{n-1} x \cos x + (n-1) U_{n-2} - (n-1) U_n.$$

$$U_n (1 + n - 1) = -\sin^{n-1} x \cos x + (n-1) U_{n-2}.$$

$$U_n = \frac{1}{n} \sin^{n-1} x \cos x + \left( \frac{n-1}{n} \right) U_{n-2}.$$

$$u = \sin^{n-1} x, \quad v' = \sin x$$

$$u' = (n-1) \sin^{n-2} x \cos x, \quad v = -\cos x.$$

$$u' = (n-1) \sin^{n-2} x \cos x.$$

Question 2.

$$\Delta V = 2\pi r h \Delta x.$$

$$= 2\pi (x-1)(8-y) \Delta x.$$

~~$$= 2\pi (8x - xy - 8 + y) \Delta x.$$~~

$$= 2\pi (x-1)(8 - (x^2 - 2x)) \Delta x.$$

$$= 2\pi (x-1)(8 + 2x - x^2) \Delta x.$$

$$= 2\pi (8x + 2x^2 - x^3 - 8 - 2x + x^2) \Delta x.$$


$$= 2\pi (3x^2 - x^3 + 6x - 8) \Delta x.$$

$$V = 2\pi \int_1^4 (-x^3 + 3x^2 + 6x - 8) dx.$$


$$= 2\pi \left[ -\frac{1}{4}x^4 + x^3 + 3x^2 - 8x \right]_1^4.$$

$$= 2\pi \left[ 16 + 4\frac{1}{4} \right]$$

$$= 40\frac{1}{2} \pi \text{ units}^3.$$



## Question 3



$$y = x(4 - x)$$

$$y = 4x - x^2$$

$$x^2 - 4x + y = 0$$

$$x_1 + x_2 = -\left(\frac{-4}{1}\right) = 4$$

$$x_1 x_2 = y$$

$$(x_2 - x_1)^2 = (x_2 + x_1)^2 - 4x_1 x_2$$

$$= 16 - 4y$$


$$x_2 - x_1 = 2\sqrt{4 - y}$$

$$\begin{aligned} \Delta V &= \pi (x_2^2 - x_1^2) \Delta y \\ &= \pi (x_2 + x_1)(x_2 - x_1) \Delta y \\ &= \pi (4)(2\sqrt{4 - y}) \Delta y \end{aligned}$$


$$\begin{aligned} V &= 8\pi \int_0^4 \sqrt{4 - y} \, dy \\ &= 8\pi \int_0^4 (4 - y)^{\frac{1}{2}} \, dy \\ &= 8\pi \left[ -\frac{2}{3}(4 - y)^{\frac{3}{2}} \right]_0^4 \\ &= 8\pi \left[ \frac{16}{3} \right] \end{aligned}$$

$$= \frac{128}{3} \pi \text{ units}^3$$

Question 4.



Typical slice =



$$(a) \quad \Delta V = (\text{Area of triangle}) \Delta x \\ = \sqrt{3} y^2 \Delta x$$

$$\text{Area of triangle} = \frac{1}{2} ab \sin C$$

$$= \frac{1}{2} (2y)(2y) \sin 60^\circ$$

$$= \frac{1}{2} (4y^2) \frac{\sqrt{3}}{2}$$

$$= \sqrt{3} y^2$$

$$(b) \quad V = \sqrt{3} \int_{-1}^1 y^2 dx \\ = 2\sqrt{3} \int_0^1 x^2 dx \\ = \sqrt{3} \left[ x - \frac{1}{3} x^3 \right]_{-1}^1 \\ = \frac{4\sqrt{3}}{3} \sqrt{3} \left[ \frac{4}{3} \right]$$

$$= \frac{4\sqrt{3}}{3} \text{ units}^3$$