



# St Catherine's School

Year: 12

Subject: Mathematics

Time allowed: 55 minutes

Date: 17<sup>th</sup> February 2006

Name: 16358959  
Natalie Giam

**Directions to candidates:**

- All questions are to be attempted.
- Marks may be deducted for careless or badly arranged work.
- Answer all questions.
- All necessary **working** must be shown in every question.
- Start a new page for every question clearly labelling the questions.

**GOOD LUCK ☺**

**Teacher's Use Only**

Question 1	/11
Question 2	/7
Question 3	/12
Question 4	/8
<b>TOTAL</b>	<b>/38</b>

**Question 1***(Start a new page)***11 Marks**

- (a) Find the equation of the tangent to the curve  $y = x^3 - x$  at the point where  $x = 2$ . 3
- (b) Differentiate with respect to  $x$ :
- (i)  $f(x) = \sqrt{25 - x^2}$  3
- (ii)  $y = \frac{(3x - 2)}{(2 - x)}$  3
- (iii)  $y = \frac{1}{\sqrt[3]{x^2}}$  2

**Question 2****Locus***(Start a new page)***7 Marks**

- (a)  $P(x, y)$  is a point which moves so that its distance from  $A(2, 1)$  is always equal to its distance from the line  $y = -1$ .
- (i) Show that the equation of the locus of  $P$  is:  $(x - 2)^2 = 4y$ . 2
- (ii) Sketch the locus of  $P$  clearly labelling the vertex, focus and the directrix. 2
- (b) Find the equation of the parabola in which the focus is  $(1, 5)$  and the equation of the directrix is  $y = -1$ . 3

**Question 3***(Start a new page)***12 Marks**

- (a) Consider the function  $f(x) = (x - 5)^2(x + 1)$
- (i) Show that  $f'(x) = 3(x^2 - 6x + 5)$   $3x^2 - 18x + 15$  2
- (ii) Find the co-ordinates of the stationary points of the curve  $y = f(x)$  and determine their nature. 3
- (iii) Find the  $x$ - and  $y$ -intercepts 2
- (iv) Sketch the graph of the curve  $y = f(x)$  showing all intercepts, and stationary points for the domain  $-3 \leq x \leq 7$ . 3
- (v) What is the minimum value of the function in this domain. 1
- (vi) For what values of  $x$ , in the given domain, is the function increasing? 1

**Question 4***(Start a new page)***8 Marks**

- (a) In an arithmetic sequence, the fourth term is 12 and the fourteenth term is 62.
- (i) Find the first term and the common difference. 3
- (ii) Calculate the sum of the first 50 terms of this sequence. 2
- (b) Find the number of terms in the geometric series 4, 12, 36,.... whose sum is 1456. 3

 $108 \quad 324 \quad 972$ **END OF TASK**

✓ means 1 mark / ✗ means 1/2 mark

Question 1:

(a)  $y = x^3 - x$

$y' = 3x^2 - 1$  ✓

when  $x = 2$ ,  $y' = 3(2)^2 - 1$

$y' = 11$  ✓

ie gradient of the tangent is 11

Equation of the tangent is

$y - y_1 = m(x - x_1)$

$y - 6 = 11(x - 2)$

$y - 6 = 11x - 22$

$11x - y - 16 = 0$  ✓

and when  $x = 2$

$y = 2^3 - 2$

$= 8 - 2$

$y = 6$  ✓

(b) i)  $f(x) = \sqrt{25 - x^2}$

$f(x) = (25 - x^2)^{1/2}$  ✓

$f'(x) = \frac{1}{2}(25 - x^2)^{-1/2} \times (-2x)$

$= -x(25 - x^2)^{-1/2}$  ✓

$= \frac{-x}{(25 - x^2)^{1/2}}$

$= \frac{-x}{\sqrt{25 - x^2}}$  ✓

ii)  $y = \frac{3x - 2}{2 - x}$

let  $u = 3x - 2$

$v = 2 - x$

$u' = 3$

$v' = -1$

$y' = \frac{u'v - v'u}{v^2}$  ✓

$= \frac{3(2 - x) - (-1)(3x - 2)}{(2 - x)^2}$  ✓

$= \frac{6 - 3x + 3x - 2}{(2 - x)^2}$

$= \frac{4}{(2 - x)^2}$  ✓

(iii)  $y = \frac{1}{\sqrt[3]{x^2}}$

$y = x^{-2/3}$  ✓

$y' = -\frac{2}{3}x^{-5/3}$  ✓


$= -\frac{2}{3x^{5/3}}$

$= -\frac{2}{3x^{5/3}}$

$= -\frac{2}{3\sqrt{x^5}}$  ✓

Question 2:

(a)



$PA = PB$

$\sqrt{(x - 2)^2 + (y - 1)^2} = (y + 1)$  ✓

$(x - 2)^2 + (y - 1)^2 = (y + 1)^2$

$x^2 - 4x + 4 + y^2 - 2y + 1 = y^2 + 2y + 1$

$x^2 - 4x + 5 - 2y - 2y - 1 = 0$

$x^2 - 4x + 4 - 4y = 0$  ✓

$$x^2 = 4x + 4 = 4y$$

$$(x-2)^2 = 4y$$

$$(ii) (x-2)^2 = 4y$$

$$(x-h)^2 = 4a(y-k)$$



$$(x-2)^2 = 4 \times \frac{1}{4} (y-0)$$

$V(h, k)$

$$\therefore V(2, 0)$$

Focal length is  $\frac{1}{4}$

focus  $(2, \frac{1}{4})$   
and directrix  $y = -\frac{1}{4}$



focal length = 3 ✓

$$(x-h)^2 = 4a(y-k)$$

$$(x-1)^2 = 4 \times 3 (y-2)$$

$$(x-1)^2 = 12(y-2) \quad \checkmark$$

### Question 3

$$(a) f(x) = (x-5)^2(x+1)$$

$$(ii) f'(x) = u'v + v'u \quad \text{let } u = (x-5)^2 \quad \left| \begin{array}{l} v = x+1 \\ u' = 2(x-5) \\ v' = 1 \end{array} \right.$$

$$= 2(x-5)(x+1) + 1(x-5)^2$$

$$= 2(x^2 + x - 5x - 5) + x^2 - 10x + 25$$

$$= 2x^2 - 8x - 10 + x^2 - 10x + 25$$

$$= 3x^2 - 18x + 15$$

$$= 3(x^2 - 6x + 5)$$

(ii) Stat. pts occur at  $f'(x) = 0$

$$\text{ie. } 3(x^2 - 6x + 5) = 0$$

$$x^2 - 6x + 5 = 0$$

$$(x-5)(x-1) = 0$$

$$\therefore x = 5, 1 \quad \checkmark$$

Test for  $x=5$

$x$	$5^-$	$5$	$5^+$
$f(x)$	$< 0$	$0$	$> 0$

↪ min

$(5, 0)$  minimum ✓  
turning pt

when  $x=5$

$$y = (5-5)^2(5+1)$$

$$= (0)(6)$$

$$= 0$$

Test for  $x=1$

$x$	$1^-$	$1$	$1^+$
$f(x)$	$> 0$	$0$	$< 0$

↪ max. ✓  
 $(1, 32)$  maximum  
turning pt

when  $x=1$

$$y = (1-5)^2(1+1)$$

$$= (-4)^2(2)$$

$$= (16)(2)$$

$$= 32$$

iii) The x-intercept (is when  $y=0$ )

$$f(x) = (x-5)^2(x+1)$$

$$0 = (x-5)(x-5)(x+1)$$

$$\therefore x = 5, 5, -1$$

The y-intercept (when  $x=0$ )

$$f(x) = (0-5)^2(0+1)$$

$$= (25)(1)$$


$$= 25$$

(v) The minimum value of the function is  $y = -128$

(vi)  $3 \leq x \leq 1$  and

$$5 \leq x \leq 7$$

(iv)



Question 4:

a) (i)  $T_4 = 12$

and  $T_{14} = 62$

$$T_n = a + (n-1)d$$

$$12 = a + (4-1)d$$

$$62 = a + (13-1)d \quad \text{--- (2) ✓}$$

$$12 = a + 3d \quad \text{--- (1) ✓}$$

$$a + 3d = 12 \quad \text{--- (1)}$$

$$-a + 13d = 62 \quad \text{--- (2)}$$

$$-10d = -50$$

$$d = 5 \quad \text{✓}$$

$$a + 3d = 12$$

$$a + 3(5) = 12$$

$$a + 15 = 12$$

$$a = 12 - 15$$

$$a = -3 \quad \text{✓}$$

$\therefore$  first term is  $-3$   
and the common difference is  $5$

(ii)  $S_n = \frac{n}{2} [2a + (n-1)d]$  ✓

$$S_{50} = \frac{50}{2} [2(-3) + (50-1)5]$$

$$= 25 [-6 + (49 \times 5)]$$

$$= 25 [239]$$

$$= 5975 \quad \text{✓}$$

$$\frac{4(3^n - 1)}{3 - 1} = 145$$

$$\frac{4(3^n - 1)}{2} = 145$$

$$2(3^n - 1) = 145$$

$$3^n - 1 = \frac{145}{2}$$

b)  $S_n = 1456$ ,  $a = 4$ ,  $r = 3$ ,  $n = ?$

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$3^n - 1 = 728$$

$$3^n = 729 \quad \text{✓}$$

$$\therefore n = 6$$