



# Mathematics Extension 1

### General Instructions

- Time allowed – 75 minutes
- Write using blue or black pen
- Board-approved calculators may be used.
- A table of standard integrals is provided.
- All necessary working should be shown in every question.
- Write on one side of the page only.
- Start each question on a new page.

### Total marks – 70

- Attempt Questions 1 – 5
- All questions are of equal value


Question	Mark
Question 1	/14
Question 2	/14
Question 3	/14
Question 4	/14
Question 5	/14
<b>Total</b>	<b>/70</b>

### Question 1 – (14 marks) – Start a new page

Marks

a) Differentiate  $x \tan^{-1} \frac{x}{2}$

3



In the triangle  $ABC$ ,  $XY = 8\text{cm}$ ,  $BC = 14\text{cm}$ ,  $AC = 18\text{cm}$  and  $XY \parallel BC$ .

(i) Prove that  $\triangle AXY$  is similar to  $\triangle ABC$

3

(ii) Find the length of  $AY$  giving reasons.

3

c) For the function  $y = \sin^{-1}\left(\frac{x}{2}\right)$ :

(i) State the domain and range

2


(ii) Sketch the graph of the function

3

**Question 2** – (14 marks) – Start a new page

Marks

a)




$ABCD$  is a rectangle.  $E$  is a point on  $DC$  such that  $AE$  bisects angle  $DEB$  and  $EF$  bisects angle  $BEC$ . Prove that angle  $AEF = 90^\circ$

b) Evaluate  $\int_0^1 \frac{1}{\sqrt{4-3x^2}} dx$

c) Find the value of the expression

$\cos^{-1}\left(-\frac{1}{2}\right) - \sin^{-1}\left(-\frac{1}{2}\right)$  in terms of  $\pi$

d) In the diagram below,  $\angle AXY = \angle AYZ$  and  $XP = YP$



(i) Copy this diagram on your sheet.

(ii) Prove that  $\triangle ABY \cong \triangle ACX$ , giving reasons.

(iii) Hence prove that  $\triangle BPC$  is isosceles.

3

2

2

1

4

2

**Question 3** – (14 marks) – Start a new page

Marks


a) (i) If  $\theta = \tan^{-1} A + \tan^{-1} B$  show that  $\tan \theta = \frac{A+B}{1-AB}$

1

(ii) Hence solve the equation  $\tan^{-1} 3x + \tan^{-1} 2x = \frac{\pi}{4}$

4

b)



The diagram shows triangle  $ABC$ . The bisector of angle  $B$  meets the line  $AE$  at  $F$  and the line  $AC$  at  $D$ .

If  $\angle ABD = x^\circ$   
 $\angle AEC = y^\circ$   
 $\angle ACB = z^\circ$

and  $AD = AF$  show that

(i)  $\angle ADF = \frac{1}{2}(y+z)$

3

(ii)  $\angle EAB = y - 2x$

2


c) Find the equation of the tangent to the curve  $y = 3 \cos^{-1} \frac{x}{2}$  at the point on the curve where  $x=0$

4

**Question 4** – (14 marks) – Start a new page

Marks

- a) In the diagram  $AE$  bisects  $\angle BAC$ ,  $BF$  is perpendicular to  $AE$  and  $G$  is the midpoint of  $BC$ .  $BF$  meets  $AC$  at the point  $D$ .




- |   |   |
|---|---|
| (i) Copy this diagram onto your answer sheet and mark in all the given information. | 1 |
| (ii) Prove that $\triangle BAF$ is congruent to $\triangle DAF$                     | 3 |
| (iii) Explain why $BF = FD$   | 1 |
| (iv) Hence prove that $FG$ is parallel to $DC$                                      | 4 |
- b) (i) Find  $\frac{d}{dx}(\sqrt{1-x^2} + x \sin^{-1} x)$  3
- (ii) Hence or otherwise evaluate  $\int_0^{\frac{1}{2}} \sin^{-1} x \, dx$  correct to 3 significant figures. 2

**Question 5** – (14 marks) – Start a new page

Marks

- a)



$ABCD$  is a quadrilateral. If  $W$  is the midpoint of  $AB$ ,  $X$  is the midpoint of  $BC$ ,  $Y$  is the midpoint of  $CD$  and  $Z$  is the midpoint of  $AD$ , prove that the quadrilateral  $WXYZ$  is a parallelogram.

- |   |   |
|---|---|
| b) Consider the function $f(x) = \frac{8}{4+x^2}$   | 4 |
| (i) Show that $f(x)$ is an even function, and the $x$ axis is a horizontal asymptote to the curve $y = f(x)$      | 3 |
| (ii) Find the coordinates and nature of the stationary point on the curve $y = f(x)$                              | 3 |
| (iii) Sketch the graph of the curve showing the above features.   | 2 |
| (iv) Find the exact area of the region in the first quadrant bounded by the curve $y = f(x)$ and the line $x = 2$ | 2 |

Q3 Michelle

Question 1

a)  $\frac{dy}{dx} = v \frac{dv}{dx} + u \frac{du}{dx}$   
 $u = x$      $v = \tan^{-1} \frac{2x}{2}$   
 $\frac{du}{dx} = 1$      $\frac{dv}{dx} = \frac{2}{4-x^2}$   
 $\frac{dy}{dx} = \tan^{-1} \frac{2x}{2} + \frac{2x}{4-x^2}$

b) (i) For  $\triangle AXZ$  and  $\triangle ABC$   
 $\angle A$  is common  
 $\angle AXZ = \angle ABC$  (corresponding angles on  $XZ \parallel BC$ )  
 $\therefore \triangle AXZ \sim \triangle ABC$  (equiangular)

(ii)  $XY : BC = 8 : 14$  sides of similar triangles  
 $= 4 : 7$


$AY : AC = 4 : 7$

$$\frac{AY}{AC} = \frac{4}{7}$$

$$\frac{AY}{18} = \frac{4}{7}$$

$$AY = \frac{72}{7} \text{ cm}$$

c) (i) Domain  $-2 \leq x \leq 2$  range  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$



Question 2

a) Let  $\angle AED = x$

$\therefore \angle AEB = x$

Let  $\angle CEF = y$

$\therefore \angle BEF = y$

$x + x + y + y = 180^\circ$  angles on a straight line

$$2x + 2y = 180$$


$$x + y = 90$$

$$\angle AEF = x + y = 90^\circ$$

b)  $\int_0^1 \frac{1}{\sqrt{4-3x^2}} dx = \left[ \frac{1}{\sqrt{3}} \sin^{-1} \frac{\sqrt{3}x}{2} \right]$   
 $= \frac{1}{\sqrt{3}} \left( \frac{\pi}{3} - 0 \right)$   
 $= \frac{\pi}{3\sqrt{3}}$

c)  $\cos^{-1} \left( -\frac{1}{2} \right) = \frac{2\pi}{3}$   
 $\sin^{-1} \left( \frac{1}{2} \right) = \frac{\pi}{6}$

$\cos^{-1} \left( -\frac{1}{2} \right) - \sin^{-1} \left( \frac{1}{2} \right) = \frac{2\pi}{3} - \frac{\pi}{6}$   
 $= \frac{5\pi}{6}$



(ii) For  $\triangle ABY$  and  $\triangle ACX$   
 $\angle PXY = \angle PYX$  angles opposite equal sides  
 $\therefore \angle AXC = \angle AYB$  sums of equal angles

$\angle A$  is common

$AX = AY$  sides opposite equal angles

$\therefore \triangle ABY \cong \triangle ACX$  (AAS)

(iii)  $BY = CX$  (corresponding sides of congruent  $\triangle$ 's)

$BY - PY = CX - PX$

$BP = CP$

$\therefore \triangle BPC$  is an isosceles  $\triangle$

Question 3

(1) a) (i)  $\tan \theta = \tan(\tan^{-1} A + \tan^{-1} B)$   
 $= \frac{\tan(\tan^{-1} A) + \tan(\tan^{-1} B)}{1 - \tan(\tan^{-1} A)\tan(\tan^{-1} B)}$   
 $= \frac{A + B}{1 - AB}$

(4) (ii)  $A = 3x$      $B = 2x$   
 $\tan^{-1} 3x + \tan^{-1} 2x = \frac{3x + 2x}{1 - 3x \cdot 2x}$   
 $= \frac{5x}{1 - 6x^2}$   
 $\frac{5x}{1 - 6x^2} = \tan \frac{\pi}{4}$   
 $\frac{5x}{1 - 6x^2} = 1$

$$5x = 1 - 6x^2$$

$$6x^2 + 5x - 1 = 0$$

$$6x^2 + 6x - x - 1 = 0$$

$$6x(x+1) - (x+1) = 0$$

$$(x+1)(6x-1) = 0$$

$x = -1$  or  $\frac{1}{6}$   
 NO SOLUTION

(3) b) (i)  $\angle AFD = \angle ADF$  angles opposite equal sides

$$\angle EAC = 180 - (y + z)$$

$$\angle AFD + \angle ADF = y + z$$

$$\therefore \angle ADF = \frac{y+z}{2}$$

(2) (ii)  $\angle ABE + \angle BAE = \angle AEC$  (exterior angle of a triangle)

$$2x + \angle AEB = y$$

$$\angle AEB = y - 2x$$


(4) (i)  $\frac{dy}{dx} = \frac{-3}{\sqrt{4-x^2}}$   
 $= \frac{-3}{2}$  when  $x = 0$

$$\left( 0, \frac{3\pi}{2} \right)$$

$$y - \frac{3\pi}{2} = -\frac{3}{2}x$$

$$3x + 2y - 3\pi = 0$$

Question 4



(ii) For  $\triangle ABF$  and  $\triangle ACF$   
 $\angle BAF = \angle CAF$  given  
 $\angle BFA = \angle CFA$  given as perpendicular

$AF$  is common

$\therefore \triangle ABF \cong \triangle ACF$  (AAS)

(iii)  $BF = CF$  corresponding sides of congruent triangles

(iv) For  $\triangle BEF$  and  $\triangle CDF$

$BF = CF$  above

$\angle B = \angle C$  given

$\angle E = \angle F$  is common

$\therefore \triangle BEF \sim \triangle CDF$  two sides in proportion and the included angle equal


$\therefore \angle BGF = \angle CDF$  corresponding angles of similar triangles

$\therefore EF \parallel DC$  (corresponding angles are equal on parallel lines)

b) (i)  $\frac{d}{dx} (\sqrt{1-x^2} + x \sin^{-1} x)$   
 $= \frac{-2x}{2\sqrt{1-x^2}} + \frac{x}{\sqrt{1-x^2}} + \sin^{-1} x$   
 $= \frac{-x}{\sqrt{1-x^2}} + \frac{x}{\sqrt{1-x^2}} + \sin^{-1} x$   
 $= \sin^{-1} x$

(ii)  $\int_0^{\frac{1}{2}} \sin^{-1} x dx$   
 $= \left[ \sqrt{1-x^2} + x \sin^{-1} x \right]_0^{\frac{1}{2}}$   
 $= \left[ \sqrt{1-\frac{1}{4}} + \frac{1}{2} \sin^{-1} \frac{1}{2} - 1 - 0 \right]$   
 $= \frac{\sqrt{3}}{2} + \frac{\pi}{12} - 1$   
 $= 0.128$

### Question 5



join BD

For  $\triangle AWZ$  and  $\triangle ABD$

$AZ = ZD$   
 $AW = WB$   
 $\angle A$  is common

$\therefore \triangle AWZ \parallel \triangle ABD$

two sides in proportion  
 and the included angles equal

$\therefore WZ \parallel BD$  equal intercepts

For  $\triangle CXY$  and  $\triangle CBD$

$CX = XB$   
 $CY = YD$   
 $\angle C$  is common

$\therefore \triangle CXY \parallel \triangle CBD$

two sides in proportion  
 and the included angles equal

$\therefore XY \parallel BD$  equal intercepts

$\therefore WZ \parallel XY$

join AC

and use similar proof of

$WX \parallel ZY$


$\therefore WXYZ$  is a parallelogram

b) (i)  $f(-x) = \frac{8}{4+(-x)^2} = \frac{8}{4+x^2} = f(x)$   
 $\therefore$  it is an even function

(ii)  $f'(x) = \frac{-16x}{(4+x^2)^2}$   
 $f'(x) = 0$  when  $x = 0$   
 $f(0) = 2$

0	0	0
red	0	red

$\therefore$  a maximum turning pt at  $(0, 2)$



(iv)  $\int_0^{\pi} \frac{8}{4+x^2} dx = \left[ 4 \tan^{-1} \frac{x}{2} \right]_0^{\pi}$   
 $= 4 \tan^{-1} 1 - 4 \tan^{-1} 0$   
 $= \pi - 0$   
 $= \pi$