



Mathematics Extension 1

*Time Allowed: 75 Minutes
(plus 5 minutes reading time)*

Instructions to Candidates

1. Write using black or blue pen.
2. Attempt all questions.
3. Start each question on a new page.
4. Show all necessary working.
5. Marks for each question are shown in the right column.
6. Complete cover sheet clearly showing:
 - your name
 - your mathematics class and teacher.

Question 1 – (14 marks) – Start a New Page

- a) Differentiate with respect to x


(i) $y = \log_e(7x + 6)$

(ii) $y = e^{x^2+2}$

(iii) $y = (e^{2x} + 3)^5$

(iv) $y = x^3 e^{2x}$

(v) $y = \frac{\log_e x}{x}$



The graphs of $y = x^2 - x - 6$ and $y = x + 2$ are shown on the diagram.

- (i) Find the coordinates of P and Q .

- (ii) Find the shaded area.

Question 2 – (14 marks) – Start a new page

a) If $\log_a 2 = 1.31$ and $\log_a 3 = 2.07$ find the value of:

(i) $\log_a 6$

(ii) $\log_a 4.5$

(iii) $\log_a \frac{8}{a^2}$

Marks

1

1


2

b) Solve the equation $3^x = 17$, giving your answer correct to 3 decimal places.

2

c) Find the equation of the tangent to the curve $y = \log_e(3x-2) + 4$ at the point where $x=1$

3



The curves $y = x^2 + 1$ and $y = \frac{2}{x}$ meet at the point A, as shown on the diagram.

(i) Show that A has coordinates (1, 2)

1

(ii) Find the value of k , given that the shaded area is $\frac{10}{3}$ units²

4

Question 3 – (14 marks) – Start a new page

a) Find the following:


(i) $\int x\sqrt{x} dx$

(ii) $\int \frac{1}{3x^2} dx$

(iii) $\int x e^{x^2+1} dx$

b) Show that $\int_{-6}^{-2} \frac{1}{x} dx = -\log_e 3$

c) Use Simpson's Rule with 3 function values to find an approximate value of $\int_0^2 \frac{1}{\sqrt{4+x^2}} dx$ correct to 3 decimal places.



$y = x^2 e^{-x}$ has a maximum turning point at A, as shown on the graph.

(i) Find the coordinates of A.

(ii) The equation $x^2 e^{-x} - k = 0$ has 3 real, distinct roots. Using the graph, or otherwise, write down the possible values of k .

Question 4 – (14 marks) – Start a new page

Marks

a) Solve $\log_3(x+3) + \log_3(x-5) = 2$

4

b) The curve $y = f(x)$ has a stationary point at the point $(2, 2)$.

If $\frac{d^2y}{dx^2} = \frac{2}{x^2}$ find $f(x)$

4

c) The section of the curve $y = \frac{x}{\sqrt{x^3+1}}$ from $x=1$ to $x=3$ is rotated about the x axis. Find the volume of the solid of revolution.

3

d) (i) Show that $f(x) = \frac{x^3}{x^2+1}$ is an odd function.

2

(ii) Hence, or otherwise, write down the value of $\int_{-2}^2 \frac{x^3}{x^2+1} dx$


1

Question 5 – (14 marks) – Start a new page

Mark

a) Evaluate $\int_0^2 (3-2x)^3 dx$

b)



A rectangular prism has dimension x , $3x$ and h metres.

(i) If the surface area is $72m^2$ show that $h = \frac{36-3x^2}{4x}$

(ii) Hence show that the maximum volume of this prism is $36m^3$

c) (i) Differentiate $y = (x+2)\sqrt{2x+1}$. Express your answer as a single fraction.

(ii) Hence or otherwise find $\int_4^{12} \frac{x+1}{\sqrt{2x+1}} dx$

Extension 1 Solutions

HSC Assessment Task I

Question 1

a) (i) $y = \log_e(7x+6)$

$$\frac{dy}{dx} = \frac{7}{7x+6}$$

(ii) $y = e^{x^2+2}$

$$\frac{dy}{dx} = 2x \cdot e^{x^2+2}$$

(iii) $y = (e^{2x} + 3)^5$

$$\frac{dy}{dx} = 5(e^{2x} + 3)^4 \cdot 2e^{2x}$$

$$= 10e^{2x}(e^{2x} + 3)^4$$

(iv) $y = x^3 e^{2x}$

$$\frac{dy}{dx} = e^{2x}(3x^2) + x^3(2e^{2x})$$

(v) $y = \frac{\log_e x}{x}$

$$\frac{dy}{dx} = x \left(\frac{1}{x} \right) - \log_e x \cdot (1)$$

$$= \frac{1 - \log_e x}{x^2}$$

b) (i) $x^2 - x - 6 = x + 2$

$$x^2 - 2x - 8 = 0$$

$$(x+2)(x-4) = 0$$

$$x = 2, 4$$

when $x=2, y=0$

when $x=4, y=6$

P is (4, 6) and Q is (-2, 0)

(ii) $A = \int_{-2}^4 x+2 - (x^2-x-6) dx$

$$= \int_{-2}^4 (2x+8-x^2) dx$$

$$= \left[x^2 + 8x - \frac{x^3}{3} \right]_{-2}^4$$

$$= (16 + 32 - \frac{64}{3}) - (4 - 16 + \frac{8}{3})$$

$$= 36 \text{ units}^2$$

Question 2

a) $\log_a 2 = 1.31$ $\log_a 3 = 2.07$

$$\begin{aligned} \text{(i)} \log_a b &= \log_a (2 \times 3) \\ &= \log_a 2 + \log_a 3 \\ &= 1.31 + 2.07 \\ &= 3.38 \end{aligned}$$

(ii) $\log_a 4.5 = \log_a \left(\frac{3^2}{2}\right)$

$$\begin{aligned} &= 2\log_a 3 = \log_a 8 \\ &= 2 \times 2.07 - 1.31 \\ &= 2.83 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \log_a 8 &= \log_a 2^3 - \log_a 2 \\ &= 3\log_a 2 = 2 \\ &= 3 \times 1.31 - 2 \\ &= 1.93 \end{aligned}$$

b.) $3^x = 17$

$$\log_e 3^x = \log_e 17$$

$$x = \frac{\log_e 17}{\log_e 3}$$

c) $y = \log_e(3x-2) + 4 \quad x=1$

$$\frac{dy}{dx} = \frac{3}{3x-2}$$

when $x=1, \frac{dy}{dx} = \frac{3}{1}$

when $x=1, y = \log_e 1 + 4$

$$= 4$$

$$m=1 \quad (1, 4)$$

$$y-4 = 1(x-1)$$

equation of the tangent

$$is \quad y = x + 3$$

d) (i) Sub. (1, 2) in $y = x^2 + 1$,

$$\text{RHS} = 2$$

$$\text{LHS}$$

$$\text{Sub. (1, 2) in } y = \frac{2}{x}$$

$$\text{RHS} = 2$$

$$= \text{LHS}$$

A has coordinates (1, 2)

since it satisfies both equations.

(ii) $\frac{16}{3} = \int_0^1 x^2 + 1 dx + \int_1^K \frac{2}{x} dx$ (c.) $\int_0^2 \frac{1}{\sqrt{4+x^2}} dx$

$$= \left[\frac{x^3}{3} + x \right]_0^1 + \left[2\log_e x \right]_1^K$$

$$= \left(\frac{1}{3} + 1 - 0 \right) + (2\log_e K - 2\log_e 1)$$

$$= \frac{4}{3} + 2\log_e K$$

$$2\log_e K = 2$$

$$\log_e K = 1$$

$$K = e$$

Questions 3

a) (i) $\int x \sqrt{x} dx = \int x^{\frac{3}{2}} dx$

$$= \frac{2x^{\frac{5}{2}}}{5} + C$$

(ii) $\int \frac{1}{3x^2} dx = \int \frac{1}{3} x^{-2} dx$

$$= \frac{1}{3} x^{-1} + C$$

$$= \frac{-1}{3x} + C$$

(iii) $\int x \cdot e^{x^2+1} dx = \frac{1}{2} e^{x^2+1}$

$$= \frac{1}{2} e^{x^2+1}$$

$$= (\log_e 2 + 1)$$

$$= \log_e \left(\frac{e}{2} \right)$$

$$= \log_e \left(\frac{1}{2} \right)$$

$$= \log_e 3^1$$

$$= -\log_e 3$$

$$\int_0^2 \frac{1}{\sqrt{4+x^2}} dx \approx \frac{2}{6} [f(0) + 4f(1) + f(2)]$$

$$= \frac{1}{3} [0.5 + 4 \times 0.447 + 0.354]$$

$$x^2 = 4$$

$$x = 2, x > 0$$

$$\frac{d^2V}{dx^2} = -\frac{27x}{2}$$

$$\text{when } x=2, \frac{d^2V}{dx^2} = -2 < 0$$

∴ max. occurs when $x = 2$, $V = 27 \times 2 - 9$

$$= 36m^3$$

$$(e) (i) y = (x+2)\sqrt{2x+1}$$

$$= (x+2)(2x+1)^{\frac{1}{2}}$$

$$\begin{aligned} b) (i) A &= 2(3x \times x) + 2(3x \times h) + 2(2x \times h) \\ &= 6x^2 + 6xh + 2xh \\ &= 6x^2 + 8xh \\ 72 &\equiv 6x^2 + 8xh \\ 8xh &= 72 - 6x^2 \\ h &= \frac{72 - 6x^2}{8x} \\ \therefore h &= \frac{36 - 3x^2}{4x} \end{aligned}$$

$$(ii) \int_4^{12} \frac{3x+3}{\sqrt{2x+1}} dx = (x+2)\sqrt{2x+1}$$

$$\begin{aligned} \int_3^4 \frac{3(3x+1)}{\sqrt{2x+1}} dx &= \int_3^4 \frac{x+1}{\sqrt{2x+1}} dx \\ &= \frac{1}{3} (6x+3) \sqrt{2x+1} \Big|_3^4 \\ &= \frac{1}{3} (6(4)+3) \sqrt{2(4)+1} - \frac{1}{3} (6(3)+3) \sqrt{2(3)+1} \\ &= \frac{1}{3} (27) \sqrt{9} - \frac{1}{3} (15) \sqrt{7} \\ &= \frac{1}{3} (14x^2 - 6x^3) \end{aligned}$$

$$= \frac{1}{3} (14 \times 5 - 6 \times 3)$$

$$(iii) \int_1^2 \frac{27x^2}{\sqrt{2x+1}} dx = 17\frac{1}{3}$$

$$d) y = x^2 e^{-x}$$

$$\begin{aligned} (i) \frac{dy}{dx} &= e^{-x}(2x) + x^2(-e^{-x}) \\ &= xe^{-x}(2-x) \end{aligned}$$

stat. pts occur when $\frac{dy}{dx} = 0$.

$$xe^{-x}(2-x) = 0$$

$$x=0, 2$$

$$\text{when } x=2, y = \frac{4}{e^2}$$

∴ A has coordinates $(2, \frac{4}{e^2})$

$$(ii) 0 < k < 2$$

Question 4

$$a) \log_3(x+3) + \log_3(x-5) = 2.$$

$$\log_3((x+3)(x-5)) = 2.$$

$$(x+3)(x-5) = 3^2$$

$$x^2 - 2x - 15 = 9.$$

$$x^2 - 2x - 24 = 0.$$

$$(x+4)(x-6) = 0.$$

$$x = -4, 6.$$

∴ $x = 6$, since $x+3 > 0$

and $x-5 > 0$

$$b) \frac{d^2y}{dx^2} = \frac{2}{x^2} \quad \text{stat. pt. @ (2,2)}$$

$$\frac{dy}{dx} = \int 2x^{-2} dx$$

$$= -2x^{-1} + C$$

$$\text{when } \frac{dy}{dx} = 0, x = 2.$$

$$dy = -2 + C$$

$$0 = -\frac{2}{2} + C_1$$

$$\therefore C_1 = 1$$

$$\frac{dy}{dx} = -\frac{2}{x} + 1$$

$$y = \int \left(-\frac{2}{x} + 1\right) dx$$

$$= -2 \log x + x + C_2$$

$$\text{when } x=2, y=2$$

$$2 = -2 \log 2 + 2 + C_2$$

$$\therefore C_2 = \log 4$$

$$f(x) = x - 2 \log x + \log 4.$$

$$c) y = \frac{x}{\sqrt{x^3+1}} \quad x=1 \text{ to } x=3$$

$$y^2 = \frac{x^2}{x^3+1}$$

$$V = \pi \int_1^3 \frac{x^2}{x^3+1} dx$$

$$= \frac{1}{3} \pi \left[\log(x^3+1) \right]_1^3$$

$$= \frac{\pi}{3} (\log 28 - \log 2)$$

$$= \frac{\pi}{3} \log 14 \text{ units}^3$$

$$d) (i) f(x) = \frac{x^3}{x^2+1}$$

$$f(-x) = \frac{-x^3}{x^2+1}$$

$$-f(x) = -\left(\frac{x^3}{x^2+1}\right)$$

$$= f(-x)$$

$$(ii) \int_2^2 \frac{x^3}{2x^2+1} dx = 0$$

Question 5

$$a) \int_2^2 (3-2x)^3 dx = \left[\frac{(3-2x)^4}{4x-2} \right]_2^2$$

$$= \left[\frac{(3-2x)^4}{-8} \right]_2^2$$

$$= -\frac{1}{8} + \frac{81}{8} \quad (1)$$

$$= 10 \quad (2)$$

$$(e) (ii) y = (x+2)\sqrt{2x+1}$$

$$= (x+2)(2x+1)^{\frac{1}{2}}$$

$$dy = (2x+1)^{\frac{1}{2}} \cdot 1 + (x+2) \cdot \frac{1}{2}(2x+1)^{-\frac{1}{2}} \cdot 2$$

$$= \sqrt{2x+1} + \frac{(x+2)}{\sqrt{2x+1}}$$

$$= \frac{2x+1+x+2}{\sqrt{2x+1}} \quad (3)$$

$$= \frac{3x+3}{\sqrt{2x+1}} \quad (3)$$

$$= \frac{3x+3}{\sqrt{2x+1}} \quad (3)$$

$$(ii) \int_4^{12} \frac{3x+3}{\sqrt{2x+1}} dx = (x+2)\sqrt{2x+1}$$

$$\int_3^4 \frac{3(3x+1)}{\sqrt{2x+1}} dx = \int_3^4 \frac{x+1}{\sqrt{2x+1}} dx$$

$$= \frac{1}{3} (6x+3) \sqrt{2x+1} \Big|_3^4$$

$$= \frac{1}{3} (6(4)+3) \sqrt{2(4)+1} - \frac{1}{3} (6(3)+3) \sqrt{2(3)+1}$$

$$= \frac{1}{3} (27) \sqrt{9} - \frac{1}{3} (15) \sqrt{7}$$

$$= \frac{1}{3} (14x^2 - 6x^3)$$

$$\text{stat. pt occurs when } \frac{dy}{dx} = 0.$$

$$27 = 27x^2$$