

SYDNEY TECHNICAL HIGH SCHOOL

YEAR 12 ASSESSMENT TASK 2

MARCH 2004

MATHEMATICS 2 UNIT

Time allowed: 70 minutes

Instructions:


- Start each question on a new page
- Full marks may not be awarded if working is incomplete or illegible.

Name: _____

Teacher: _____

Q1	Q2	Q3	Q4	Q5	Total
/10	/10	/10	/10	/10	/50

Question 1



- a) Copy this diagram above onto your answer page

- b) Find the length of OA in surd form

1

- c) Find the gradient of OA

1

- d) Find the equation of OA in general form
- e) K is on OA such that BK \perp OA. Use your answer in part (c) to find the gradient of BK and hence the equation of BK.
- f) Find the perpendicular distance BK and hence the area of \triangle OAB.
- g) Find the coordinates of a point C such that OABC is a parallelogram

2

2

3

1

Question 2

- a) Given $\frac{dy}{dt} = 12t^2 + t$ and that $y = 20$ when $t = 2$, find y in terms of t

2


- b) (i) Find $\int(x^2 + \frac{3}{x^2})dx$

2

- (ii) Evaluate $\int_1^3 \frac{1}{\sqrt{x}} dx$

2

- c) Two circles are such that the sum of their radii is constant at 10 cm.



Let the radii of the two circles be x cm and y cm.


- (i) Show that the sum of their areas is given by $A = 2\pi x^2 - 20\pi x + 100\pi$

1

- (ii) Show that the sum of their areas will be a minimum when the radii are equal

3

Question 3



- a) Find the x values of the points of intersection of the graphs of $y = x^2 - 6x$ and $y = 2x$.

1

- b) (i) Find the area between the two graphs above.

3

- c) Find the area between the curve $y = x^3$, the y -axis and the lines $y=1$ and $y=8$

3

Question 4

- a) Find a primitive of $(3x + 2)^4$ 1

b) For the curve $y = 6x^2 - x^3 + 9$:

 - (i) Find the stationary points and determine their nature 4
 - (ii) Determine any point (s) of inflection 2
 - (iii) Sketch the curve over the domain $-3 \leq x \leq 5$ 2
 - (iv) In this domain, determine the maximum value of the function. 1


Question 5

- a) Sketch a curve on a number plane that satisfies all conditions below. 3


$$f(0) = 2$$

When $x < 0$, $f'(x) < 0$ and $f''(x) > 0$

When $x > 0$, $f'(x) < 0$ and $f''(x) < 0$.

- b)  Find the shaded area

$$y = x^2 - 1$$




- c) The graphs of $y = 2 - x$ and $y = \sqrt{4 - x^2}$ are shown.

Find the volume generated when

the shaded area shown

is rotated about the x -axis.



Solutions.

D) b) $OA = \sqrt{6^2 + 10^2}$
 $= \sqrt{136} \quad \textcircled{1}$

c) $M_{AB} = \frac{10}{6}$
 $= \frac{5}{3} \quad \textcircled{1}$

d) $y = \frac{5}{3}x$, $3y = 5x$
 $\therefore 5x - 3y = 0 \quad \textcircled{1}$

e) $M_{OK} = -\frac{3}{5} \quad \textcircled{1}$

\therefore eqn. of BK is
 $y - 4 = -\frac{3}{5}(x - 8)$
 $5y - 20 = -3x + 24$
 $\therefore 3x + 5y - 44 = 0 \quad \textcircled{1}$

f) line $5x - 3y = 0$
point $(8, 4)$

\therefore p.d. = $\frac{\sqrt{5^2 + 3^2}}{\sqrt{5^2 + 3^2}}$
 $= \frac{28}{\sqrt{34}} \quad \textcircled{1}$ correct idea
 \therefore correct applic.

\therefore area $\Delta OAB = \frac{1}{2} \times \sqrt{136} \times \frac{28}{\sqrt{34}}$
 $= 28u^2 \quad \textcircled{1}$

g) $C(2, -6) \quad \textcircled{1}$

② a) $y = 4x^3 + \frac{x^2}{2} + c \quad \textcircled{1}$

$y = 20, x = 2 :$
 $\therefore 20 = 32 + 2 + c \quad (c = -14)$
 $\therefore y = 4x^3 + \frac{x^2}{2} - 14 \quad \textcircled{1}$

b) (i) $\int (x^2 + 3x^{-2}) dx$
 $= \frac{x^3}{3} + 3x^{-1} + c \quad \textcircled{1}$

$= \frac{x^3}{3} - \frac{3}{x} + c \quad \textcircled{1}$

(ii) $\int_1^3 x^{-\frac{1}{2}} dx = \left[\frac{x^{\frac{1}{2}}}{\frac{1}{2}} \right]_1^3 \quad \textcircled{1}$
 $= [2\sqrt{x}]_1^3$
 $= 2\sqrt{3} - 2 \quad \textcircled{1}$

c) (i) $A = \pi x^2 + \pi y^2$
 $\quad (\text{and } x+y=10)$
 $\therefore y = 10-x$

$\therefore A = \pi x^2 + \pi (10-x)^2$
 $= \pi(x^2 + 100 - 20x + x^2) \quad \textcircled{1}$
 $= \pi x^2 + 100\pi - 20\pi x + \pi x^2$
 $= 2\pi x^2 - 20\pi x + 100\pi$

(ii) min. area when $\frac{dA}{dx} = 0$ $\textcircled{1}$

$\frac{dA}{dx} = 4\pi x - 20\pi = 0$
 $4\pi(x-5) = 0$
 $\therefore x = 5 \quad \textcircled{1}$

x	5	5	5
$\frac{dA}{dx}$	-	0	+

 $\textcircled{1} \cancel{x}$

\therefore area must be a minimum when $x = 5$ and $y = 5$, i.e. equal radii.

③ a) $A = 2 \times \int_0^3 (3x^2 + 2) dx$
 $= 2[x^3 + 2x]_0^3 \quad \textcircled{1}$
 $= 2[27+6-(0+0)]$
 $= 66u^2 \quad \textcircled{1}$

b) (i) At intersection,
 $x^2 - 6x = 2x$
 $x^2 - 8x = 0$
 $x(x-8) = 0$
 $\therefore x = 0 \text{ or } 8 \quad \textcircled{1}$


(ii) $A = \left| \int_0^8 (x^2 - 6x - 2x) dx \right|$
 $= \left| \int_0^8 (x^2 - 8x) dx \right|$
 $= \left| \left[\frac{x^3}{3} - 4x^2 \right]_0^8 \right| \quad \textcircled{1}$

$= \left| \left(\frac{512}{3} - 256 \right) - (0 - 0) \right|$

$= |170\frac{2}{3} - 256|$

$= |-85\frac{1}{3}|$

$= 85\frac{1}{3}u^2 \quad \textcircled{1}$



c) $y = x^{1/3}$
 $x^3 = y$
 $\therefore x = y^{1/3}$

$A = \int_1^8 y^{1/3} dy \quad \textcircled{1}$
 $= \left[\frac{3}{4} y^{4/3} \right]_1^8 \quad \textcircled{1}$
 $= \frac{3}{4} (8^{4/3} - 1^{4/3})$
 $= \frac{3}{4} (16 - 1)$
 $= \frac{3}{4} \times 15$
 $= 11\frac{1}{4}u^2 \quad \textcircled{1}$

④ a) $\frac{(3x+2)^5}{5 \times 3} + C$
 $= \frac{(3x+2)^5}{15} + C \quad \textcircled{1}$

b) i) S.P.'s when $\frac{dy}{dx} = 0$ ①

$$\frac{dy}{dx} = (2x - 3x^2) = 0$$

$$3x(4-x) = 0$$

$$\therefore x = 0, 4. \quad ①$$

x	0^-	0	0^+
$\frac{dy}{dx}$	$-$	0	$+$

x	4^-	4	4^+
$\frac{dy}{dx}$	$+$	0	$-$

~~+~~ ~~+~~

\therefore min. T.P. at $(0, 9)$ ① \therefore max. T.P. at $(4, 41)$ ①


ii) P. of I. when $\frac{d^2y}{dx^2} = 0$
and concavity changes

$$\frac{d^2y}{dx^2} = (2 - 6x) = 0$$


$$\therefore x = 2 \quad ①$$

x	2^-	2	2^+
$\frac{d^2y}{dx^2}$	$+$	0	$-$

\therefore concavity changes
 \therefore P. of I. at $(2, 25)$



iv) maximum value is 90
or $y = 90 \quad ①$



$$\begin{aligned}
 &= \pi \left[2x^2 - \frac{2x^3}{3} \right]_0^2 \quad ① \\
 &= \pi \left[(8 - 1\frac{2}{3}) - (0 - 0) \right] \\
 &= \pi \times 2^{\frac{2}{3}} \\
 &= 8\pi u^3 \quad ①
 \end{aligned}$$

b) $A = \left| \int_0^1 (x^2 - 1) dx \right| + \int_1^2 (x^2 - 1) dx$ ①

$$\begin{aligned}
 &= \left| \left[\frac{x^3}{3} - x \right]_0^1 \right| + \left[\frac{x^3}{3} - x \right]_1^2 \\
 &= \left| \left(\frac{1}{3} - 1 \right) - (0 - 0) \right| \quad ① \\
 &\quad + \left(\frac{8}{3} - 2 \right) - \left(\frac{1}{3} - 1 \right) \\
 &= \left| -\frac{2}{3} \right| + \frac{2}{3} - \left(-\frac{2}{3} \right) \\
 &= \frac{2}{3} + \frac{2}{3} + \frac{2}{3} \\
 &= 2u^2 \quad ①
 \end{aligned}$$

c) $V = \pi \int_0^2 ((4-x^2)^2 - (2-x)^2) dx$ ①

$$\begin{aligned}
 &= \pi \int_0^2 (4-x^2 - (4-4x+x^2)) dx \\
 &= \pi \int_0^2 (4-x^2 - 4+4x-x^2) dx \\
 &= \pi \int_0^2 (4x-2x^2) dx \quad ①
 \end{aligned}$$