

Name: ..... Maths Teacher: .....

Answers to be done on the multiple choice answer sheet in your answer booklet.

## SYDNEY TECHNICAL HIGH SCHOOL



Year 11

## Mathematics

Assessment 2

JULY, 2015

*Time allowed: 90 minutes*○ **General Instructions:**

- Marks for each question are indicated on the question.
- Approved calculators may be used
- All necessary working should be shown
- Full marks may not be awarded for careless work or illegible writing
- *Begin each question on a new page*
- Write using black or blue pen
- All answers are to be in the writing booklet provided

- Section I Multiple Choice
  - Questions 1-5
  - 5 Marks
- Section II Questions 6-13
  - 63 Marks

1. What are the solutions of  $2x^2 - 5x - 1 = 0$ ?

- (A)  $x = \frac{-5 \pm \sqrt{17}}{4}$
- (B)  $x = \frac{5 \pm \sqrt{17}}{4}$
- (C)  $x = \frac{-5 \pm \sqrt{33}}{4}$
- (D)  $x = \frac{5 \pm \sqrt{33}}{4}$

2. Which inequality defines the domain of the function  $f(x) = \frac{1}{\sqrt{x+3}}$ ?


- (A)  $x > -3$
- (B)  $x \geq -3$
- (C)  $x < -3$
- (D)  $x \leq -3$

3. Find the values of  $m$  for which  $24 + 2m - m^2 \leq 0$ 

- (A)  $m \leq -4$  or  $m \geq 6$
- (B)  $m \leq -6$  or  $m \geq 4$
- (C)  $-4 \leq m \leq 6$
- (D)  $-6 \leq m \leq 4$

Section II


4.



The shaded region is best described by the inequality.

- (A)  $2x - y - 4 \geq 0$
- (B)  $2x - y - 4 \leq 0$
- (C)  $2x - y - 4 > 0$
- (D)  $2x - y - 4 < 0$

5.



If  $AB \parallel CD$ ,  $ED = FD$  and  $\angle DFG = 112^\circ$  then  $\angle BED =$

- (A)  $112^\circ$
- (B)  $24^\circ$
- (C)  $68^\circ$
- (D)  $44^\circ$

Question 6 – (8 marks)

Mark


- a) Evaluate  $\sqrt[3]{\frac{651}{4\pi}}$  to four significant figures 2
- b) Solve  $2 - 3x \leq 8$  and sketch your solution on a number line 2
- c) Solve  $x^2 - 6x = 0$  2
- d) Solve  $4 < 4x - 3 < 9$  2

Question 7 – (8 marks) – Start a new page

- a) Express  $\frac{a^{-1}+b^{-1}}{a+b}$  in simplest fraction form without using negative indices. 2
- b) Solve  $|5x - 2| = |3x + 4|$  2
- c) Solve  $\frac{5}{8}(x+4) = 4x - \frac{1}{2}$  2
- d) Express  $\frac{3\sqrt{2}}{3\sqrt{2}+2\sqrt{3}}$  in the form  $a + b\sqrt{6}$  2

Question 8 – (8 marks) – Start a new page


- a) The diagram shows a line  $\ell_1$  with equation  $3x + 4y - 12 = 0$ , which intersects the y axis at B. A second line  $\ell_2$  with equation  $4x - 3y = 0$ , passes through the origin O and intersects  $\ell_1$  at E.



- (i) Show that coordinates of B are (0, 3). 1
- (ii) Show that  $\ell_1$  is perpendicular to  $\ell_2$ . 2
- (iii) Show that the perpendicular distance from O to  $\ell_1$  is  $\frac{12}{5}$  units. 1
- (iv) Using Pythagoras' theorem, or otherwise, find the length of the interval BE. 1
- (v) Hence, or otherwise, find the area of  $\triangle BOE$ . 1

b) Simplify  $\frac{x^2-1}{x^2-1} + \frac{3x^2+3x+3}{x^2-4x-5}$

**Question 9** – (7 marks) – Start a new page



XY || TP and XP || YT  
 Redraw the diagram in your answer booklet.  
 Find  $x$  giving reasons for your answer.

b) A function is defined as follows


$$f(x) = \begin{cases} 0 & \text{if } x \leq -3 \\ -1 & \text{if } -3 < x < 0 \\ x & \text{if } x \geq 0 \end{cases}$$

Find

- i)  $f(-3)+f(-2)+f(2)$  1
- ii)  $f(a^2)$  1

- c) i) Sketch  $y = |x - 1|$  and  $y = x + 1$  on the same axes. 2  
 Use a ruler and label each function carefully. Show any points of intersection with the  $x$  and  $y$  axes. Your sketch should be approximately half a page.
- ii) Hence solve  $|x - 1| > x + 1$  1

**Question 10** – (8 marks) – Start a new page



Let  $\angle ABP = x$   
 BP bisects  $\angle ABC$  and  $AB \parallel PQ$   
 Redraw this diagram in your answer booklet. Use a ruler.  
 Your diagram should be approximately half a page in size.  
 Prove that  $BQ = PQ$


- b) Find the exact value of 1
  - i)  $\sin 225^\circ$  1
  - ii)  $\tan(-30^\circ)$  1
- c) If  $\theta$  is obtuse and  $\tan \theta = \frac{-1}{5}$  find the exact value of  $\cos \theta$  1
- d) Prove  $\frac{1}{\sin \theta \cdot \cos \theta} \cdot \tan \theta = \cot \theta$  3

**Question 11** – (8 marks) – Start a new page

- a) Solve the following in the domain  $0^\circ \leq x \leq 360^\circ$ .  
(write your answers correct to the nearest minute)
- i)  $\tan 2\theta = -1$  2
  - ii)  $3 \sin^2\theta + 2 \sin\theta = 0$  2
  - iii)  $3 \sin\theta = 2 \cos\theta$  2
- b) Find  $\lim_{x \rightarrow 3} \frac{x-3}{x^2-9}$  2

**Question 12** – (8 marks) – Start a new page

- a) Differentiate the following
- i)  $y = 4x^3 - x + 5$  1
  - ii)  $y = (3x^2 - 4)^4$  2
  - iii)  $y = \frac{x+1}{x-1}$  2




- i) Redraw the diagram showing the size of angles  $\hat{BAC}$ ,  $\hat{ABC}$  and  $\hat{BCA}$ . 1
- ii) Hence find the length of AC to the nearest metre. 2

**Question 13** – (8 Marks) – Start a new page


Mark

- a) In the figure  $CD = 6\text{cm}$ ,  $AC = 10\text{cm}$ , angle  $ACB = 50^\circ$  and angle  $ABC = 90^\circ$ . Find:



- i) AD to the nearest cm 2
  - ii) Area of  $\triangle ACD$  to the nearest  $\text{cm}^2$ . 1
- b) i) Show that  $k = 2$  is a solution to the equation  $2k^3 - 3k^2 - 4 = 0$  1


- ii) The diagram shows a tangent at the point where  $x = k$  (where  $k > 0$ ) to the cubic  $y = x^3$



- $\alpha$ . Find the gradient of the tangent at  $x = k$  1
- $\beta$ . Find the equation of the tangent at  $x = k$   $3x^2 - y - 5 = 0$  2
- $\gamma$ . If the tangent is found to pass through  $(1, -4)$  find the value of  $k$ . 1

Q1	D
2	A
3	A
4	D
5	D

Question 6

- 1)  $3 \cdot 728$  (4 sig. fig.)
- 2)  $2 - 3x \leq 8$   
 $-3x \leq 6$   
 $x \geq -2$
- 
- 3)  $x^2 - 6x = 0$   
 $x(x-6) = 0$   
 $x = 0, x = 6$
- 4)  $4 < 4x - 3 < 9$   
 $7 < 4x < 12$   
 $\frac{7}{4} < x < 3$

Question 7

- 1)  $(\frac{1}{a} + \frac{1}{b}) \div (a+b)$   
 $\frac{(\frac{b+a}{ab}) \times \frac{1}{(a+b)}}{\frac{1}{ab}}$
- 2)  $5x - 2 = 3x + 4$      $5x - 2 = -(3x + 4)$   
 $2x = 6$      $5x - 2 = -3x - 4$   
 $\therefore x = 3$      $8x = -2$   
 and  $x = -\frac{1}{4}$
- 3)  $\frac{5}{8}(x+4) = 4x - \frac{1}{2}$   
 $5(x+4) = 32x - 4$

$$50x + 20 = 32x - 4$$

$$24 = 27x$$

$$x = \frac{24}{27}$$

$$\therefore x = \frac{8}{9}$$

d)  $\frac{3\sqrt{2}}{3\sqrt{2}+2\sqrt{3}} \times \frac{3\sqrt{2}-2\sqrt{3}}{3\sqrt{2}-2\sqrt{3}}$


$$\frac{3\sqrt{2}(3\sqrt{2}-2\sqrt{3})}{18-12}$$

$$\frac{18-6\sqrt{6}}{6}$$

$$\frac{6(3-\sqrt{6})}{6}$$

$$\therefore \frac{3\sqrt{2}}{3\sqrt{2}+2\sqrt{3}} = 3-\sqrt{6}$$


Question 8



i) sub.  $x=0$  into  $l_1: 3x+4y-12=0$   
 $4y = 12$   
 $y = 3$   
 $\therefore B(0, 3)$

ii)  $m_1 = -\frac{3}{4}$      $m_2 = \frac{4}{3}$   
 since  $-\frac{3}{4} \cdot \frac{4}{3} = -1$   
 $\therefore l_1 \perp l_2$

ii)  $p = \left| \frac{3 \cdot 0 + 4 \cdot 0 - 12}{\sqrt{9+16}} \right|$   
 $l_1: 3x+4y-12=0$   
 $p = \frac{-12}{5}$   
 $\therefore p = \frac{12}{5}$  units



iii)  $3^2 - (\frac{12}{5})^2 = (BE)^2$   
 $9 - \frac{144}{25} = (BE)^2$   
 $\frac{81}{25} = (BE)^2$   
 $\therefore BE = \frac{9}{5}$  units


v) Area  $\triangle BOE = \frac{1}{2} \left( \frac{12}{5} \times \frac{9}{5} \right)$   
 $= \frac{54}{25}$  units<sup>2</sup>

b)  $\frac{x^3-1}{x^2-1} \times \frac{x^2-4x-5}{3x^2+3x+3}$

$$\frac{(x-1)(x^2+x+1)}{(x-1)(x+1)} \times \frac{(x-5)(x+1)}{3(x^2+x+1)}$$

$$\frac{x-5}{3}$$


Question 9



a)  $\angle XST = 120^\circ$   
 $\angle YST = 55^\circ$   
 (opposite angles of parallelogram equal)  
 $\therefore x = 65^\circ$  (exterior)


b) i)  $f(-3) + f(-2) + f(2)$   
 $= 0 + -1 + 2$   
 $= 1$

ii)  $f(a^2) = a^2$  since  $a^2 \geq 0$

c) i) 

ii)  $x < 0$


Question 10



a)  $\hat{PBQ} = x$  (BP bisects  $\hat{ABC}$ )  
 $\hat{BPQ} = x$  (alternate angles  $AB \parallel PQ$ )  
 $\therefore PQ = BQ$  (sides opposite equal angles in isosceles triangle)

b) i)  $\sin 225^\circ = \sin (180+45)$   
 $\frac{S}{A}$   
 $\frac{T}{C}$   
 $= -\sin 45^\circ$   
 $= -\frac{1}{\sqrt{2}}$

ii)  $\tan(-30^\circ) = \tan(360-30)$   
 $\frac{S}{A}$   
 $\frac{T}{C}$   
 $= -\tan 30^\circ$   
 $= -\frac{1}{\sqrt{3}}$


c)  $\tan \theta = -\frac{1}{5}$   $\frac{S}{T} | \frac{A}{C}$   
  
 $\therefore \cos \theta = -\frac{5}{\sqrt{26}}$

d) LHS =  $\frac{1}{\sin \theta \cdot \cos \theta} - \tan \theta$   
 $= \frac{1}{\sin \theta \cdot \cos \theta} - \frac{\sin \theta}{\cos \theta}$   
 $= \frac{1 - \sin^2 \theta}{\sin \theta \cdot \cos \theta}$   
 $= \frac{\cos^2 \theta}{\sin \theta \cdot \cos \theta}$   
 $= \frac{\cos \theta}{\sin \theta}$   
 $= \cot \theta$   
 = RHS

b)  $\lim_{x \rightarrow 3} \frac{1(x-3)}{(x^2/3)(x+3)}$   
 $= \frac{1}{6}$


Question 12.

a) i)  $\frac{d}{dx} (4x^3 - x + 6) = 12x^2 - 1$   
 ii)  $\frac{d}{dx} (3x^2 - 4)^4 = 4 \cdot 6x(3x^2 - 4)^3 = 24x(3x^2 - 4)^3$   
 iii) Let  $u = x + 1$   $v = x - 1$   
 $u' = 1$   $v' = 1$   
 $\therefore \frac{d}{dx} \left( \frac{x+1}{x-1} \right) = \frac{1(x-1) - 1(x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$




ii)  $\frac{AC}{\sin 116^\circ} = \frac{270}{\sin 22^\circ}$   
 $\therefore AC = \frac{270 \sin 116^\circ}{\sin 22^\circ}$   
 $AC = 648 \text{ m (nearest m)}$

Question 13



ii) Area  $\Delta ACD = \frac{1}{2} \cdot 6 \cdot 10 \cdot \sin 130^\circ = 23 \text{ cm}^2$  (nearest  $\text{cm}^2$ )  
 b) i) sub  $k = 2$  into  $2k^3 - 3k^2 - 4 = 0$   
 $LHS = 16 - 12 - 4 = 0 = RHS$   
 $\therefore k = 2$  is a solution



ii)  $y = x^3$   
 $\frac{dy}{dx} = 3x^2$   
 $\therefore m = 3k^2$  where  $x = k$

pt.  $(k, k^3)$   
 tangent:  $y - k^3 = 3k^2(x - k)$   
 $y - k^3 = 3xk^2 - 3k^3$   
 $y = 3xk^2 - 2k^3$

8. sub  $(1, -4)$  into tangent  
 $-4 = 3k^2 - 2k^3$   
 $\therefore 2k^3 - 3k^2 - 4 = 0$   
 $\therefore$  from part i)  
 $k = 2$

QUESTION 11

a) i)  $\tan 2\theta = -1$   $\frac{S}{T} | \frac{A}{C}$   
 acute  $2\theta = 45^\circ$   
 $\therefore 2\theta = 135^\circ, 315^\circ, 495^\circ, 675^\circ$   
 $\theta = 67\frac{1}{2}^\circ, 157\frac{1}{2}^\circ, 247\frac{1}{2}^\circ, 337\frac{1}{2}^\circ$   
 or  $67^\circ 30', 157^\circ 30', 247^\circ 30', 337^\circ 30'$

ii)  $3 \sin^2 \theta + 2 \sin \theta = 0$   
 $\sin \theta (3 \sin \theta + 2) = 0$   
 $\sin \theta = 0$   $\sin \theta = -\frac{2}{3}$   
 $\theta = 0^\circ, 180^\circ, 360^\circ$  and  $\theta = 221^\circ 44', 318^\circ 11'$

iii)  $3 \sin \theta = 2 \cos \theta$   
 $\frac{\sin \theta}{\cos \theta} = \frac{2}{3}$   $\frac{S}{T} | \frac{A}{C}$   
 $\tan \theta = \frac{2}{3}$   
 $\therefore \theta = 22^\circ 44', 212^\circ 44'$