

# SYDNEY BOYS' HIGH SCHOOL

MOORE PARK, SURRY HILLS



TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION 2001

# MATHEMATICS

## EXTENSION 1

*Time allowed:* 2 Hours  
(plus five minutes reading time)  
*Examiner:* E. Choy

### DIRECTIONS TO CANDIDATES

- ALL questions may be attempted.
- All necessary working should be shown in every question. Full marks may not be awarded for careless or badly arranged work.
- Approved calculators may be used.
- Start each question on a new answer sheet.
- Additional answer sheets may be obtained from the supervisor upon request.

**NOTE:** This is a trial paper only and does not necessarily reflect the content or format of the final Higher School Certificate examination paper for this subject.

Sydney Boys High Extension 1 Trial 2001

### Question 1. (12 marks)

- (a) Find the acute angle (correct to the nearest minute) between the lines  $3x + 2y = 7$  and  $4x - 3y = 2$ . 2
- (b) Using the expansion of  $\tan(\alpha - \beta)$ , or otherwise, show that  $\tan(-15^\circ) = \sqrt{3} - 2$ . 2
- (c) Find  $\lim_{x \rightarrow 0} \left( \frac{\sin 4x + \tan x}{x} \right)$ . 2
- (d) Differentiate with respect to  $x$ : 2
- (i)  $y = \ln(\cos x)$
- (ii)  $y = \tan^{-1} 3x$
- (e) Solve  $2\cos^2 x + 3\sin x - 3 = 0$ , where  $0 \leq x \leq 2\pi$ . 2
- (f) Find the co-ordinates of the point  $P$  that divides the interval joining the points  $A(-3, 4)$  and  $B(-1, 0)$  externally in the ratio 4:3. 2


**Question 2.** (12 marks)

- (a) Find the general solution of  $\tan x = \sqrt{3}$ . Give your answer in a concise, general form. 2
- (b) How many different 9-letter "words" can be made from the letters of *ISOSCELES*? 2
- (c) Find the domain and range of the function  $y = \sin^{-1}(1 - \sqrt{x})$ . 1
- (d) Evaluate  $\int_0^{\sqrt{3}} \frac{dx}{\sqrt{3-x^2}}$ . 2
- (e) Find all solutions to  $\frac{x}{x^2-1} > 0$ . 2

(f) Given  $AB = AC$ , and that the tangent at  $A$  is parallel to  $PQ$ .


Prove:

- (i)  $AP = AQ$
- (ii)  $BC$  is parallel to the tangent at  $A$ .
- (iii)  $PCBQ$  is a cyclic quadrilateral.




**Question 3.** (12 marks)

- (a) Find the exact area bounded by the curve  $y = \sin^{-1} x$ , the  $x$ -axis, and the ordinate  $x = \frac{1}{2}$  as shown in the diagram. 4



(b)



The elevation of the top of a hill ( $B$ ) from a place  $P$  due east of it is  $43^\circ$ , and from a place  $Q$ , due south of  $P$ , it is  $22^\circ$ . The distance from  $P$  to  $Q$  is 400m. If  $h$  is the height of the hill, show that

$$h^2 = \frac{160000}{\cot^2 22^\circ - \cot^2 43^\circ}$$

- (c) Find  $\int \sec^2 x \cdot \tan^2 x \, dx$  using the substitution  $u = \tan x$ . 4

Question 4. (12 marks)

- (a) The points  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  lie on the parabola  $x^2 = 4ay$ . 8
- (i) Find the co-ordinates of  $A$ , the point of intersection of the tangents to the parabola at  $P$  and  $Q$ .  
(You may use the fact that equation of the tangent to the parabola  $x^2 = 4ay$  at the point  $T(2at, at^2)$  is  $y = tx - at^2$ .)
- (ii) Suppose further that  $A$  lies on the line containing the focal chord which is perpendicular to the axis of the parabola.
- ( $\alpha$ ) Show that  $pq = 1$ .
- ( $\beta$ ) Show that the chord  $PQ$  meets the axis of the parabola on the directrix.



- (b) If  $y = x^3 - 2x^2 + 3$ : 4
- (i) find the equation of the tangent to the curve at  $(2, 3)$ , and
- (ii) find the point at which the tangent meets the curve again.

Question 5. (12 marks)

- (a) Prove by mathematical induction that for positive integral  $n$ ,  $3^{3n} + 2^{n+2}$  is divisible by 5. 4
- (b) By considering the function  $f(x) = x^3 - 7$ , use one step of Newton's method to find a better approximation to  $\sqrt[3]{7}$  than 2. Leave your answer in exact fractional form. 3
- (c) The speed  $v$  m/s of a point moving along the  $x$ -axis is given by  $v^2 = 90 - 12x - 6x^2$ , where  $x$  m is the displacement of the point from the origin. 3
- (i) Prove that the motion is simple harmonic.
- (ii) Find the period, the centre of motion, and the amplitude.
- (d) (i) Prove that  $\cos 2\theta = \frac{1-x^2}{1+x^2}$ , where  $x = \tan \theta$ . 2
- (ii) Use the above result to deduce that  $\tan \frac{\pi}{8} = \sqrt{2} - 1$ .


## Question 6. (12 marks)

- (a) Given  $y = \sin^{-1}(\cos x)$ : 4
- (i) Find  $\frac{dy}{dx}$ .
- (ii) Evaluate  $y = \sin^{-1}(\cos x)$  if  $x = \pi$ .
- (iii) Sketch  $y = \sin^{-1}(\cos x)$  for  $-\pi \leq x \leq \pi$ .
- (b) Whilst playing tennis, Eric serves a ball from a height of 1.8 metres. If he hits the ball in a horizontal direction at a speed of 35 m/s, find (using  $g = 10\text{ms}^{-2}$ ): 6
- (i) How long before the ball hits the ground.
- (ii) How far the ball will travel before bouncing.
- (iii) By how much the ball clears the net, which is 0.95 m high and 14 metres distant.

- (c) (i) Find  $\frac{d}{dx}(xe^x)$ . 2
- (ii) Use the result in Part (i) to evaluate  $\int_0^1 xe^x dx$

## Question 7. (12 marks)

- (a) A street lamp is 8 m high. A small object 2 m away from the lamp falls vertically downward. 6
- (i) Show that when the object has fallen  $y$  metres, the shadow it casts on the horizontal ground is  $\frac{16}{y}$  metres from the base of the lamp.
- (ii) When the object has fallen 6 m, it is travelling at 10 m/s. At what speed is its shadow moving?
- (iii) At what height does the object have the same speed as its shadow?



- (b) A function  $f(x)$  is defined by the rule  $f(x) = (e^x - 1)\ln x$  for  $0 < x \leq 1$ . 6
- (i) Evaluate  $f'(1)$ .
- (ii) Using the fact that  $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$ , show that  $f'(x) \rightarrow -\infty$  as  $x \rightarrow 0$ .
- (iii) Hence or otherwise show that  $f(x)$  has a stationary value, and determine its nature.

1)  $\tan x = \sqrt{3}$

$\therefore x = 180n + \tan^{-1}(\sqrt{3})$   
 $\left[ x = 180n + 60 \right]$  or  $\left[ x = n\pi + \frac{\pi}{3} \right]$

repetition of S (x3)  
 repetition of E (x2)

30240

domain:  $-1 \leq 1 - \sqrt{x} \leq 1$

$\therefore -2 \leq -\sqrt{x} \leq 0$

$\therefore 0 \leq \sqrt{x} \leq 2$

$\therefore 0 \leq x \leq 4$  (1)

range:  $-\frac{\pi}{2} \leq \sin^{-1} u \leq \frac{\pi}{2}$


$\therefore \left[ -\frac{\pi}{2} \leq y \leq \frac{\pi}{2} \right]$  (2)

$\int_0^{\sqrt{3}} \frac{dx}{\sqrt{3-x^2}} = \sin^{-1} \left( \frac{x}{\sqrt{3}} \right) \Big|_0^{\sqrt{3}}$   
 $= \sin^{-1}(1) - \sin^{-1}(0)$   
 $= \frac{\pi}{2}$

1)  $\frac{x}{x^2-1} > 0$   $\left[ x > 1 \right]$

$\therefore \frac{x}{(x+1)(x-1)} > 0$   
 $\left[ x > 1 \right]$

$\therefore (x+1)(x-1) > 0$   
 $\therefore \left[ -1 < x < 0, x > 1 \right]$



(f) i)  $\therefore AB = AC$  (base angle of isos.  $\Delta$ )

$\therefore \angle ABC = \angle ACB$  (alternate angles theorem)

$\therefore \angle ACP = \angle AQP$  (alternate angles)

$\therefore \angle AQP = \angle ACB$


$\therefore PQ \parallel BC$  (corresponding angles equal)

$\therefore \angle APQ = \angle AQP$

$\therefore AP = AQ$  (isos.  $\Delta$ )

(ii)  $BC \parallel PQ$  &  $PQ \parallel AX$   
 $\therefore BC \parallel AX$

(iii)  $\angle APQ = \angle ACB$  (from (i))  
 $\therefore$  exterior angle equals opposite interior angle  
 $\therefore PQCB$  is cyclic quad.



3 UNIT: 2001 TRIAL

Q1 a)  $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$   
 gradient  $m_1 = \frac{2}{3}$ ,  $m_2 = \frac{4}{3}$   
 $\tan \theta = \frac{\frac{2}{3} - \frac{4}{3}}{1 + \frac{2}{3} \cdot \frac{4}{3}} = \frac{-\frac{2}{3}}{1 + \frac{8}{9}} = \frac{-\frac{2}{3}}{\frac{17}{9}} = -\frac{2}{3} \cdot \frac{9}{17} = -\frac{6}{17}$   
 $\theta = 70.54^\circ$


(e)  $2(1 - \sin x) + 3 \sin x - 3 = 0$   
 $2 - 2 \sin x + 3 \sin x - 3 = 0$   
 $-2 \sin x + 3 \sin x + 1 = 0$   
 $(2 \sin x - 1)(\sin x - 1) = 0$   
 $\sin x = \frac{1}{2}$  or  $\sin x = 1$   
 $x = \frac{\pi}{6}, \frac{5\pi}{6}$  or  $\frac{\pi}{2}$

(b)  $\tan(30-45) = \frac{\tan 30 - \tan 45}{1 + \tan 30 \tan 45}$   
 $= \frac{\frac{1}{\sqrt{3}} - 1}{1 + \frac{1}{\sqrt{3}} \cdot 1} = \frac{\frac{1 - \sqrt{3}}{\sqrt{3}}}{\frac{\sqrt{3} + 1}{\sqrt{3}}} = \frac{1 - \sqrt{3}}{\sqrt{3} + 1} \cdot \frac{\sqrt{3} - 1}{\sqrt{3} - 1} = \frac{(1 - \sqrt{3})(\sqrt{3} - 1)}{3 - 1} = \frac{1 - \sqrt{3} - \sqrt{3} + 3}{2} = \frac{4 - 2\sqrt{3}}{2} = 2 - \sqrt{3}$

(c)  $\lim_{x \rightarrow 0} \frac{\sin x}{x} + \lim_{x \rightarrow 0} \frac{\cos x}{x}$   
 $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ ,  $\lim_{x \rightarrow 0} \frac{\cos x}{x} = \infty$   
 $\therefore \infty$

(d) (i)  $y = \ln(\cos x)$   
 $\frac{dy}{dx} = \frac{1}{\cos x} \cdot (-\sin x) = -\tan x$

(ii)  $y = \tan^{-1} 3x$   
 $\frac{dy}{dx} = \frac{1}{1 + 9x^2}$



Q4 (a) (i)  $y = px - ap^x$  — (1)  
 $y = qx - aq^x$  — (2)

$0 = (p-q)x - a(p^x - q^x)$   
 $x = \frac{a(p^x - q^x)}{p-q}$   
 $x = a(p+q)$   
 $y = apq$

∴ A is  $(a(p+q), apq)$  3

(ii)  $y = a$  then  $apq = a$   
 $pq = 1$

(iii) check  $y = t(p+q)x - apt^x$

when  $y = ap^x = \frac{ap^x - aq^x}{p-q} = \frac{p+q}{p-q}$

$y - ap^x = (p+q)x - 2ap(p+q)$

$y - ap^x = p+q x - ap^x - apt^x$   
 $y = \frac{p+q}{p-q} x - apt^x$  2


when if  $x = 0$   $y = -apt^x$   
 $y = -a$  because  $pq = 1$  2

-8-8+3

(b)  $y = x^3 - 2x^2 + 3$   
 $\frac{dy}{dx} = 3x^2 - 4x \downarrow \downarrow$   
 $m = 3x + -4x^2 \downarrow$   
 $m = 4$

$\frac{y-3}{x-2} = 4$   
 $y-3 = 4x-8$   
 $y = 4x-5$   
 2

Solve  $x^3 - 2x^2 + 3 = 4x - 5$   
 $x^3 - 2x^2 - 4x + 8 = 0$   
 restrain 2, 2, d  
 $2+2+d = 2$   
 $d = -2$   
 $\therefore (-2, -13)$  2



Now area rectangle is  $\frac{1}{2} \times \frac{\pi}{6} = \frac{\pi}{12}$   
 area required is  $\frac{\pi}{12} - (1 - \frac{\sqrt{3}}{2}) = \frac{\pi}{12} - 1 + \frac{\sqrt{3}}{2}$  2

(b) In  $\Delta BAQ$ ,  $\tan 22^\circ = \frac{h}{PQ}$ ,  $\tan 22^\circ = \frac{h}{PQ}$   
 $PQ = \frac{h}{\tan 22^\circ} = h \cot 22^\circ$   
 In  $\Delta BAP$ ,  $\tan 43^\circ = \frac{h}{AP}$   
 $h = AP \tan 43^\circ$   
 $AP = \frac{h}{\tan 43^\circ} = h \cot 43^\circ$

Now In  $\Delta BPQ$ ,  $AP^2 + PQ^2 = BQ^2$   
 $h^2 \cot^2 43^\circ - h^2 \cot^2 22^\circ = 160000$   
 $h^2 = \frac{160000}{\cot^2 43^\circ - \cot^2 22^\circ} = \frac{160000}{\cot^2 43^\circ - \cot^2 22^\circ}$

3  
 2  
 1

(c)  $\int \sec^2 x \tan x dx$   
 $= \int u^2 du$   
 $= \frac{u^3}{3} + C$   
 $= \frac{\tan^3 x}{3} + C$

Let  $u = \tan x$   
 $\frac{du}{dx} = \sec^2 x$   
 $du = \sec^2 x dx$

4

4

### Question 6

(a)

$$y = \sin^{-1}(\cos x)$$


(i)

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{\sqrt{1-\cos^2 x}} \cdot -\sin x \\ &= \frac{-\sin x}{\sqrt{\sin^2 x}} \\ &= \frac{-\sin x}{|\sin x|} \\ &= -1 \text{ for } 0 < x < \pi \\ &= 1 \text{ for } \pi < x < 2\pi \end{aligned}$$

(ii)  $y = \sin^{-1}[\cos \pi]$

$$\begin{aligned} &= \sin^{-1}[-1] \\ &= -\sin^{-1}[1] \\ &= -\frac{\pi}{2} \end{aligned}$$

(iii)



(b)  $x = v \cos \theta = 35 \cos 0 = 35$   
 $y = 35 \sin 0 - 10t = -10t$   
 $y = 1.8 - 5t^2$   
 $x = vt \cos \theta = 35t$

(i) Strikes ground when  $y=0$   
 $\therefore 0 = 1.8 - 5t^2$   
 $t = 3/5 \text{ sec.}$

(ii)  $x = 35 \times \frac{3}{5} = 21 \text{ m}$

(iii) When  $x=14$ ,  $14 = 35t$   
 $\therefore t = 2/5$

When  $t = 2/5$ ,  $y = 1.8 - 5(2/5)^2$   
 $\therefore y = 1 \text{ m}$

$\therefore$  clears net by  $1 - 0.95 \text{ m} = 5 \text{ cm.}$

(c)

(i)  $\frac{d}{dx}(xe^x) = xe^x + 1 \cdot e^x = e^x(x+1)$

(ii)  $\frac{d}{dx}(xe^x - e^x) = xe^x$

$\therefore \frac{d}{dx}[xe^x - e^x] = xe^x \cdot \frac{1}{2}$

$\Rightarrow \int xe^x dx = (xe^x - e^x) \cdot \frac{1}{2}$

$\therefore \int_0^1 xe^x dx = \left[ \frac{1}{2}(xe^x - e^x) \right]_0^1$   
 $= \frac{1}{2}(e - 0 - (0 - e))$   
 $= 1$

5(a) Start for  $n=1$ ,  $S_1 = 3^2 + 2^2$

$$= 27 + 8$$

$= 35$  which is divisible by 5.

Assume true for  $n=k$ , i.e.  $S_k = 3^{2k} + 2^{2k+1}$

Now let for  $n=k+1$ , i.e.  $S_{k+1} = 3^{2(k+1)} + 2^{2(k+1)+1}$

$$S_{k+1} = 27(3^{2k} - 2^{2k+2}) + 2 \cdot 2^{2k+2}$$

$$= 5 \cdot 27P - (27-2)2^{2k+2}$$

$$= 5 \{ 27P - 5 \cdot 2^{2k+2} \}$$

$= 5Q$  where  $Q \in \mathbb{Z}$ .

$\therefore$  True for  $n=k+1$  if true for  $n=k$ .

Now true for  $n=1$  so true for  $\forall n \in \mathbb{Z}^+$  and so on for all  $n$ .

(b)  $f(x) = x^3 - 7$ ,  $f'(x) = 3x^2$

$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$

$$= 2 - \frac{2-7}{3 \cdot 4}$$

$$= 23/12$$

(c)  $\frac{d^2x}{dt^2} = 45 - 6x - 3x^2$

$$\dot{x} = \frac{dx}{dt} \left( \frac{dx}{dt} \right)$$

$$= -6 - 6x$$

$$= -6(x+1)$$

$$= -6(\sqrt{x})^2$$

$\therefore$  Motion is SHM with centre of motion  $= -1$ .

$$n = \sqrt{6} \text{ so period} = \frac{2\pi}{\sqrt{6}} = \frac{\pi\sqrt{3}}{3}$$

$$v^2 = -6(x^2 + 2x + 1) + 90 + 6$$

$$= 96 - 6(x+1)^2$$

$$= 6 \{ 4^2 - (x+1)^2 \}$$

5(a) (i) RHS =  $1 - \frac{\tan^2 \theta}{\cos^2 \theta}$

$$= \frac{1 + \tan^2 \theta}{\cos^2 \theta}$$

$$= \frac{1 + \frac{\sin^2 \theta}{\cos^2 \theta}}{\cos^2 \theta}$$

$$= \frac{1 + \frac{\sin^2 \theta + \cos^2 \theta}{\cos^2 \theta}}{\cos^2 \theta}$$

$$= \frac{1 + \frac{1}{\cos^2 \theta}}{\cos^2 \theta}$$

$$= \frac{1 + \sec^2 \theta}{\cos^2 \theta}$$

$$= \frac{1 + 1 + \tan^2 \theta}{\cos^2 \theta}$$

$$= \frac{2 + \tan^2 \theta}{\cos^2 \theta}$$

$$= \frac{2 + \frac{\sin^2 \theta}{\cos^2 \theta}}{\cos^2 \theta}$$

$$= \frac{2 \cos^2 \theta + \sin^2 \theta}{\cos^4 \theta}$$

$$= \frac{2 \cos^2 \theta + 1 - \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$


$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$

$$= \frac{1 + \cos^2 \theta}{\cos^4 \theta}$$



(i) Triangles are similar

$$\therefore \frac{x}{2} = \frac{8}{y} \quad 2$$

$$x = \frac{16}{y}$$

(ii)

$$\frac{dx}{dt} = \frac{dx}{dy} \cdot \frac{dy}{dt} \quad 2$$

$$= -\frac{16}{y^2} \cdot 10$$

When  $y = 6$ ,

$$\frac{dx}{dt} = -\frac{16}{36} \cdot 10$$

$$= -\frac{40}{9} \text{ m/s.}$$

(iii)

$$\frac{dy}{dt} = \frac{dx}{dt} \quad \frac{dy}{ds} = 1$$

$$\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt} = \frac{dx}{dt} \quad -\frac{16}{y^2} = -1 \text{ (Spec)}$$

$$y = 4 \cdot 2$$


(b)

(i)  $f(x) = (e^x - 1) \ln x$

$$f'(x) = \frac{e^x - 1}{x} + e^x \ln x \quad 2$$

$$f'(1) = e - 1 + 0 = e - 1$$

(ii) As  $x \rightarrow 0$ ,  $f'(x) \rightarrow \infty$  1  
 since  $\ln x \rightarrow -\infty$  as  $x \rightarrow 0$



$f'(x) = 0$  for  $0 < x < 1$

Let this root be  $\alpha$ .

For  $x < \alpha$   $f'(x) < 0$

$x > \alpha$   $f'(x) > 0$

$\therefore$  the stationary point is a local minimum