

SYDNEY TECHNICAL HIGH SCHOOL



TRIAL HIGHER SCHOOL CERTIFICATE

2003

MATHEMATICS EXTENSION 2

General Instructions:

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Approved calculators may be used
- All necessary working should be shown in every question
- A table of standard integrals is supplied at the back of this paper
- Start each question on a new page
- Attempt all Questions 1 – 8
- All questions are of equal value
- **Total marks 120**

Name: Tommy Lim

Class: _____

| Question 1 | Question 2 | Question 3 | Question 4 | Question 5 | Question 6 | Question 7 | Question 8 | TOTAL |
|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|-------|
| 14 | 15 | 12½ | 15 | 14 | 13 | 15 | 7 | 105 |

106.

Question 1**Marks**

- a) (i) Find $\int \frac{1}{\cos x + 2} dx$ using the substitution $t = \tan \frac{x}{2}$ 3

Evaluate:

(ii) $\int_2^4 \frac{dx}{x^2 - 4x + 8}$ 3

(iii) $\int_{-1}^1 \frac{4+x^2}{4-x^2} dx$ 4

- b) Let n be a positive integer and let

$$I_n = \int_1^2 (\log_e x)^n dx$$

(i) Prove that $I_n = 2(\log_e 2)^n - nI_{n-1}$ 2

(ii) Hence evaluate $\int_1^2 (\log_e x)^4 dx$ as a polynomial in terms of $\log_e 2$ 3

Question 2

- a) The complex number z and its conjugate \bar{z} satisfy the equation $z\bar{z} + 2iz = 12 + 6i$. Find the possible values of z . 4

- b) On an Argand diagram shade the region containing all points representing complex numbers z such that $\operatorname{Re}(z) \leq 1$ and $0 \leq \arg(z+i) \leq \frac{\pi}{4}$ 3

- c) Express $z_1 = \frac{7+4i}{3-2i}$ in the form $a+ib$ where a and b are real. 1

- d) On an Argand diagram sketch the locus of the point representing the complex number z such that $|z - 3 - i| = \sqrt{10}$. Find the greatest value of $|z|$ subject to this condition. 3

- e) (i) Given that w is a complex root of the equation $x^3 = 1$, show that w^2 is also a root of this equation. 2

- (ii) Show that $1+w+w^2=0$, and $1+w^2+w^4=0$. 2

on 3

| Marks | | Marks |
|-------|--|-------|
|-------|--|-------|

The ellipse E has Cartesian equation $\frac{x^2}{25} + \frac{y^2}{16} = 1$

a) Find

(i) the coordinates of the foci S and S'

1

(ii) Show that any point P on E can be represented by the coordinates $(5 \cos \theta, 4 \sin \theta)$ and hence or otherwise prove that $PS + PS'$ is a constant.

3

(iii) Show that the equation of the normal at the point P on the ellipse is

3


$$\frac{5x}{\cos \theta} - \frac{4y}{\sin \theta} = 9$$

(iv) If this normal meets the x axis at M and the y axis at N , prove that

4

$$\frac{PM}{PN} = \frac{16}{25}$$

b) The region shaded below is rotated about the y -axis to form a solid of revolution.



Using the method of cylindrical shells to calculate the volume of this solid, show that:

(i) The volume δV of a shell at x is given by

$$\delta V = 2\pi(4x^3 - x^5)\delta x$$

2

(ii) Hence find the volume of this solid.


2

Question 4

- a) Let $f(x) = -x^2 + 8x - 12$. On separate diagrams, and without calculus, sketch the following graphs. Indicate clearly any asymptotes and intercepts with the axes.
- (i) $y = f(x)$ 2
 - (ii) $y = |f(x)|$ 2
 - (iii) $y^2 = f(x)$ 2
 - (iv) $y = \frac{1}{f(x)}$ 2
 - (v) $y = e^{f(x)}$, giving the coordinates of any turning points by not using calculus. 3
- b) Given $p + q \geq 2\sqrt{pq}$ if p and q are positive real numbers
- (i) Show that $e^a + e^b \geq 2e^{\frac{a+b}{2}}$ for all real a and b 2
 - (ii) Hence find the minimum value of $e^{-2x} + e^{-x} + 1 + e^x + e^{2x}$ for all real x . 2

Question 5

a)



A stone building of height H metres has the shape of a flat topped square 'pyramid' with curved sides as shown in the figure. The cross-section at height h metres is a square with sides parallel to the sides of the base and of length l , $l = \frac{L}{\sqrt{h+1}}$ where L is the side length of the square base in metres.

- (i) Write an expression for the volume of a slice at height h metres. 2
- (ii) Hence find the volume of the building in terms of L and H . 2

■■■■■ 5

- b) The Fibonacci Sequence, F_n , is defined by:

$$F_1 = 1$$

$$F_2 = 1$$

$$F_{n+2} = F_{n+1} + F_n \text{ for all } n \geq 1$$

- (i) Write down the first 12 terms of the sequence

1

- * (ii) Prove, by mathematical induction, that for all positive integers, n , F_{4n} is divisible by 3.

5

- c) Find $\int \frac{\sin^3 \theta}{\cos^4 \theta} d\theta$

3

- * d) Consider the function of $y = \tan^{-1}(\tan x)$

- (i) What is its period?

1

- (ii) Hence sketch the function for $-2\pi \leq x \leq 2\pi$

1

Question 6

The equation $x^3 + 2x - 1 = 0$ has roots α , β , and γ . In each of the following cases, find an equation with integer coefficients having the roots stated below.

- a) (i) $-\alpha, -\beta, -\gamma$ 1

- * (ii) $\alpha, -\alpha, \beta, -\beta, \gamma, -\gamma$ 2

- (iii) $\alpha^2, \beta^2, \gamma^2$ 3


- b) (i) Prove that 1 and -1 are both roots of multiplicity 2 of the polynomial $P(x) = x^6 - 3x^2 + 2$ 2

- (ii) Express $P(x)$ as the product of irreducible factors over the field of

- (α) rational numbers 1

- (β) complex numbers 1

c)



In the Argand diagram above, $\triangle OQR$ is constructed similar to $\triangle OAP$.

Show that

$$(i) \quad |z_3| = |z_1| |z_2|$$

2

$$(ii) \quad \arg z_3 = \arg z_1 + \arg z_2$$


2

(iii) What is the significance of these results? ~~biangular interplay~~

1

Question 7

The figure shows two towns located at B and C . BCD is an equilateral triangle. A road junction is to be placed at P , somewhere on the minor arc BC of the circumscribed circle of the triangle BCD .



Let BP, CP and DP have lengths x , y , z respectively.

The point Q is on the line PC, extended so that \overline{BP} and \overline{CQ} have the same length x . Let $\angle PBC = \alpha$.

7 (cont)

- | | | |
|-------|--------------------------------------------------------------|---|
| (i) | Show that $\angle \text{BPD} = \angle \text{CPD} = 60^\circ$ | 2 |
| (ii) | Find $\angle \text{DCQ}$ in terms of α | 1 |
| (iii) | Prove $\Delta \text{PBD} \equiv \Delta \text{QCD}$. | 2 |
| (iv) | Prove ΔDPQ is equilateral | 2 |
| (v) | Now show that $z = x + y$ | 1 |

- b) Owing to the tides, the depth of water in an estuary may be assumed to rise and fall with time in simple harmonic motion.

At a certain place there is a danger of flooding when the depth of the water is above 1.25m. One day high tide was 1.5m at 1am and the following low tide was 0.5m at 7:30am.

- | | | |
|------|--------------------------------------------------------------------------|---|
| (i) | Find the amplitude in metres and period in minutes of this tidal motion. | 2 |
| (ii) | Hence find between what times after 1am was there no danger of flooding. | 3 |

- c) Find $\int \frac{1-x}{1-\sqrt{x}} dx$

Question 8

- a) (i) Find the 1st and 2nd derivatives of $P(x) = \frac{x^4}{4} + \frac{x^3}{3} + \frac{x^2}{2} + x + c$

- (ii) Hence or otherwise show that $P(x) = 0$ has not real roots if $c > \frac{7}{12}$

- b) (i) Write down in mod-arg form, the five roots of $z^5 - 1 = 0$

- (ii) By combining appropriate pairs of these roots, show that for $z \neq 1$,

$$\frac{z^5 - 1}{z - 1} = (z^2 - 2z \cos \frac{2\pi}{5} + 1)(z^2 - 2z \cos \frac{4\pi}{5} + 1)$$

- (iii) Deduce that $\cos \frac{2\pi}{5}$ and $\cos \frac{4\pi}{5}$ are the roots of the equation

$$4x^2 + 2x - 1 = 0$$

Teacher's Name:

Student's Name/N^o:

Solutions to 2003 S.T.H.S. 4U Trial

$$\text{i. a. (i)} \int \frac{1}{\cos x + 2} dx$$

$$= \int \frac{1}{\frac{1-t^2}{1+t^2} + 2} \times \frac{2dt}{1+t^2} \quad (1)$$

$$= \int \frac{2dt}{1-t^2 + 2(1+t^2)}$$

$$= \int \frac{2dt}{3+t^2} \quad (1)$$

$$= \frac{2}{\sqrt{3}} \tan^{-1} \frac{t}{\sqrt{3}} + C$$

$$= \frac{2}{\sqrt{3}} + \tan^{-1} \left(\frac{\tan \frac{x}{2}}{\sqrt{3}} \right) + C \quad (1)$$

$$\text{(iii)} \quad \frac{4+x^2}{4-x^2} = \frac{8-(4-x^2)}{4-x^2} = \frac{8}{4-x^2} - 1$$

$$\text{Let } \frac{8}{4-x^2} = \frac{A}{2-x} + \frac{B}{2+x}$$

$$\text{Then } 8 = A(2+x) + B(2-x)$$

$$\text{Let } x = -2, \therefore B = 2 \quad (1)$$

$$x = 2, \therefore A = 2 \quad (1)$$

$$\text{(ii)} \quad \int_2^4 \frac{dx}{x^2 - 4x + 8}$$

$$= \int_2^4 \frac{dx}{(x-2)^2 + 4} \quad (1)$$

$$= \left[\frac{1}{2} \tan^{-1} \frac{x-2}{2} \right]_2^4 \quad (1)$$

$$= \frac{1}{2} \tan^{-1} 1 - \frac{1}{2} \tan^{-1} 0$$

$$= \frac{\pi}{8} \quad (1)$$

$\int_{-1}^1 \frac{4+x^2}{4-x^2} dx$ becomes

$$\int_{-1}^1 \frac{2}{2-x} + \frac{2}{2+x} - 1 dx$$

$$= \left[-2 \log_e(2-x) + 2 \log_e(2+x) - x \right]_{-1}^1 \quad (1)$$

$$= -2 \log_e 1 + 2 \log_e 3 - 1 - (-2 \log_e 3 + 2 \log_e 1 + 1)$$

$$= 4 \log_e 3 - 2 \quad (1)$$

$$\text{b (i)} \quad I_n = \int_1^2 (\log_e x)^n dx$$

$$= \int_1^2 (\log_e x)^n \frac{d}{dx} x dx$$

$$= \left[x(\log_e x)^n \right]_1^2 - \int_1^2 x \cdot n(\log_e x)^{n-1} \cdot \frac{1}{x} dx \quad (1)$$

$$= 2(\log_e 2)^n - n I_{n-1} \quad \text{by parts} \quad (1)$$

$$\text{(ii)} \quad \int_1^2 (\log_e x)^4 dx = I_4$$

$$= 2(\log_e 2)^4 - 4I_3$$

$$= 2(\log_e 2)^4 - 4[2(\log_e 2)^3 - 3I_2]$$

$$= 2(\log_e 2)^4 - 8(\log_e 2)^3 +$$

$$12(2(\log_e 2)^2 - 2I_2) \quad (1)$$

$$\text{Now } I_1 = \int_1^2 \log_e x \, dx$$

$$= [x \log_e x - x]_1^2, \text{ by parts } \textcircled{1}$$

$$= 2 \log_e 2 - 1$$

$$\therefore \int_1^2 (\log_e x)^4 \, dx = 2(\log_e 2)^4 - 8(\log_e 2)^3 + 24(\log_e 2)^2 - 48 \log_e 2 + 24 \quad \textcircled{1}$$

$$2. \textcircled{a} \text{ Let } z = x + iy$$

$$\therefore z\bar{z} + 2iz = 12 + 6i \text{ becomes}$$

$$(x+i)(x-i) + 2i(x+iy) = 12 + 6i$$

$$x^2 + y^2 + 2ix - 2y = 12 + 6i$$

$$\therefore x^2 + y^2 - 2y = 12 \text{ and } 2x = 6 \quad \textcircled{1}$$

$$\therefore x^2 + y^2 - 2y = 12 \quad \therefore x = 3 \quad \textcircled{1}$$

$$y^2 - 2y - 3 = 0.$$

$$(y+1)(y-3) = 0$$


$$y = -1 \text{ or } 3$$

$$\therefore z = 3 - i \text{ or } 3 + 3i \quad \textcircled{2}$$

$$\textcircled{b} \quad \operatorname{Re}(z) \leq 1$$

$$0 \leq \arg(z+i) \leq \frac{\pi}{4}$$

$$0 \leq \arg(z - (0-i)) \leq \frac{\pi}{4} \quad \textcircled{1}$$



\textcircled{2}


\textcircled{2}

$$\textcircled{b} \quad z_1 = \frac{7+4i}{3-2i} \times \frac{3+2i}{3+2i}, \quad \textcircled{1}$$

$$= \frac{13+26i}{13} = 1+2i$$

$$\textcircled{d} \quad |z - 3-i| = \sqrt{10}$$

$$\therefore |z - (3+i)| = \sqrt{10} \quad \text{is a circle centre } (3,1) \text{ radius } \sqrt{10} \quad \textcircled{1}$$



(0,0) lies on the circle. Greatest value of $|z| = OP$

$$= 2\sqrt{10} \quad \textcircled{1}$$

Q(i) If ω is a root of $x^3=1$ then ω satisfies the equation

$$\text{ie: } \omega^3 = 1$$

$$\therefore (\omega^3)^2 = \omega^6 = 1 \quad \text{ie: } (\omega^2)^3 = 1$$

ie: ω^2 also satisfies $x^3=1$.

(ii) The sum of the roots of

$$x^3 - 1 = 0 \quad \text{is} \quad -\frac{b}{a} \quad \text{ie: } 0$$

$$\therefore 1 + \omega + \omega^2 = 0.$$

$$\text{Since } \omega^3 = 1,$$

$$\omega^4 = \omega$$

$$\therefore 1 + \omega^4 + \omega^2 = 0.$$

3. Q(i) $a = 5$ $b = 4$

$$e^2 = 1 - \frac{b^2}{a^2}$$

$$= 1 - \frac{16}{25}$$

$$e^2 = \frac{9}{25}$$

$$e = \frac{3}{5}$$


5

(i) $S(ae, 0)$ $S'(-ae, 0)$

$S(3, 0)$ $S'(-3, 0)$

①

6



②

①

①

(iii) Parametric Form of an ellipse

$$(a \cos \theta, b \sin \theta)$$

$$a = 5, b = 4$$

$$\therefore P(5 \cos \theta, 4 \sin \theta) \quad ①$$

$$PS + PS'$$

$eP_{\text{directrix}_1} + eP_{\text{directrix}_2}$

$$e\left(\frac{25}{3} - 5\cos\theta\right) + e\left(5\cos\theta + \frac{25}{3}\right) \stackrel{(1)}{\text{from the}}$$

$$= 2e \times \frac{25}{3}$$

$$= 2 \times \frac{2}{5} \times \frac{25}{3}$$

$$\underline{= 10} \quad (1)$$

$$(iii) \frac{x^2}{25} + \frac{y^2}{16} = 1$$

Differentiating implicitly,

$$\frac{2x}{25} + \frac{2y \frac{dy}{dx}}{16} = 0$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{-2x}{25} \times \frac{16}{2y} \\ &= \frac{-16x}{25y} \end{aligned}$$

$$\text{at } P, \frac{dy}{dx} = M_{\text{tangent}}$$

$$= \frac{-16x \cdot 5\cos\theta}{25 \cdot 4\sin\theta}$$

$$= \frac{-4\cos\theta}{5\sin\theta}$$

$$\therefore M_{\text{normal}} = \frac{5\sin\theta}{4\cos\theta} \quad (1)$$

' Eq'n of normal:

$$y - 4\sin\theta = \frac{5\sin\theta}{4\cos\theta}(x - 5\cos\theta) \quad (1)$$

$$4y\cos\theta - 16\sin\theta\cos\theta = 5\sin\theta x - 25\sin\theta\cos\theta$$

$$9\sin\theta\cos\theta = 5\sin\theta x - 4\cos\theta y$$

$$9 = \frac{5\sin\theta x - 4\cos\theta y}{\sin\theta\cos\theta}$$

$$9 = \frac{5x}{\cos\theta} - \frac{4y}{\sin\theta} \quad (1)$$

(iv) Cuts x axis when $y=0$

$$\text{i.e. } 9 = \frac{5x}{\cos\theta}$$

$$x = \frac{9\cos\theta}{5} \quad (1)$$

Cuts y axis when $x=0$

$$9 = -\frac{4y}{\sin\theta}$$

$$y = -\frac{9\sin\theta}{4} \quad (1)$$

$P(5\cos\theta, 4\sin\theta)$

$M\left(\frac{9\cos\theta}{5}, 0\right)$

$N\left(0, -\frac{9\sin\theta}{4}\right)$

PM
PN



$$\begin{aligned}
 &= \frac{\sqrt{(5\cos\theta - \frac{7}{5}\cos\theta)^2 + (4\sin\theta)^2}}{\sqrt{(5\cos\theta)^2 + (4\sin\theta + (\frac{9}{4}\sin\theta))^2}} \\
 &= \frac{\sqrt{(\frac{16}{5}\cos\theta)^2 + (16\sin^2\theta)}}{\sqrt{25\cos^2\theta + (\frac{25}{4}\sin\theta)^2}} \\
 &= \frac{\sqrt{\frac{256}{25}\cos^2\theta + 16(1-\cos^2\theta)}}{\sqrt{25\cos^2\theta + \frac{625}{16}(1-\cos^2\theta)}} \\
 &= \sqrt{\frac{-5\frac{19}{25}\cos^2\theta + 16}{-14\frac{1}{16}\cos^2\theta + \frac{625}{16}}} \times \frac{400}{400} \quad (1) \\
 &= \sqrt{\frac{-2304\cos^2\theta + 6400}{-5625\cos^2\theta + 15625}} \\
 &= \sqrt{\frac{256(25 - 9\cos^2\theta)}{625(25 - 9\cos^2\theta)}} \\
 &= \cancel{\frac{16}{25}} \quad \text{as req'd.} \quad (1)
 \end{aligned}$$

✓ 3b. (i) $SV = \pi((x + \delta x)^2 - x^2) \cdot h \quad (1)$ ✓
 $= \pi(x^2 + 2x\delta x + \delta x^2 - x^2)(4x^2 - x^4)$
 $= 2\pi(4x^3 - x^5)\delta x \quad \text{since } \delta x^2 \rightarrow 0 \quad (1)$


(ii) $V = \int_0^a 2\pi(4x^3 - x^5)dx$ where a is
where the curve cuts the x -axis
ie: $4x^2 - x^4 = 0$
 $x^2(4 - x^2) = 0$
 $x = 2$

$$\begin{aligned}
 \therefore V &= 2\pi \int_0^2 4x^3 - x^5 dx \quad (1) \\
 &= 2\pi \left[x^4 - \frac{x^6}{6} \right]_0^2 \\
 &= 2\pi \left[16 - \frac{64}{6} \right] \\
 &= \frac{32\pi}{3} \text{ units}^3 \quad (1)
 \end{aligned}$$



$$\begin{aligned} \text{4(i)} \quad f(x) &= -x^2 + 8x - 12 \\ &= -(x^2 - 8x + 12) \\ &= -(x-2)(x-6) \end{aligned}$$



(iv) $y = e^{-x^2+8x-12}$



(iii) $y^2 = f(x)$
 $\Rightarrow y = \pm \sqrt{-x^2 + 8x - 12}$



Q6 (i) If $p = e^a$, $q = e^b$, both real
 $e^a + e^b \geq 2(e^a \times e^b)^{\frac{1}{2}}$ given ①
 $e^a + e^b \geq 2e^{\frac{a+b}{2}}$ ②

$$(a) e^{-2x} + e^{-x} + 1 + e^x + e^{2x} \geq 2e^{\frac{-x+x}{2}} + 1 + 2e^{\frac{-2x+2x}{2}} \quad (b) (i) 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144 \quad (ii) 14$$

$$\geq 2 + 1 + 2$$

$$\geq 5.$$

∴ Min. value is 5 (i)

(ii) Show true for $n=1$.

$$F_4 = 3 \\ 3 = 3$$

(i)

Assume true for $n=k$

5. Consider a slice, of thickness δh , at height h .

$$\text{Area of cross-section} = l^2 = \frac{L^2}{h+1} \text{ m}^2 \quad (i)$$

$$\therefore \text{Volume of slice} = \frac{L^2}{h+1} \delta h \text{ m}^3 \quad (i)$$

$$\begin{aligned} \therefore \text{Volume of solid} &= \int_0^H \frac{L^2}{h+1} dh \quad (i) \\ &= \left[L^2 \log_e(h+1) \right]_0^H \\ &= L^2 \log_e(H+1) \end{aligned}$$

i.e. $F_{4k} = 3K$ where K is a positive integer (i)

Need now to show result holds for $n=k+1$


i.e. $F_{4(k+1)} = 3L$ where L is a positive integer.

$$\begin{aligned} \text{LHS } F_{4k+4} &= F_{4k+3} + F_{4k+2} \text{ from def'n.} \quad (i) \\ &= F_{4k+2} + F_{4k+1} + F_{4k+1} + F_{4k} \\ &= F_{4k+1} + F_{4k} + F_{4k+1} + F_{4k+1} + F_{4k} \\ &= 3F_{4k+1} + 2F_{4k} \quad (i) \\ &= 3F_{4k+1} + 6K \text{ (from assumption)} \\ &= 3[F_{4k+1} + 2K] \\ &= 3L \text{ as } F_{4k+1} + 2K \text{ is integral} \quad (i) \end{aligned}$$

\therefore Since result is true for $n=1$, (15)
 it must also be true for $n=1+1=2$,
 $n=2+1=3$ etc. for all positive integral
 values of n .

$$\begin{aligned} \textcircled{(c)} \quad & \int \frac{\sin^3 \theta}{\cos^4 \theta} d\theta \\ &= \int \frac{\sin^2 \theta}{\cos^4 \theta} \cdot \sin \theta d\theta \\ &= \int \frac{1-\cos^2 \theta}{\cos^4 \theta} \sin \theta d\theta \quad \textcircled{1} \\ &\text{Let } v = \cos \theta \quad \therefore dv = -\sin \theta d\theta \\ &= \int \frac{1-v^2}{v^4} \cdot -dv \quad \textcircled{1} \\ &= \int \frac{1}{v^2} - \frac{1}{v^4} dv \\ &= \frac{-1}{v} + \frac{1}{3v^3} + C \\ &= \frac{1}{3} \sec^3 \theta - \sec \theta + C \quad \textcircled{1} \end{aligned}$$

D) (i) Period is π (same as for $\tan x$) (1)



6. (a) $x^3 + 2x - 1 = 0$ has roots α, β, γ . (1b)

$$\therefore (-x)^3 + 2(-x) - 1 = 0$$

$$\underline{\text{ie: } x^3 + 2x + 1 = 0} \quad \textcircled{1}$$

$$\text{(iii)} \quad (x^3 + 2x + 1)(x^3 + 2x - 1) = 0 \quad \text{from given equation and (i)} \quad \textcircled{1}$$

$$\underline{\text{ie: } x^6 + 4x^4 + 4x^2 - 1 = 0} \quad \textcircled{1}$$

$$(x^{\frac{1}{2}})^3 + 2(x^{\frac{1}{2}}) - 1 = 0$$

$$x^{\frac{3}{2}} + 2x^{\frac{1}{2}} = 1$$

$$x^{\frac{1}{2}}(x+2) = 1 \quad \textcircled{1}$$

$$x(x+2)^2 = 1 \quad \textcircled{1}$$

$$x(x^2 + 4x + 4) = 1$$

$$\underline{x^3 + 4x^2 + 4x - 1 = 0} \quad \textcircled{1}$$

$$\textcircled{(b) (i)} \quad P(x) = x^6 - 3x^2 + 2$$

$$P'(x) = 6x^5 - 6x$$

$$= 6x(x^4 - 1)$$

$$= 6x(x^2 - 1)(x^2 + 1)$$

$$P'(x) = 6x(x-1)(x+1)(x^2 + 1) \quad \textcircled{1}$$

Since $(x-1)$, $(x+1)$ are factors of $P(x)$ then $x=\pm 1$ are roots of multiplicity 2. ①

$$\Rightarrow P(x) = (x-1)^2(x+1)^2(x^2+2)$$

$$(i) P(x) = (x-1)(x+1)(x-1)(x+1)(x^2+2) \quad ①$$

$$(ii) P(x) = (x-1)^2(x+1)^2(x+\sqrt{2}i)(x-\sqrt{2}i) \quad ①$$

③ (i) By similar Δ 's ①

$$\frac{|z_3|}{|z_1|} = \frac{|z_2|}{1}$$

$$\therefore |z_3| = |z_1||z_2| \quad ①$$

$$(ii) \angle ROQ = \angle POA \quad (\text{corr. } \angle's \text{ in similar } \Delta's) \quad ①$$

$$\therefore \arg z_3 - \arg z_2 = \arg z_1$$

$$\therefore \arg z_3 = \arg z_1 + \arg z_2 \quad ①$$

(iii) The construction can be used to multiply complex numbers. ①

$$\begin{aligned} & \text{(i) } \angle CPD = \angle CBD \quad (\text{Angles in same segment}) \\ & \qquad \qquad \qquad = 60^\circ \quad (\text{as } \triangle CBD \text{ is equilateral}) \quad ① \end{aligned}$$

$$\begin{aligned} & \angle BPD = \angle BCD \quad (\text{Angles in same segment}) \\ & \qquad \qquad \qquad = 60^\circ \quad (\text{as } \triangle CBD \text{ is equilateral}) \quad ① \end{aligned}$$

$$(ii) \angle BCQ = \angle CBP + \angle BPC \quad (\text{exterior angle})$$

$$\angle DCQ + 60^\circ = \alpha + 120^\circ \quad \text{from (i)}$$

$$\therefore \underline{\angle DCQ = \alpha + 60^\circ} \quad ①$$

$$(iii) BD = DC \quad (\text{equilateral triangle})$$

$$BP = CQ = x \quad (\text{given}) \quad ①$$

$$\angle DBP = \angle DCQ = \alpha + 60^\circ \quad (\text{from (ii)})$$

$$\therefore \triangle PBD \cong \triangle QCD \quad (\text{SAS}) \quad ①$$

civ) $\angle DPQ = \angle DBC$ (same segment)

$$\therefore \angle DPQ = 60^\circ$$

$$\angle DQC = 180 - \angle DCQ - \angle CDQ$$

$$= 180 - (\alpha + 60) - \angle BDP \quad \textcircled{1}$$

$$= 180 - (\alpha + 60) - (180 - (\alpha + 60) - 60)$$

$$= 180 - \alpha - 60 - (60 - \alpha)$$


$$= 180 - 120$$

$$= 60^\circ \quad \textcircled{1}$$

$\therefore \triangle DPQ$ is equilateral (2 L's of 60°).

v) $Z = x + y$ (equal sides of equilateral \triangle). $\textcircled{1}$

19 7.b.



(i) amplitude = 0.5 m $\textcircled{1}$

$$\text{Period} = 13 \text{ hrs} \times 60$$

$$= 780 \text{ minutes.}$$

(ii) Let equation of motion be

$$x = A \cos(nt) + C$$

$$T = \frac{2\pi}{n}$$

$$780 = \frac{2\pi}{n} \therefore n = \frac{2\pi}{780} = \frac{\pi}{390}$$

$$\therefore x = 0.5 \cos\left(\frac{\pi t}{390}\right) + C$$

$$\text{When } t=0, x=1.5$$

$$\therefore 1.5 = 0.5 \cos 0 + C$$

$$1.5 = 0.5 + C$$

$$\therefore C = 1$$

$\therefore x = 0.5 \cos\left(\frac{\pi t}{390}\right) + 1$ is the equation $\textcircled{1}$ representing the tidal motion.

Need to find t when $x=1.25$

$$1.25 = 0.5 \cos\left(\frac{\pi t}{390}\right) + 1$$

$$\therefore \cos\left(\frac{\pi t}{390}\right) = 0.5 \quad (1)$$

$$\frac{\pi t}{390} = \frac{\pi}{3}, \frac{5\pi}{3}$$

$\therefore t = 130$ minutes, 650 minutes after 1am

no danger of flooding between

3:10 am and 11:50 am (1)

21 8 @ (i) $P'(x) = x^3 + x^2 + x + 1$

$$P''(x) = 3x^2 + 2x + 1 \quad (1)$$

(ii) $P'(x) = 0$

$$x^3 + x^2 + x + 1 = 0$$

$$a=3, \Delta < 0$$

$$x^2(x+1) + 1(x+1)$$

\therefore positive definite

$$(x^2+1)(x+1) = 0$$

when $x = -1$. (1)

\therefore curve is always concave up. (1)

So curve must have a minimum turning point at $x = -1$.

$$\begin{aligned} P(-1) &= \frac{1}{4} - \frac{1}{3} + \frac{1}{2} - 1 + C \\ &= -\frac{7}{12} + C \end{aligned} \quad (1)$$

\therefore if $C > \frac{7}{12}$, curve will always be above the x -axis and have no real roots.

(b) (i) $Z^5 = 1 = \cos 0 + i \sin 0$

$$Z = r(\cos \theta + i \sin \theta)$$

$$\therefore Z^5 = r^5 (\cos 5\theta + i \sin 5\theta) \quad (1)$$

$$\therefore r^5 = 1 \quad \therefore r = 1$$

$$5\theta = 0 + 2k\pi$$

$$\theta = \frac{2k\pi}{5}$$

$$\therefore z^5 = \cos \frac{2k\pi}{5} + i \sin \frac{2k\pi}{5} \quad \text{for } k=0,1,2,3,4 \quad \textcircled{1}$$

When

$$k=0, z_1 = \cos 0 + i \sin 0$$

$$k=1, z_2 = \cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}$$

$$k=2, z_3 = \cos \frac{4\pi}{5} + i \sin \frac{4\pi}{5}$$

$$k=3, z_4 = \cos \frac{6\pi}{5} + i \sin \frac{6\pi}{5} = \bar{z}_3$$

$$k=4, z_5 = \cos \frac{8\pi}{5} + i \sin \frac{8\pi}{5} = \bar{z}_2 \quad \textcircled{1}$$

$$\text{(ii)} z^5 - 1 = (z - z_1)(z - z_2)(z - z_3)(z - z_4)(z - z_5)$$

$$z_1 = 1, z_4 = \bar{z}_3, z_5 = \bar{z}_2 \quad \textcircled{1}$$

$$\therefore z^5 - 1 = (z - 1)(z - z_2)(z - \bar{z}_2)(z - z_3)(z - \bar{z}_3) \quad \textcircled{1}$$

$$\begin{aligned} \therefore \frac{z^5 - 1}{z - 1} &= [z^2 - z(\bar{z}_2 + z_2) + z_2 \bar{z}_2][z^2 - z(z_3 + \bar{z}_3) + z_3 \bar{z}_3] \\ &= [z^2 - z(2\cos \frac{2\pi}{5}) + 1][z^2 - z(2\cos \frac{4\pi}{5}) + 1] \quad \textcircled{1} \\ &= (z^2 - 2z\cos \frac{2\pi}{5} + 1)(z^2 - 2z\cos \frac{4\pi}{5} + 1) \quad \textcircled{1} \end{aligned}$$

23

(iii) For $z^5 = 1$,

Sum of roots is

$$z_1 + z_2 + z_3 + z_4 + z_5 = 0$$

$$z_1 + z_2 + \bar{z}_2 + \bar{z}_3 + z_3 = 0$$

$$1 + 2\cos \frac{2\pi}{5} + 2\cos \frac{4\pi}{5} = 0$$

$$\cos \frac{2\pi}{5} + \cos \frac{4\pi}{5} = -\frac{1}{2}. \quad \textcircled{1}$$

Sum of pairs of roots for $z^5 - 1 = 0$

$$z_1 z_2 + z_1 \bar{z}_2 + z_3 z_1 + z_1 \bar{z}_3 + z_2 \bar{z}_2 + z_2 z_3,$$

$$+ z_2 \bar{z}_3 + \bar{z}_2 z_3 + \bar{z}_2 \bar{z}_3 + z_3 \bar{z}_3 = 0$$

$$\begin{aligned} 1 + 2\cos \frac{2\pi}{5} + 2\cos \frac{4\pi}{5} + 1 + z_2(2\cos \frac{4\pi}{5}) + \bar{z}_2(2\cos \frac{4\pi}{5}) \\ + 1 = 0 \quad \textcircled{1} \end{aligned}$$

$$-1 + 1 + (2\cos \frac{4\pi}{5})(2\cos \frac{2\pi}{5}) + 1 = 0 \quad (\text{from above}),$$

$$\therefore \cos \frac{4\pi}{5} \cos \frac{2\pi}{5} = -\frac{1}{4}. \quad \textcircled{1}$$

∴ The quadratic equation whose roots are $\cos \frac{2\pi}{5}, \cos \frac{4\pi}{5}$ is

$$x^2 - \left(-\frac{1}{2}\right)x + \left(-\frac{1}{4}\right) = 0$$

$$\text{i.e. } 4x^2 + 2x - 1 = 0 \quad \textcircled{1}$$

24